

**SUMMARY OF THE ZETA REGULARIZATION METHOD  
APPLIED TO THE CALCULATION OF DIVERGENT SERIES  $\sum_{n=1}^{\infty} n^s$   
AND DIVERGENT INTEGRALS  $\int_0^{\infty} x^s dx$**

Jose Javier Garcia Moreta

Graduate student of Physics at the UPV/EHU (University of Basque country)  
In Solid State Physics

Address: Practicantes Adan y Grijalba 2 6 G  
P.O 644 48920 Portugalete Vizcaya (Spain)  
Phone: (00) 34 685 77 16 53  
E-mail: josegarc2002@yahoo.es

MSC: 97M50, 40G99 , 40G10 , 81T55

- **ABSTRACT:** We study a generalization of the zeta regularization method applied to the case of the regularization of divergent integrals  $\int_0^{\infty} x^s dx$  for positive 's', using the Euler Maclaurin summation formula, we manage to express a divergent integral in term of a linear combination of divergent series , these series can be regularized using the Riemann Zeta function  $\zeta(s)$   $s > 0$  , in the case of the pole at  $s=1$  we use a property of the Functional determinant to obtain the regularization  $\sum_{n=0}^{\infty} \frac{1}{(n+a)} = -\frac{\Gamma'(a)}{\Gamma(a)}$ , with the aid of the Laurent series in one and several variables we can extend zeta regularization to the cases of integrals  $\int_0^{\infty} f(x)dx$  , we believe this method can be of interest in the regularization of the divergent UV integrals in Quantum Field theory since our method would not have the problems of the Analytic regularization or dimensional regularization
- **Keywords:** = Riemann Zeta function, Functional determinant, Zeta regularization, divergent series .

**ZETA REGULARIZATION FOR DIVERGENT INTEGRALS:**

Sometimes in mathematics and physics , we must evaluate divergent series of the form

$\sum_{n=1}^{\infty} n^k$  , of course this series is divergent unles  $\text{Re}(k) > 1$  , however cases like  $k=1$  or

$k=3$  appear in several calculations of string theory and Casimir effect , for the case of Casimir effect [3] the result  $\sum_{n=1}^{\infty} n^3 = \frac{1}{120}$  appears to give the correct result for the

Casimir force  $\frac{F_c}{A} = -\frac{\hbar c \pi^2}{240 a^4}$  here A is the area and 'd' the separation between the 2 plates , c and  $\hbar$  are the speed of light and the Planck's constant. The idea behind the Zeta regularization method is to take for granted that for every 's' the identity

$\sum_{n=1}^{\infty} n^s = \zeta(s)$ , follows although this formula is valid just for  $\text{Re}(s) > 1$  , to extend the definition of the Riemann Zeta function to negative real numbers, one need to use the functional equation for the Riemann function

$$\zeta(1-s) = 2(2\pi)^{-s} \Gamma(s) \cos\left(\frac{\pi s}{2}\right) \zeta(s) \quad \Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin(\pi s)} \quad (1)$$

This gives the expressions  $\sum_{n=1}^{\infty} n^0 = -\frac{1}{2}$  ,  $\sum_{n=1}^{\infty} n = -\frac{1}{12}$  and  $\sum_{n=1}^{\infty} n^2 = 0$  due to the pole at  $s=1$  , the Harmonic series  $\sum_{n=1}^{\infty} n^{-1}$  is NOT zeta regularizable, although it can be given

a finite value  $\sum_{n=1}^{\infty} n^{-1} = \gamma = 0.577215..$  , this value can be justified by using the theory of Zeta-regularized infinite products (determinants) , as we shall see later in the paper

o *Zeta regularization for divergent integrals:*

Let be  $f(x) = x^{m-s}$  with  $\text{Re}(m-s) < -1$  , then the Euler-Maclaurin summation formula for this function reads

$$\int_a^{\infty} x^{m-s} dx = \frac{m-s}{2} \int_a^{\infty} x^{m-1-s} dx + \zeta(s-m) - \sum_{i=1}^a i^{m-s} + a^{m-s} - \sum_{r=1}^{\infty} \frac{B_{2r} \Gamma(m-s+1)}{(2r)! \Gamma(m-2r+2-s)} (m-2r+1-s) \int_a^{\infty} x^{m-2r-s} dx \quad a \in N \quad (2)$$

Here in formula (2) all the series and integrals are convergent, formula (2) is usually worthless , since it is trivial to prove that  $\int_a^{\infty} x^{-k} dx = \frac{a^{1-k}}{k-1}$  for  $\text{Re}(k) > 1$  , and the Riemann

zeta function  $\zeta(m-s) = \sum_{i=1}^{\infty} i^{m-s}$  , so nothing new can be obtained from (2) , the idea is

to use the Functional equation (1) for the Riemann and Zeta function to extend the definition of equation (2) to the whole complex plane except  $s=1$  , in case (m-s) is positive there will be no pole at  $x=0$  , so we can put  $a=0$  and take the limit  $s \rightarrow 0^+$

$$\int_0^{\infty} x^m dx = \frac{m}{2} \int_0^{\infty} x^{m-1} dx + \zeta(-m) - \sum_{r=1}^{\infty} \frac{B_{2r} m! (m-2r+1)}{(2r)! (m-2r+1)!} \int_0^{\infty} x^{m-2r} dx \quad (3)$$

Formula (3) is the Analytic continuation of formula (2) with  $a=0$  and can be used to obtain a finite definition for otherwise divergent integrals, apparently this recurrence equation has an infinite number of terms but the Gamma function has a pole at  $x=0$  and at  $x$  being some negative integer, some examples of formula (3)

$$I_0 = \zeta(0) + 1 = \int_0^{\infty} dx \quad I_1 = \frac{I_0}{2} + \zeta(-1) = \int_0^{\infty} x dx$$

$$I_2 = \left( \frac{I_0}{2} + \zeta(-1) \right) - \frac{B_2}{2} a_{21} I_0 = \int_0^{\infty} x^2 dx \quad (4)$$

$$I_3 = \frac{3}{2} \left( \frac{1}{2} (I_0 + \zeta(-1)) - \frac{B_2}{2} a_{21} I_0 \right) + \zeta(-3) - B_2 a_{31} I_0 = \int_0^{\infty} x^3 dx$$

So our method can provide finite ‘regularization’ to divergent integrals, with the Aid of the zeta regularization algorithm. Also our formulae (2) (3) and (4) are consistent with the usual summation properties, in fact if  $\int_0^{\Lambda} x^m dx$  is finite for finite  $\Lambda$  and we use the property of the Riemann and Hurwitz Zeta function [ ] to get the sum of the  $k$ -th powers of  $n$  on the interval  $[0, \Lambda]$   $\sum_{i=0}^{\Lambda-1} i^m = \zeta(-m) - \zeta(-m, \Lambda)$ ,  $\zeta(s, \Lambda) = \sum_{n=0}^{\infty} (n + \Lambda)^{-s}$  defined for  $\text{Re}(s) > 1$  (of course for positive ‘ $s$ ’ as  $\Lambda \rightarrow \infty$  the second term goes to 0)

$$\int_0^{\Lambda} x^m dx = \frac{m}{2} \int_0^{\Lambda} x^{m-1} dx + \zeta(-m) - \zeta(-m, \Lambda) - \sum_{r=1}^{\infty} \frac{B_{2r} m! (m-2r+1)}{(2r)! (m-2r+1)!} \int_0^{\Lambda} x^{m-2r} dx \quad (5)$$

For integer ‘ $m$ ’  $\zeta_H(-m, x) = -\frac{B_{m+1}(x)}{m+1}$  we find the Bernoulli Polynomials, the powers of  $\Lambda$  would cancel the integral  $\int_0^{\Lambda} x^m dx = \frac{\Lambda^{m+1}}{m+1}$ , so in the end in formula (5) we would

get the usual definition of Zeta regularization  $\zeta_H(-m) = -\frac{B_{m+1}(0)}{m+1}$  for integer ‘ $m$ ’. Of course one could argue that a ‘simpler’ regularization of the divergent integrals should be  $I(s) = \int_0^{\infty} dx (x+a)^s = -\frac{a^{s+1}}{s+1}$  and  $I(-1) = \int_0^{\infty} dx (x+a)^{-1} = -\log a$ , this is just dropping

out the term proportional to  $\log \infty$  or  $\infty^{s+1}$  inside the integral to make it finite, however if we plugged this result into the Euler-Maclaurin summation formulae (2) (3) or (5) the terms involving ‘ $a$ ’ would cancel and we would finally find that  $\zeta_H(-m) = 0$  for every ‘ $m$ ’ which clearly is against the definition of zeta regularization of a series, for the case

of the logarithmic divergence, obtained from differentiation with respect to the external parameter 'a' this is a result of taking the finite part of the integral, which apparently works. For the case of the integrals  $\int_a^\infty x^{m-s} \log^k(x) dx$ , we can simply differentiate k-times with respect to regulator 's' in order to obtain finite values in terms of  $\zeta(-s)$  and  $\zeta'(-s)$  for negative values of 's' unless  $m=-1$  (for other negative values of m we can make a change of variable  $xq=1$ ), this is treated in the next section

○ *Zeta-regularized determinants and the Harmonic series:*

Given an operator A with an infinite set of nonzero Eigenvalues  $\{\lambda_n\}_{n=0}^\infty$  we can define a Zeta function and a Zeta-regularized determinant, Voros [10]

$$\text{Tr}\{A^{-s}\} = \zeta_A(s) = \sum_{n=0}^\infty \lambda_n^{-s} \quad \det(A) = \prod_{n=0}^\infty \lambda_n = \exp\left(-\frac{d\zeta_A(0)}{ds}\right) \quad (6)$$

The proof of the second formula inside (6) is pretty easy, the derivative of the Generalized zeta function will be  $\zeta_A'(s) = -\sum_{n=0}^\infty \frac{\log \lambda_n}{\lambda_n^s}$  now let  $s=0$ , use the property of the logarithm  $\log(a.b) = \log a + \log b$  and take the exponential on both sides.

For the case of the Eigenvalues of a simple Quantum Harmonic oscillator in one dimension [10]  $\lambda_n = n + a$ , the Zeta function is just the Hurwitz Zeta function, so we can define a zeta-regularized infinite product in the form

$$\prod_{n=0}^\infty (n + a) = \exp\left(-\frac{d\zeta_H(0, a)}{ds}\right) \quad \frac{d\zeta_H(0, a)}{ds} = \log \Gamma(a) - \log(\sqrt{2\pi}) \quad (7)$$

In case we put  $a=1$  we find the zeta-regularized product of all the natural numbers

$$\prod_{n=0}^\infty (n + 1) = \sqrt{2\pi}, \text{ see [5] if we take the derivative with respect to 'a', we would find}$$

$$\text{the same regularized Value Ramanujan did [2] precisely } \sum_{n=0}^\infty \frac{1}{(n + a)} = -\frac{\Gamma'(a)}{\Gamma(a)} \quad a > 0$$

Harmonic series appear due to a logarithmic divergence of the integral  $\int_0^\infty \frac{dx}{(n + a)}$ , if we

put  $m = -1$  inside formula (2), using a regulator 's',  $s \rightarrow 0^+$  we have the Euler Maclaurin summation formula

$$\int_0^\infty \frac{dx}{(n + a)^{s+1}} = -\frac{1}{2a} + \sum_{n=0}^\infty \frac{1}{(n + a)^{1+s}} + \sum_{r=1}^\infty \frac{B_{2r}}{(2r)!} \frac{\partial^{2r-1}}{\partial u^{2r-1}} \left( \frac{1}{(x + a)^{s+1}} \right)_{x=0} \quad (8)$$

Since  $s > 0$  the integral and the series inside (8) will be convergent, now we can integrate over 'a' inside (8) and use the definition of the logarithm  $\lim_{s \rightarrow 0^+} \frac{x^s - 1}{s} = \log x$ , to

regularize the integral  $\int_0^{\infty} \frac{dx}{(n+a)^{s+1}}$  as  $s \rightarrow 0^+$  in terms of the function  $-\frac{\Gamma'}{\Gamma}(a)$  plus some finite corrections due to the Euler-Maclaurin summation formula.

A faster method is just simple differentiate with respect to 'a' inside the integral

$$\int_0^{\infty} \frac{dx}{(n+a)^2} = -\frac{dI}{da}, \text{ now this integral is convergent for every 'a' and equal to } \frac{1}{a},$$

integration over 'a' again gives the value  $-\log a + c$  plus a constant 'c' that will not depend on the value of a inside the integral in question, the proof that 'c' is unique no

matter what a is comes from the fact that the difference  $\int_0^{\infty} dx \left( \frac{1}{x+a} - \frac{1}{x+b} \right) = \log \left( \frac{b}{a} \right)$ .

For the case  $a=0$ , the derivative of the Hurwitz Zeta is  $\frac{d\zeta_H(0,0)}{ds} = -\log(\sqrt{2\pi})$  so if

we approximate the divergent integral by a series, then we can get the regularized result

$$\int_0^{\infty} \frac{dx}{x} \approx \sum_{n=0}^{\infty} \frac{1}{n} = 0. \text{ Apparently it seems that using two different regularizations we get}$$

some different results, the idea is that if we use the Stirling asymptotic formula approximation for the logarithm of the Zeta function

$$\log \Gamma(z) = \left( z - \frac{1}{2} \right) \log z - z + \frac{1}{2} \log(2\pi) + \sum_{r=1}^{\infty} \frac{B_{2r} z^{1-2r}}{2r(2r-1)} \quad (9)$$

If we take the derivative with respect to 'z' inside (9), is now more apparent that for

the logarithmic derivative  $\int_0^{\infty} \frac{dx}{x+a} \approx \log \left( \frac{\mu}{a} \right)$  here  $c = \log \mu$  is a constant obtained from

differentiation with respect to 'a' to regularize the divergent integral, this constant 'c' must be related to some physical constant or in case the quantity 'a' has dimension of Energy then  $\mu$  must have also dimensions of energy so the logarithm is dimensionless, this constant 'c' would be the only free adjustable parameter that would appear inside our calculations to regularize integrals. If 'a' is negative there is an extra term due to the value  $\log(-1) = \pi i$ , for more complex logarithmic integral one can use the definition

$$\int_0^{\infty} \frac{\log^k(x+a) dx}{x+a} \approx \frac{1}{k+1} \log^{k+1} \left( \frac{\mu}{a} \right) \text{ with the same energy scale } c = \log \mu$$

o *Regularization of divergent integrals*  $\int_0^{\infty} dx f(x) :$

In general, the divergent integrals that appear in Quantum Field Theory [ ] are invariant

under rotations, for example  $\int \frac{d^4 p}{(p^2 + m^2)^2}$  or  $\int \frac{d^4 p}{((p-q)^2 + m^2)} \frac{1}{p^2}$ , if we use 4-

dimensional polar coordinates we can reduce these integrals to the case

$\frac{2\pi^{d/2}}{\Gamma(d/2)} \int_0^\infty dr f(r) r^{d-1}$  then the UV divergences appear when  $r \rightarrow \infty$ , here  $d=4$  is the dimension of the spacetime, depending on the value of 'd' we can have several types of divergences  $\int_0^\Lambda dr f(r) r^{d-1} \approx a\Lambda^{m+1} + b \log \Lambda$ , if  $b=0$  for  $m=2$  the UV divergences are quadratic if  $m=0$  the divergences are linear, in case  $a=0$  and  $b=1$  the divergences are of logarithmic type, for example  $\int \frac{d^4 p}{(p^2 + m^2)^2}$  has only a logarithmic divergence in dimension 4, for a lower value of the dimension ( $d=3$ ) this integral exists.

To study the rate of divergence, we can expand the function into a Laurent series valid for  $z \rightarrow \infty$ ,  $f(x) = \sum_{n=-\infty}^{n=k} c_n (x+a)^n$  'k' is a finite number and means that the function  $f(x)$  has a power law divergence for big 'x', then the idea to compute a divergent integral would be this, we add and subtract a Polynomial plus a term proportional to  $\frac{1}{x+a}$  to split the integral into a finite part and another divergent integrals

$$\int_0^\infty dx \left( f(x) - \sum_{n=0}^k b_n x^n - \frac{b_{-1}}{x+a} \right) + \sum_{n=0}^k b_n \int_0^\infty x^n dx + b_{-1} \int_0^\infty \frac{dx}{x+a} = \int_0^\infty f(x) dx \quad (10)$$

The number of terms 'k' is chosen so the first integral is FINITE, this first integral can be computed by Numerical or exact methods and yields to a finite value, the rest of the integrals are just the logarithmic and power-law divergences, they can be regularized with the aid of formulae (2) (3) (4) (6) (8) to get a finite value involving a linear

combination of  $\zeta(-m)$   $m=0,1,2,\dots,k$  and another value proportional to  $\frac{\partial}{\partial s} \frac{\partial \zeta_H(a,0)}{\partial a}$

or  $\int_0^\infty \frac{dx}{x+a} \approx \log\left(\frac{\mu}{a}\right)$  for example we can analyze this simple divergent integral  $a > 0$

$$\int_a^\infty \frac{x^2 dx}{x+1} = \int_a^\infty dx \left( \frac{x^2}{x+1} - 1 + x + \frac{1}{x} \right) + \frac{\zeta(0)}{2} - a + \frac{a^2}{2} + \frac{1}{2a} + \frac{\Gamma'(a)}{\Gamma(a)} - \sum_{r=1}^\infty \frac{B_{2r}}{(2r)!} \frac{\partial^{2r-1}}{\partial u^{2r-1}} \left( \frac{1}{x+a} \right)_{x=0} - \zeta(-1) + \frac{1}{2} \quad (11)$$

The first integral in (11) is convergent and have an exact value of  $\log\left(\frac{a+1}{a}\right)$ , in order to regularize the logarithmic integral we have used the result  $\sum_{n=0}^\infty \frac{1}{(n+a)} = -\frac{\Gamma'(a)}{\Gamma(a)}$  plus the Euler-Maclaurin summation formula. The mathematical justification of this is the following, given a divergent integral  $\int_a^\infty dx f(x)$  we introduce a regulator

$F(s) = \int_a^{\infty} f(x) \frac{dx}{x^s}$  so the integral  $F(s)$  exists for some big 's', if we add and subtract powers of the form  $x^{k-s}$  for integer  $k$  and  $(x+a)^{s+1}$ , we can split  $F(s)$  into a convergent

integral  $I(s)$  valid for  $s \rightarrow 0^+$  and some divergent integrals of the form  $\int_a^{\infty} x^{m-s} dx$  and  $\int_0^{\infty} \frac{dx}{(x+a)^{s+1}}$ , using formulae (2) (3) (4) and (8) we can express these integrals in terms

of the series  $\sum_{n=0}^{\infty} \frac{1}{(n+a)^{s+1}}$  and  $\sum_{n=0}^{\infty} \frac{1}{(n+a)^{s-m}}$ , which will be convergent for

$\text{Re}(s-m) > 1$  and  $\text{Re}(s+1) > 1$ , now using the Functional equation for the Hurwitz and Riemann Zeta function we can make the analytic continuation of both series to  $s \rightarrow 0^+$  avoiding the pole at  $s=1$  by the use of Riemann Zeta function at negative integers  $\zeta(-n)$  plus some corrections involving  $-\frac{\Gamma'}{\Gamma}(a)$  of course the rules for change

of variable and still valid so  $\int_0^{\infty} dx f(x+a) = \int_a^{\infty} du f(u)$  this can be used to avoid some IR

divergences at  $x=0$  by splitting the integral into an IR divergent part and an UV

divergent part  $\int_0^{\infty} du = \int_0^a du + \int_a^{\infty} du$ . For other types of divergent integrals like

$\int_a^{\infty} dx \log^{\beta}(x) x^{\alpha}$  for positive  $\alpha$  and  $\beta$  one could differentiate with respect to 'm' or 's'

inside formula (2) in order to obtain a recurrence equation for the integrals

$\int_a^{\infty} dx \log^{\beta}(x) x^{\alpha}$ , this recurrence equation is finite (approximately) since for  $\text{Re}(p) > 1$

$\int_a^{\infty} dx \frac{\log^{\beta}(x)}{x^p}$  is finite and do not need to be regularized provided  $a > 0$ . Other useful

identities can be  $(1+x)^{1/2} \approx 1 + \frac{x}{2} - \frac{x^2}{2.4}$  or the expansion of the logarithm valid for any

$x > 0$   $\log x = 2 \sum_{n=0}^{\infty} \frac{1}{2n+1} \left( \frac{x-1}{x+1} \right)^{2n+1}$  to make logarithms more tractable, also we could

use Laurent expansions to handle complicate non-Polynomial expressions like  $(x^n + \mu^n)^k$  by expanding it for big 'x' into asymptotic (inverse) power series.

- *Regularization of integrals in the form  $\int_0^{\infty} \frac{dx}{x^m}$  and  $\int_0^{\infty} \frac{f(x)dx}{(x-a)^m (x-b)^m}$ :*

Until now, we have only considered the UV divergent integrals, the integrals whose integrand  $f(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , from the definition of an improper integral

$$\int_{\varepsilon}^{\infty} \frac{dx}{x^m} = \frac{\varepsilon^{m-1}}{m-1} = F_{reg}(m) \quad \int_0^{1/\varepsilon} x^{m-2} dx = \frac{\varepsilon^{-(m-1)}}{m-1} = F_{reg}(2-m) \quad \varepsilon = \frac{1}{N} \quad (12)$$

As  $N \rightarrow \infty$ , this imply that in our regularization producedure  $F_{reg}(s) = F_{reg}(2-s)$ , for the case  $s > 0$  we can use formulae (2) and (3) to regularize the divergent integral, and by the formula relating  $s$  and  $(2-s)$  one could also regularize IR (infrared) divergent

integrals  $\int_0^{\infty} \frac{dx}{x^m}$  in a similar way we did for  $\int_0^{\infty} x^m dx$ , except for the case  $m = 1$

(logarithmic integral), for the case of this integral  $\int_0^{\infty} \frac{dx}{x}$  one could split it

into  $\int_0^a \frac{dx}{x} + \int_a^{\infty} \frac{dx}{x}$  now we make the change of variables,  $x \rightarrow x+a$   $x \rightarrow x+a^{-1}$  and

$x \rightarrow \frac{1}{x}$  to rewrite this as  $\int_0^{\infty} \frac{dx}{x+a} + \int_0^{\infty} \frac{dx}{x+1/a}$ , these are again logarithmic divergent

integrals and can be regularized with the aid of the zeta regularized product

$\prod_{n=0}^{\infty} (n+b) = e^{-\zeta'_H(0,b)}$  plus the Euler-Maclaurin summation formula. For the case of a more general divergent integral like

$$\int_0^a \frac{f(x)dx}{(x-c)^m} \rightarrow \int_0^a \frac{f(x) - \sum_k f^{(k)}(c)(x-c)^k}{(x-c)^m} dx + \sum_k \int_0^a \frac{f^{(k)}(c)dx}{(x-c)^{m-k}} \quad a > c > 0 \quad (13)$$

First integral inside (13) is finite and after several manipulations the other divergent

integrals can be written as  $\int_0^{\infty} \frac{x^r dx}{(x^2 - (c+i\varepsilon)^2)^{m-k}}$  for some real and positive parameters

'r' 'c' 'm' and 'k', by multiplying both numerator and denominator by  $(x+c)^{m-r}$

Another possibility is to avoid the pole at a certain point  $x = a$  by using the Analytic continuation of the integral involving several parameters

$$\int_0^{\infty} \frac{dx}{x^m} = \int_0^{\infty} \frac{dx(x+a)^m}{x^m(x+a)^m} = \sum_{i=0}^m \binom{m}{i} \int_0^{\infty} \frac{x^{m-i} a^i dx}{(x-\alpha_1)^2 + \alpha_2^2} = F(\alpha_1, \alpha_2) \quad (14)$$

The main idea is to calculate the integral (14) that will depend on two parameters  $\alpha_i$

$i=1,2$  and finally set  $\alpha_1 = -\frac{a}{2}$ ,  $\alpha_2 = \pm \frac{ai}{2}$  if  $F\left(-\frac{a}{2}, \frac{\pm ai}{2}\right)$  exists, this can be

regarded as the regularized value of the integral, of course (14) may be divergent as  $x \rightarrow \infty$  so we may need to add and substract terms to make it convergent in a similar

way we did in (10), another form to regularize (14) is defining  $F\left(-\frac{a}{2}, \alpha_2\right)$  and then



calculate this integral for a general value of  $\alpha_2$ , in the end we would put  $\alpha_2 = \pm \frac{ai}{2}$ .

Another useful identity to regularize infrared divergences whenever  $x = a$  is (tables of integrals by Amabrowitz and Stegun [1] )

$$\int_0^{\infty} \frac{x^m dx}{(x^2 - (a + i\varepsilon)^2)^r} = \frac{(-1)^{r-1} \pi (-ia)^{m+1-2r} \Gamma\left(\frac{m+1}{2}\right)}{2 \sin\left(\frac{m\pi + \pi}{2}\right) (r-1)! \Gamma\left(\frac{m+1}{3-r}\right)} = I(a, m, r) \quad \varepsilon \rightarrow 0 \quad (15)$$

Integral inside (14) will be convergent whenever  $2r-m > 1$  if this is not the case we could use the Euclidean algorithm to split this integral into a convergent term defined as (14)

and some divergent integrals  $\int_0^{\infty} dx (x+a)^m$   $m = -1, 0, 1, 2, \dots$  the main idea to justify why

the infrared divergences are easier to regularize than ultraviolet ones, is that for the infrared, you could insert a small complex term  $i\varepsilon$  to regularize it and make it convergent, so there are some complex values of 'a' that make (14) to be well-defined, however for the ultraviolet divergences this is not the case since there is no value of 'a'

that makes  $\int_0^{\infty} \frac{x^4 dx}{(x+a)^2}$  convergent, unless we use some kind of regularization

## REGULARIZATION OF MULTIPLE INTEGRALS:

Until now, we have only considered integrals in one variable (after change to polar coordinates), then it arises the question if one can apply our method of zeta regularization to more complicate integrals like

$$I(s) = \int d^4 q_1 \int d^4 q_2 \dots \int d^4 q_n \prod_{i=1}^{\infty} \frac{1}{(1+q_i^2)} F(q_1, q_2, \dots, q_n) (R(q_1, q_2, \dots, q_n))^{-s} \quad (16)$$

Here we have introduced a regulator depending on an external parameter 's' in order the integral (16) to converge for big 's' and then use the analytic regularization to take the limit  $s \rightarrow 0^+$ , this regulator must be chosen with care in order not to spoil any symmetries of the Physical system this regulator may be of the form

$$R(q_1, q_2, \dots, q_n) = 1 + \sum_{i=1}^n q_i^2 \quad R(q_1, q_2, \dots, q_n) = \prod_{i=1}^n (1+q_i) \quad (17)$$

Our first ansatz would be to define n-dimensional polar coordinates so we can rewrite

(16) as a multiple integral depending on 'r'  $\sqrt{\sum_{i=1}^n q_i^2} = r$  and several angles  $\theta_i$   $i =$

1,2,3,4,..., n-1 in the form

$$I(s) = \int_{\Omega} d\Omega \int_0^{\infty} dr G(r, \theta_i) r^{n-1} (R(r, \theta_i))^{-s} \quad d\Omega = \prod_{i=1}^{n-1} d\theta_i \sin^{n-i-1}(\theta_i) \quad (18)$$

We may choose the first regulator inside (17) so it does not depend on the angular coordinates, the idea is that in case (16) has an ultraviolet divergence this divergence will appear whenever  $r \rightarrow \infty$ , so if we perform the integral over the angular variables  $d\Omega = \prod_{i=1}^{n-1} d\theta_i \sin^{n-i-1}(\theta_i)$  we are left with an integral  $I(s) = \int_0^{\infty} dr U(r) r^{n-1} (1+r)^{-s}$ , in order to regularize this we define a convergent integral (by subtraction) plus some divergent terms

$$I(s) = \int_0^{\infty} dr (1+r)^{-s} \left( U(r) r^{n-1} - \sum_{i=-1}^k a_i (1+r)^i \right) + \sum_{i=-1}^k a_i \int_0^{\infty} (1+r)^{i-s} dr \quad (19)$$

$U(r)$  is the function obtained after integration over the angles, and 'k' is a finite number to perform the minimal subtraction of terms in order the first integral to be convergent even for  $s = 0$ , if the integral over the angles is too complicate to have an exact form we could replace this integral over the angles by an approximate finite sum  $d\Omega \rightarrow \sum_i$  (sum over all the angular variables) in order to make the integral easier to calculate, this can be using Montercarlo methods of integration.

○ *Substraction method:*

Once we have made the change of variable to spherical coordinates inside our integral  $I(q_1, q_2, \dots, q_n)$  one could substract some terms to render the integral finite

$$I(s) = \int_{\Omega} d\Omega \int_0^{\infty} dr \left( G(r, \theta_i) r^{n-1} - \sum_{j=-1}^k f_j(\theta_i) (1+r)^{j-s} \right) + \int_{\Omega} \sum_{j=-1}^k f_j(\theta_i) d\Omega \int_0^{\infty} dr (1+r)^{j-s} \quad (20)$$

We chose the number 'k' and the functions  $f_j(\theta_i)$  so the first integral inside (20) is convergent, for the second integral we could perform integration over the angular variables and then use formulae (2) and (3) to regularize  $\int_0^{\infty} (1+r)^m dr$ .

Another method is to consider the multiple integral as an interate integral and then make the substraction for every variable for example

$$\int \partial q_n \left( F(q_1, q_2, \dots, q_n) - \sum_{i=-1}^k a_i(q_1, q_2, \dots, q_{n-1}) (1+q_n)^i \right) + \int_0^{\infty} \partial q_n \sum_{i=-1}^k a_i(q_1, q_2, \dots, q_{n-1}) (1+q_n)^i \quad (21)$$

The symbol  $\partial q_n$  means that the integral is made over the variable  $q_n$  keeping the other variables constant, the number 'k' is chosen so the first integral is finite, this integral

will depend on  $I(q_1, q_2, \dots, q_{n-1})$ , the divergent integrals (even for the logarithmic case  $i=-1$ ) can be regularized.

Now we have regularized the first integral, we have reduced in one variable the multiple integral, repeating the iterative process for the functions  $a_i(q_1, q_2, \dots, q_{n-1})$

$$\int \partial q_{n-1} \left( a_i(q_1, q_2, \dots, q_{n-1}) - \sum_{j=1}^k b_j(q_1, q_2, \dots, q_{n-2})(1+q_{n-1})^j \right) + \int_0^\infty \partial q_{n-1} \sum_{j=1}^k b_j(q_1, q_2, \dots, q_{n-2})(1+q_{n-1})^j \quad (22)$$

Using (21) and (22) for every step we can reduce the dimension of the integral until we reach to the one dimensional case, which is easier to handle. As an example

$$\int_0^\infty dx \int_0^\infty dy \frac{xy}{x+y+1} = \int_0^\infty dx \int_0^\infty dy \left( \frac{xy}{x+y+1} - x + \frac{x+x^2}{y+1} \right) + \int_0^\infty x dx \int_0^\infty dy - \int_0^\infty (x+x^2) dx \int_0^\infty \frac{dy}{y+1} \quad (23)$$

$$\int_0^\infty dx (f(x) - bx^2 + (a-b)x) \quad f(x) = \int_0^\infty \frac{dy}{(y+1)} \frac{x^3+x^2}{(x+y+1)} \quad (24)$$

With  $a = \left( \int_0^\infty dx \right)_{reg}$   $b = \left( \int_0^\infty \frac{dy}{y+1} \right)_{reg}$  so for an initial given integral with an overlapping

divergence as  $x \rightarrow \infty$   $y \rightarrow \infty$  we have made a subtraction to get a finite integral over 'y' (23) repeating the same process we can regularize the integral over 'x', in order to integrate the finite part of the integral we can use several numerical methods. For

example, the integral  $f(x) = \int_0^\infty \frac{dy}{(y+1)} \frac{x^3+x^2}{(x+y+1)}$  can be calculated numerically to give

$f(x) \approx \sum_j \frac{1}{(y_j+1)} \frac{x^3+x^2}{(x+y_j+1)}$  in order to avoid terms with  $\log(x)$ ,  $\arctan(x)$  or similar

ones inside (23) and (24), another method to calculate (23) would be to introduce a regulator in the form  $(1 + \sqrt{x^2 + y^2})^{-s} = R(s, x, y)$ , that will make (23) to converge for certain values of 's', if we use polar coordinates in (x,y) this becomes

$$\int_0^\infty dx \int_0^\infty dy (xy) \frac{R(x, y, s)}{x+y+1} = \int_0^\infty dr \int_0^{\pi/2} du \frac{r^2 \sin(2u)(1+r)^{-s}}{2(\cos(u) + \sin(u) + r^{-1})} \quad (25)$$

Integration over the angular variable 'u' can be carried by numerical methods, to produce some new divergent integrals, that will only depend on the value of 'r'

$\int_0^\infty dr g_i(r)(1+r)^{-s}$ , this integrals can be regularized with the same methods we used to

give a finite meaning to 1-D integrals, unlike the dimensional regularization, the 'regulator' is not the dimension of space-time, so we have no problem whenever changing to spherical coordinates in d=4 to overcome the UV divergencies.

If the integrand  $F(q_1, q_2, \dots, q_n)$  had no singularities for every  $q_j > 0$ , we may expand this integrand into a multiple Laurent series of several variables, and then perform the subtraction  $\sum_{m_1, m_2, \dots, m_n = -1}^{s_1, s_2, \dots, s_n} C_{m_1, m_2, \dots, m_n} (q_1 + b_1)^{m_1} (q_2 + b_2)^{m_2} \dots (q_n + b_n)^{m_n}$  in order to define a finite part of the integral

$$\int d^4 q_1 \int d^4 q_2 \dots \int d^4 q_n \left( F - \sum_{m_1, m_2, \dots, m_n = -1}^{s_1, s_2, \dots, s_n} C_{m_1, m_2, \dots, m_n} (q_1 + b_1)^{m_1} (q_2 + b_2)^{m_2} \dots (q_n + b_n)^{m_n} \right) \quad (26)$$

Plus some corrections due to divergent integrals  $\int_0^\infty (q_i + b_i)^m dq_i$   $m = -1, 0, 1, \dots$ . In many cases although the integrals given in (21) and (22) are finite they will have no exact expression or the exact expression will be too complicate, in this case we can use the Gauss-Laguerre Quadrature formula (in case the interval is  $[0, \infty)$ ) to approximate the integral by a sum over the zeros of Laguerre Polynomials  $\sum_{i=0}^n w_i f(q_1, q_2, \dots, q_{n-1}, x_i)$  with the weight expressed in terms of Laguerre Polynomials and their roots

$$w_i = \frac{x_i}{(n+1)^2 (L_{n+1}(x_i))^2}, \quad L_n(x_i) = 0$$

## FRACTIONAL DERIVATIVES FOR IR AND UV DIVERGENCES:

Another form to deal with IR divergences using our Zeta regularization algorithm (2) and (3) is the following, the term  $(p^2 + i\varepsilon - m^2)^{-1/2}$   $\varepsilon \rightarrow 0$  can be regarded as the derivative with respect to  $-m^2$  of the UV divergent expression  $\sqrt{p^2 + i\varepsilon - m^2}$ , so if we wish to calculate the following integral  $\int_0^\infty dx \frac{x^k}{\sqrt{x^2 + i\varepsilon - m^2}} = J(k, -m^2)$  we must use our zeta-regularization algorithm in order to give a finite meaning to  $\int_0^\infty x^k dx \sqrt{x^2 + i\varepsilon - m^2}$ ,

we split the integrand into  $\int_0^{m+1} + \int_{m+1}^\infty$  the first integral is no longer problematic at  $x = m$

the second integral may be written as  $\int_{m^2+1}^\infty x^{k+1} dx \sqrt{1 + \frac{i\varepsilon - m^2}{x^2}}$ , now we can use the

Binomial theorem to expand the square root in powers of  $\left(\frac{i\varepsilon - m^2}{x^2}\right)$  and valid for

$x > m$  and use formulae (2) and (3) to regularize the divergent integrals  $\int_{m+1}^\infty x^{k+1+i-s} dx$

.For other non-integer exponent  $(p^2 + i\varepsilon - m^2)^{-\alpha}$  we may simply use the formula of fractional calculus  $\frac{\Gamma(3/2)}{\Gamma(1-\alpha)} x^{-\alpha} = \frac{d^{1/2+\alpha} \sqrt{x}}{dx^{1/2+\alpha}}$  with this method we turn an UV divergence into an IR one by formal differentiation-integration, for the case of ‘n’ integer  $(p^2 + i\varepsilon - m^2)^{-n}$ , we only have to study the case n=1 since by differentiation with respect to  $-m^2$  other cases can be obtained, we use partial fraction decomposition and division of Polynomials to set the following identities

$$(p^2 - m^2)^{-1} = \frac{1}{2m} \left( \frac{1}{p-m} - \frac{1}{p+m} \right) \quad \frac{p^k}{p \pm m} = F_{\pm}(p) + \frac{C_{\pm}}{p \pm m} \quad (27)$$

with F a polynomial on ‘p’ and C a constant, now in the sense of Principal value

$$\int_0^{\infty} \frac{dx}{x-a} = \log\left(\frac{\mu}{a}\right) + i\pi \quad \int_0^{\infty} \frac{dx}{x+a} = \log\left(\frac{\mu}{a}\right) \quad (28)$$

Here  $c = \log \mu$  (Energy or UV scale) is a divergent constant obtained in the regularization of the logarithmic divergent integral, due to the pole at  $s=1$  we must recall the definition of Functional determinant to define a zeta-regularized product

$e^{-\zeta_H'(0,a)} = \prod_{n=0}^{\infty} (n+a)$  and take the logarithmic derivative with respect to ‘a’ and apply the expansion of the Digamma function  $\frac{\Gamma'}{\Gamma}(a)$  plus the Euler-Maclaurin formula that relates series and integrals.

If we wished to calculate an integral involving the term  $(p^2 - m^2)^{-1}$  we would have a problem since  $\frac{\Gamma(\varepsilon)}{\Gamma(3/2)} \frac{\partial^{3/2-\varepsilon}}{\partial(-m^2)^{3/2-\varepsilon}} \sqrt{p^2 - m^2} = (p^2 + i\varepsilon - m^2)^{-1+\varepsilon}$  become singular as  $\varepsilon \rightarrow 0^+$ , if we expand the Gamma function  $x\Gamma(x) = \Gamma(x+1)$  near its poles in the form  $\Gamma(\varepsilon) \approx \frac{1}{\varepsilon} - \gamma + O(\gamma)$  and introduce an IR cut-off  $\varepsilon = \varepsilon_{IR}$ , then we will be able to

define a ‘regularized’ fractional derivative as  $\frac{-2\gamma}{\sqrt{\pi}} \frac{d^{3/2} \sqrt{x}}{dx^{3/2}} =_{reg} \frac{1}{x}$  so we can turn UV

divergent quantities such as  $\sqrt{p^2 - m^2 + i\varepsilon}$  into IR divergent quantities at  $p=m$

$(p^2 - m^2 + i\varepsilon)^{-1}$ , this would explain a certain relationship we conjectured before involving an IR divergent integral with another UV divergent integral by using formal fractional differentiation with respect to external parameters.

○ *Multiple integrals:*

For the case of multiple integrals, we can make use of Polar coordinates in any integer dimension  $n$  and then replace the integral over the angles  $\int d\Omega$  by a multiple sum

$$\int dVF(q_1, \dots, q_n) \approx \sum_{\Omega_i} \int_0^\infty r^{n-1} dr \varphi(r, \Omega_i) (1+r)^{-s} \prod_{i=0}^n \sqrt{b_i(\Omega) - m_i^2} \sqrt{f(r, c(\Omega))} \quad (29)$$

The functions ‘a’ ‘b’ and ‘c’ depend only on the ANGLES but not on the variable ‘r’, for multiple integrals we can not always obtain exact results for the integration over the angles so we should settle for approximations.

For the case of the logarithmic divergent integral, we can still apply another trick we use the identities  $\frac{1}{x} = \frac{1}{\sqrt{x}} \cdot \frac{1}{\sqrt{x}}$  and  $(\sqrt{x} + 1)(\sqrt{x} - 1) = x - 1$  in order to obtain a

Rational approximation (Padé approximant) for the square root of ‘x’  $\frac{1}{\sqrt{x}} \approx \frac{P(x)}{Q(x)}$  with

$P(x)$  and  $Q(x)$  Polynomials of degree ‘m’ and ‘n’ respectively so the logarithmic integral becomes a more tractable integral that can be regularized by using (2)

$$\int_c^\infty \frac{dx}{\sqrt{x}} \frac{x^{-s}}{\sqrt{x}} \approx \int_c^\infty \frac{dx}{x^{1/2+s}} \left( \frac{P(x)}{Q(x)} - \sum_{j=-1}^{m-n} c_j x^j \right) + \int_c^\infty \frac{dx}{x^{1/2+s}} \left( \sum_{j=-1}^{m-n} c_j x^j \right) \quad (30)$$

Due to the factor  $\sqrt{x}$ , the Riemann the zeta function terms  $\zeta\left(j - \frac{1}{2}\right)$  with  $j=-1, 0, 1, \dots$  will not include the pole at  $s=1$  when we use the formule (2) and (3) to regularize the divergent integrals  $\int_c^\infty x^{m-1/2-s} dx$ , the set of  $\{c_j\}$  is chosen so the first integral inside (30)

is finite for every  $\text{Re}(s) > 0$ , in order to regularize the second divergent integral we will need formula (2), ‘c’ is chosen in order the Polynomial  $Q(x)$  has no roots on  $[c, \infty)$ .

For other kind of logarithmic divergent integrals like  $\int_c^\infty \frac{\log^k(p^2 - m^2)}{p} dp$ , differentiation

with respect  $-m^2$  will make it convergent at the expense of introducing a new parameter (constant) ‘u’, see [12] Zeidler for further details on the calculation of UV integrals and Zeta regularization.

## CONCLUSIONS AND FINAL REMARKS:

We have extended the definition of the zeta regularization of a series to apply it to the

Zeta regularization of a divergent integral  $\int_0^\infty x^m dx$   $m > 0$  by using the Zeta

regularization technique combined with the Euler Maclaurin summation formula. For a good introduction to the Zeta regularization techniques, there is the book by Elizalde

[4] or the Book by Brendt based on the mathematical discoveries of Ramanujan and its method of summation equivalent to the Zeta regularization algorithm [2] , another good reference (but a bit more advanced) is Zeidler [12] , for the case of Zeta-regularized determinants [7] is a good online reference describing also the process of Zeta regularization via analytic continuation and how it can be applied to prove the identity

$\prod_{n=0}^{\infty} (n+1) = \log \sqrt{2\pi}$  . Apparently there is a contradiction, since the Riemann Zeta function has a pole at  $s=1$  so the Harmonic series could not be regularized, however using the definition of a functional determinant  $\prod_{n=0}^{\infty} \frac{E_n}{\mu}$   $E_n = n+a$  one gets the finite

result for the Harmonic (generalized) series  $\sum_{n=0}^{\infty} \frac{1}{n+a} = -\frac{\Gamma'(a)}{\Gamma(a)}$ , with the aid of the Euler-maclaurin summation formula this result for the Harmonic series can be used to give an approximate regularized value of the logarithmic integral  $\int_0^{\infty} dx \frac{1}{x+a}$  , for the

case of other types of divergent integrals  $\int_0^{\infty} dx (x+a)^m$  we can use again Euler-

Maclaurin summatio formula to express this divergent integrals in terms of the negative values of the Hurwitz or Riemann Zeta function  $\zeta_H(s,1) = \zeta(s)$   $\zeta_H(-m,1)$  (UV)  $m=0,1,2,3,4,\dots$  and the value of the derivative of Hurwitz zeta function along  $s=0$   $\partial_s \zeta_H(0,a)$  (logarithmic UV), these values encode the UV divergences [11] . For the

case of the IR (infrared ) divergences in the form  $\int_0^{\infty} \frac{dx}{x^{m-s}}$  one could make a change of

variable  $x \rightarrow \frac{1}{q}$  to re-interpretate these integrals as  $\int_0^{\infty} q^{m-2-s} dq$  for the case  $m=1$  we

have a logarithmic divergence both at  $x=0$  and as  $x \rightarrow \infty$  so we must split the integral into a IR and an UV divergent part  $\int_0^{\infty} \frac{dx}{x} = \int_0^{1/a} \frac{dx}{x} + \int_{1/a}^{\infty} \frac{dx}{x}$  after a few simple calculations

this integral will be equal to  $\int_0^{\infty} \frac{dx}{x} = 2 \log \mu$  , since we can simply introduce a formal UV

and IR regulator so  $\lim_{\Lambda \rightarrow \infty} \int_{\Lambda^{-1}}^{\Lambda} \frac{dx}{x} = 2 \log(\Lambda_{UV})$ , an UV regulator is introduced to ensure

that the integral will be convergent . We also believe that a similar procedure can be applied to extend our Zeta regularization algorithm to multiple (multi-loop) integrals  $\int d^4 q_1 \int d^4 q_2 \dots \int d^4 q_n F(q_1, q_2, \dots, q_n)$ , one of the main advantages of this algorithm is that the dimension of the space does not appear explicitly so our method does not have the same problems as dimensional regularization, and can be used when the Dirac matrices  $\gamma_5 = i\gamma_0\gamma_1\gamma_2\gamma_3$  appear . The imposition in formula (2) that ‘a’ must be a natural number is in order to avoid oddities in the process of Zeta regularization with the Zeta and Hurwitz Zeta function, since unless ‘a’ is a positive integer the equality

$$\zeta(-1, a) = \sum_{n=0}^{\infty} (n+a) \neq \sum_{n=0}^{\infty} n + \sum_{n=0}^{\infty} a = \frac{-1}{12} - \frac{a}{2}$$
 does not hold

## APPENDIX A: REGULARIZATION METHODS AND EXPANSIONS

In order to calculate a divergent integral  $\int^{\Lambda} dx f(x) \approx \Lambda^k$ , for any real and positive 'k' we must expand the integrand into a Laurent series of positive and negative terms

$$(x+a)^{-s} f(x) = \sum_{j=1}^{\infty} \frac{c_j}{(x+a)^{j+s}} + \sum_{m \geq 0} c_m (x+a)^{m-s} \quad c_n = \frac{1}{2\pi i} \int_C \frac{dz}{(z+a)^{n+1}} f(z) \quad (\text{A.1})$$

For this purpose the Binomial theorem and the expansion of the logarithm should be useful, in order to find this Laurent expansion (in general for our divergent integrals there will be only a finite number of terms since the integrals will diverge at most as a power of the regulator  $\Lambda$ ).

$$(a+x)^r = \sum_{u=0}^{\infty} \binom{r}{u} x^u a^{r-u} \quad \log(p^2 + m^2) \approx 2 \log p + 1 + \sum_{n=1}^{\infty} \frac{m^{2n} (-1)^n}{p^{2n}} \quad (\text{A.2})$$

Once we have expanded the function  $f(x)$  we may use formula (2) to regularize the divergent integrals  $\int_a^{\infty} x^{m-s} dx$  for 'm' integer or real, the case  $m = -1$  is just the

logarithmic integral, and can be regularize by using the definition of Functional determinant (infinite products of all the positive integers) so  $\sum_{n=0}^{\infty} \frac{1}{n+a} = -\frac{\Gamma'(a)}{\Gamma(a)}$ , in any

case if the integral is only logarithmic divergent  $\int^{\Lambda} dx f(x, a) \approx \log \Lambda$ , formal

differentiation with respect to external parameter 'a' will make it convergent, in this case we must introduce a new parameter inside our theory  $c_a$ .

This regularization model can be applied to almost any divergent integral that appears in Quantum Field theory, using the trick of 'Wick rotation' for example with the integral

$$\int_{-\infty}^{\infty} dE \int_{R^3} \frac{d^3 p}{(E^2 - |p|^2 c^2 - m^2 c^4)^2} \rightarrow \int_{-i\infty}^{i\infty} dE \int_{R^3} \frac{d^3 p}{(E^2 - |p|^2 c^2 - m^2 c^4)^2} = i \int_{-\infty}^{\infty} \frac{d^4 q}{(q^2 + m^2)^2} \quad (\text{A.3})$$

The main idea of this Wick rotation is to change to 'imaginary' energies by replacing the real axis by the imaginary one and then make the substitution  $E = iq_0$  and  $cp_i = q_i$  with  $i=1,2,3$ , (for more explanation see Zeidler [12]) the last integral in (A.3) is divergent but using a simple change to polar coordinates, it can be turned into the one dimensional (divergent) integral  $i \int_0^{\infty} \frac{x^3 dx}{(x^2 + m^2)^2}$ , this is still divergent, but can be

regularized using the technique of zeta regularization for integrals introduced in this paper.



For multiple integrals  $F(q_1, q_2, \dots, q_n)$  we can make a change of variable to n-dimensional polar (spherical) coordinates, and replace the integral over the angular variables  $\int d\Omega$  by a sum  $\int_0^\infty dr \sum_{\Omega} r^{n-1} F(r, \theta_1, \theta_2, \dots, \theta_{n-1})$ , this sum will clearly

depend only in the variable 'r', expanding each term into a Laurent series on powers of (r+a) for some positive 'a' we can regularize multiple integrals in an approximate way, (note that replacing an integral by a sum is legitimate and there are many Numerical Quadrature methods for this purpose) even in the case that the integrand is not invariant under rotations (unlike the dimensional regularization method or other methods).

For the case of integrals that have also an IR divergence, we can define several parameters  $\mu^2 > 0$   $\alpha^2 > 0$ , in this case the expressions  $\frac{1}{p_1^2 + \alpha^2 p_2^2}$ ,  $\frac{1}{p^m + \alpha^2 + 1}$  and

$\left((p_1 - p_2)^2 + \mu^2\right)^{-1}$  are free of IR divergencies, after regularizing the integral to cure the UV divergencies, we can make the analytic continuation in our parameters to the cases  $\mu^2 = i\varepsilon - m^2$  and  $\alpha^2 = -1 + i\varepsilon$  for a small parameter  $\varepsilon \rightarrow 0$ , so the IR divergencies appear. Another useful formula is

$\int_0^\infty \frac{dp}{(p^2 + i\varepsilon - m^2)^{s+1}} = \frac{1}{2\Gamma(s+1)} \frac{\partial^s}{\partial(m^2)^s} \left( \frac{\pi i}{\sqrt{m^2}} \right)$ , for

example by division of Polynomials integrals of the form  $\int_0^\infty \frac{p^k dp}{p^2 - m^2}$  can be reduced to

the above integrals plus some UV divergent integrals like  $\int_0^\infty \frac{dp}{p+m}$ . Another brute force

alternative is to consider the general propagator  $\left((p_1 - p_2)^2 - m^2\right)^\lambda$ , whenever lambda takes the value -1 this integral has an IR divergence at  $p_1 = p_2 \pm m$  but for lambda positive this integral has no IR divergences, the same would hold for  $(p^2 + i\varepsilon)^\lambda$  that is IR divergent for negative values of parameter lambda, we can use our zeta

regularization algorithm, by expanding the quantities  $(p^2 + i\varepsilon)^\lambda$   $\left((p_1 - p_2)^2 - m^2\right)^\lambda$

with  $\lambda \in (-2, 2)$  into a convergent Laurent series of powers  $\sum_{m=-\infty}^\infty \frac{c_m(\lambda)}{(x+a)^m}$  and then set

$\lambda = -1$  after having regularized the divergent integrals by means of the zeta function  $\zeta(s - m - 2\lambda)$  in order to obtain finite UV results.

For the case of fractional derivatives and IR divergences, we can always define a 3/2 fractional derivative relating UV and IR divergences in the form

$\frac{2}{\sqrt{\pi}} \left( \frac{1}{\varepsilon} - \gamma \right) \frac{d^{3/2-\varepsilon} \sqrt{\Delta + \lambda}}{d\lambda^{3/2-\varepsilon}} \Big|_{\lambda=0} = \frac{1}{\Delta^{1-\varepsilon}}$  with  $\Delta(q_1, q_2, \dots, q_n)$  a function of n-

variables whose zeros are precisely the IR divergences in our theory.

For a more complicated IR divergent integral, if we could use the 'sector method' so we can isolate the divergences in the form

$$\int_0^1 dx_1 \dots \int_0^1 dx_n \frac{f(x_1, \dots, x_n)}{(K(x_1, \dots, x_n))^{\alpha + \beta \varepsilon}} \rightarrow \int_0^1 dy_1 \dots \int_0^1 dy_n \frac{g(y_1, \dots, y_n)}{(C + P(y_1, \dots, y_n))^{\alpha + \beta \varepsilon}} y_1^{-(a_1 + b_1 \varepsilon)} \dots y_n^{-(a_n + b_n \varepsilon)} \quad (\text{A.4})$$

With K and P polynomials of several variables ‘C’ is a constant and f and g are smooth functions near the origin, the coefficients  $\{a_i + b_i \varepsilon\} \geq 0$  so we can factorize the IR divergences as  $\varepsilon \rightarrow 0$ , with a simple change of variable  $y_i = \frac{1}{1 + q_i}$  for  $i=1,2,3,4,\dots,n$  the last integral in (A.4) becomes an UV divergent integral

$$\int_0^\infty dq_1 \dots \int_0^\infty dq_n (1 + q_1)^{-(a_1 + b_1 \varepsilon - 2)} \dots (1 + q_n)^{-(a_n + b_n \varepsilon - 2)} \frac{g((1 + q_1)^{-1}, \dots, (1 + q_n)^{-1})}{(C + P((1 + q_1)^{-1}, \dots, (1 + q_n)^{-1}))^{\alpha + \beta \varepsilon}} \quad (\text{A.5})$$

Although we have only studied multiple integrals, the same can be applied to Fourier transform, for example if the function inside the Fourier transform depends only on the modulus of position vector  $f(\vec{r})$  then we can make a change of variable to polar

coordinates to get  $\int_{R^n} dr f(\vec{r}) e^{i\vec{k} \cdot \vec{r}} \approx \sum_i \int_0^\infty dr f(r) r^{n-1} g(\theta_i) e^{ikr \cos \theta_i}$ , here the angle is

introduced by the scalar product  $\vec{k} \cdot \vec{r} = k.r \cos \theta$ , the function ‘g’ will depend only on the angle  $\theta$ , so we may replace a multiple Fourier transform by an approximate sums of Fourier transform in one dimension for the function  $f(r)r^{n-1}$

## References

- [1] Abramowitz, M. and Stegun, I. A. (Eds.). "Riemann Zeta Function and Other Sums of Reciprocal Powers." §23.2 in *"Handbook of Mathematical Functions"*. New York: Dover, pp. 807-808, 1972.
- [2] Berndt B. "Ramanujan's Notebooks: Part II" Springer; 1 edition (1993) ISBN-10: 0387941096
- [3] Dowling Jonathan P, "The Mathematics of the Casimir Effect", *Math. Mag.* **62**, 324-331 (1989).
- [4] Elizalde E. ; "Zeta-function regularization is well-defined", *Journal of Physics* **27** (1994), L299-304.
- [5] Elizalde E. ; "Zeta regularization techniques with applications" Ed. World Scientific Press ISBN 981-02-1441-3
- [6] Mizuno, Y. "Generalized Lerch Formulas: Examples of Zeta-Regularized Products." *J. Number Th.* **118**, 155-171, 2006.

- [7] Muñoz García, E. and Pérez Marco, R. "The Product Over All Primes is  $4\pi^2$ ." Preprint IHES/M/03/34. May 2003.  
<http://inc.web.ihes.fr/prepub/PREPRINTS/M03/Resu/resu-M03-34.html>.
- [8] Pavel E "Note on dimensional regularization", *Quantum fields and strings: a course for mathematicians, Vol. 1*, (Princeton, NJ, 1996/1997), Providence, R.I.: Amer. Math. Soc., pp. 597–607, ISBN 978-0-8218-2012-4 (1999)
- [9] Prellberg Thomas, " *The mathematics of the Casimir Effect* " a talk available at e-print at <http://www.maths.qmul.ac.uk/~tp/talks/casimir.pdf>
- [10] Voros A. " *Zeta functions over zeros of zeta functions* " Ed. SPRINGER (2010) ISBN: 978-3-642-05202-6
- [11] Zee, A. (2003). " *Quantum Field Theory in a Nutshell* . "Princeton University ISBN ISBN 0-691-01019-6.
- [12] Zeidler E. " *Quantum Field Theory I : Basis in Physics and Mathematics* " Ed : Springer Verlag ISBN: 3540347623