

# About incorrect application of the equations in cosmology (Neumann-Seeliger paradox and the Big Bang theory)

Valery Zhukov <sup>1)</sup>

## Abstract

*The application of the Poisson's equation solutions in the Neumann-Seeliger paradox for an infinite universe is incorrect. Also it is incorrect to use the metric and all the equations of the Big Bang theory for distances greater than the age of the universe, i.e. outside the event horizon for any point of the universe.*

The Neumann-Seeliger paradox is a paradox of the classical Newton's theory of gravitation and was one of the impetuses to the development of the Big Bang theory.

Neumann-Seeliger paradox:

***In an infinite Universe with Euclidean geometry and non-zero average density of matter, the gravitational potential has an indefinite value at any point.***

If the density of matter  $\rho$  is randomly distributed in space, in the classical theory the gravitational field created by it is determined by the gravitational potential  $\varphi$ . To find this potential, you have to solve the Poisson's equation

$$\Delta\varphi = -4\pi G\rho \quad (1)$$

Here  $G$  is a gravitational constant. The general solution of this equation can be written as:

$$\varphi = -G \int_V \frac{\rho dV}{r} + C \quad (2)$$

**It was found that if the average density of matter in the Universe is non-zero, the integral diverges.**

Let us consider the Poisson's equation. Gravitational potential, by definition, is a scalar function, which is numerically equal to the work the field produces when transferring unit mass from any starting point to a given point. A more general solution of Equation (2) for the body with a radius of  $r_0$ , where  $R$  is a distance to the center of the body ( $R > r_0$ ):

$$\varphi = \varphi(R) - G \int_0^{r_0} \frac{\rho dV}{r} \quad (3)$$

When  $r_0 = \infty$ , the application of the equation is incorrect, as an infinite Universe has no boundaries and the potential at the point  $R$ , with respect to which the calculation is made, can not be determined. Point  $R$  does not belong to the considered universe. And the potential between any two points belonging to the considered universe is zero and has an infinite value.

For a limited static universe, Equation (3) is correct.

But for a limited dynamic universe, the event horizon for each point of the universe is no more than the age of the universe. According to the current estimates of the Big Bang theory, the size of the universe is 160 billion. light-years, the age of the universe 14 billion. light-years, and the application of Equation (3) in terms of the special theory of relativity is incorrect.

It should be noted that the use of all the equations of the Big Bang theory in the context of the special theory of relativity can be correct only for distances of no more than 14 billion. light-years, the rest of space is outside the event horizon. Negative time (before the Big Bang) in the Big Bang theory is not considered. I.e. in this case, as well as in the Neumann-Seeliger paradox, the equations operate on the indefinite space within this theory (beginning from the use in the Friedmann equations of mass and size of the universe).

---

<sup>1)</sup> valeryzhukov@mail.ru / Valery Zhukov, Netanya, Israel