

## On Improper Integrals

Anamitra Palit

palit.anamitra@gmail.com

### Abstract

The writing intends to point out aspects of conflict regarding some standard improper integrals

### Introduction

Two standard integrals frequently used in physics have been considered and the results have been analyzed to bring out some conflicting aspects

### Section I

We consider the standard integral<sup>[1]</sup>

$$I = \int_{-\infty}^{+\infty} \frac{dk_0}{k^2 + s - i\varepsilon} \quad (1)$$

$$I = \int_{-\infty}^{+\infty} \frac{dk_0}{k_0^2 - |\vec{k}|^2 + s - i\varepsilon} = \frac{i\pi}{\sqrt{|\vec{k}|^2 + s}}$$

$$I = \int_{-\infty}^{+\infty} \frac{dk_0}{k_0^2 - A^2 - i\varepsilon}; A^2 > 0 \quad (2)$$

While integration with respect to  $k_0$  the variable  $|\vec{k}|^2$  is held constant[asides s]

When  $A^2 = |\vec{k}|^2 - s > 0$

$$I = \int_{-\infty}^{+\infty} \frac{dk_0}{k_0^2 + A^2 - i\varepsilon}; A^2 > 0 \quad (3)$$

We evaluate (2) and (3) ignoring the complex part

Evaluation of (2'), ignoring the imaginary part:

We evaluate the following improper integral by using limit concepts :

$$I = \int_{-\infty}^{+\infty} \frac{1}{x^2 - a^2} dx \quad (2')$$

Indefinite integral

$$\int \frac{1}{x^2 - a^2} dx = \ln \frac{x - a}{x + a}$$

The integral represented by (2') may be interpreted as

$$\begin{aligned} I &= \frac{1}{2a} \lim_{q \rightarrow s, M \rightarrow \infty} \left[ \left[ \ln \frac{x - a}{x + a} \right]_{-M}^{-q} + \left[ \ln \frac{x - a}{x + a} \right]_{-q}^{+q} + \left[ \ln \frac{x - a}{x + a} \right]_q^M \right] \\ &= \frac{1}{2a} \lim_{q \rightarrow s, M \rightarrow \infty} \left[ \ln \left| \frac{-q - a}{-q + a} \right| - \ln \left| \frac{-M - a}{-M + a} \right| + \ln \left| \frac{q - a}{q + a} \right| - \ln \left| \frac{-q - a}{-q + a} \right| + \ln \left| \frac{M - a}{M + a} \right| - \ln \left| \frac{q - a}{q + a} \right| \right] \\ &= \frac{1}{2a} \lim_{q \rightarrow s, M \rightarrow \infty} \left[ \ln \frac{|q + a|}{|q - a|} - \ln \frac{|M + a|}{|M - a|} + \ln \frac{|q - a|}{|q + a|} - \ln \frac{|q + a|}{|q - a|} + \ln \frac{|M - a|}{|M + a|} - \ln \frac{|q - a|}{|q + a|} \right] \\ &= \frac{1}{2a} \lim_{q \rightarrow s, M \rightarrow \infty} \left[ \ln \frac{|q + a|}{|q - a|} - \ln \frac{|q + a|}{|q - a|} + \ln \frac{|q - a|}{|q + a|} - \ln \frac{|q - a|}{|q + a|} + \ln \frac{|M - a|}{|M + a|} - \ln \frac{|M + a|}{|M - a|} \right] \\ &= \frac{1}{2a} \lim_{M \rightarrow \infty} \ln \frac{|M - a|}{|M + a|} - \frac{1}{2a} \lim_{M \rightarrow \infty} \frac{|M + a|}{|M - a|} \\ &= \frac{1}{2a} [\ln 1 - \ln 1] = 0 \end{aligned}$$

$$\int_{-\infty}^{+\infty} \frac{1}{x^2 - a^2} dx ; a^2 > 0 \quad (4)$$

Next we pass on to the evaluation of

$$\int_{-\infty}^{+\infty} \frac{1}{x^2 + a^2} dx ; a^2 > 0 \quad (5)$$

The indefinite integral

$$\int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \text{Sin}^{-1} \frac{x}{a} + C$$

Since the integrand an even function and positive everywhere on the x-axis

$$\int_{-\infty}^{+\infty} \frac{1}{x^2 + a^2} dx = 2 \int_0^{+\infty} \frac{1}{x^2 + a^2} dx \rightarrow \infty \quad (5)$$

[The indefinite integral, in fact, is not required to come to this conclusion since we know that the integrand is positive everywhere on the x axis]

## Section II

Standard result<sup>[2]</sup>

$$\begin{aligned}
I &= \int \frac{d^4k}{(2\pi)^4} \frac{1}{(k^2 + ns - i\varepsilon)^3} = \frac{i}{32\pi^2 ns} \quad (4) \\
I &= \int \frac{d^4k}{(2\pi)^4} \frac{1}{(k^2 + ns - i\varepsilon)^3} \\
&= \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns + i\varepsilon)^3}{(k^2 + ns - i\varepsilon)^3 (k^2 + ns + i\varepsilon)^3} \\
&= \int \frac{d^4k}{(2\pi)^4} \frac{[(k^2 + ns) + i\varepsilon]^3}{[(k^2 + ns)^2 + \varepsilon^2]^3} \\
&= \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)^3 - i\varepsilon^3 + 3i\varepsilon(k^2 + ns)(k^2 + ns + i\varepsilon)}{[(k^2 + ns)^2 + \varepsilon^2]^3} \\
&= \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)^3 - i\varepsilon^3 + 3i\varepsilon(k^2 + ns)^2 - 3\varepsilon^2(k^2 + ns)}{[(k^2 + ns)^2 + \varepsilon^2]^3} \\
&= \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)^3 - 3\varepsilon^2(k^2 + ns)}{[(k^2 + ns)^2 + \varepsilon^2]^3} - i \int \frac{d^4k}{(2\pi)^4} \frac{\varepsilon^3 - 3\varepsilon(k^2 + ns)^2}{[(k^2 + ns)^2 + \varepsilon^2]^3} = \frac{i}{32\pi^2 ns} \\
I_1 &= \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)^3 - 3\varepsilon^2(k^2 + ns)}{[(k^2 + ns)^2 + \varepsilon^2]^3} = 0; I_2 = \int \frac{d^4k}{(2\pi)^4} \frac{\varepsilon^3 - 3\varepsilon(k^2 + ns)^2}{[(k^2 + ns)^2 + \varepsilon^2]^3} = \frac{i}{32\pi^2 ns} \\
I_1 &= \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)^3 - 3\varepsilon^2(k^2 + ns)}{[(k^2 + ns)^2 + \varepsilon^2]^3} \\
&= \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)^3}{[(k^2 + ns)^2 + \varepsilon^2]^3} - 3\varepsilon^2 \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)}{[(k^2 + ns)^2 + \varepsilon^2]^3} \\
I_2 &= \int \frac{d^4k}{(2\pi)^4} \frac{\varepsilon^3 - 3\varepsilon(k^2 + ns)^2}{[(k^2 + ns)^2 + \varepsilon^2]^3} = \varepsilon \int \frac{d^4k}{(2\pi)^4} \frac{\varepsilon^2 - 3(k^2 + ns)^2}{[(k^2 + ns)^2 + \varepsilon^2]^3}
\end{aligned}$$

Calculations based on  $I_1$ 

$$I_1 = \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)^3}{[(k^2 + ns)^2 + \varepsilon^2]^3} - 3\varepsilon^2 \int \frac{d^4k}{(2\pi)^4} \frac{(k^2 + ns)}{[(k^2 + ns)^2 + \varepsilon^2]^3} = 0$$

For  $\epsilon \rightarrow 0$ ,

$$I_1 = \int \frac{d^4k}{(2\pi)^4} \frac{1}{(k^2 + ns)^3}$$

Since

$I_1 = 0$  we have for  $\epsilon \rightarrow 0$

$$\int \frac{d^4k}{(2\pi)^4} \frac{1}{(k^2 + ns)^3} = 0 \quad (A)$$

Differentiating (A) with respect to  $s$  we have

$$\int \frac{d^4k}{(2\pi)^4} \frac{1}{(k^2 + ns)^4} = 0 \quad (B)$$

Calculations based on  $I_2$

$$I_2 = \epsilon \int \frac{d^4k}{(2\pi)^4} \frac{\epsilon^2 - 3(k^2 + ns)^2}{[(k^2 + ns)^2 + \epsilon^2]^3}$$

Asides the fact that  $\epsilon \rightarrow 0$  we have the additional strength of (B)

For  $\epsilon \rightarrow 0$  [and recalling (B)]

$$I_2 = \epsilon \int \frac{d^4k}{(2\pi)^4} \frac{\epsilon^3 - 3\epsilon(k^2 + ns)^2}{[(k^2 + ns)^2 + \epsilon^2]^3} = -3\epsilon \times \int \frac{d^4k}{(2\pi)^4} \frac{1}{(k^2 + ns)^4} = 0 \neq \frac{i}{32\pi^2 ns}$$

Asides the fact that  $\epsilon \rightarrow 0$  we have the additional fact that (B) does not tend to infinity in which case there would have been a possibility of the integral becoming convergent. On the contrary it evaluates to zero with  $\epsilon \rightarrow 0$ .

$I_2$  does not work out to its standard value as given by (4)

### Conclusion

As claimed, we have arrived at some conflicts with the two the standard integrals

### References

1. Sakurai J. J., Advanced Quantum Mechanics, Pearson Education, India, Appendix E,p327
2. Sakurai J. J., Advanced Quantum Mechanics, Pearson Education, India, Appendix E,p327

