

# Chen's Formulas of the Fine-structure Constant

Gang Chen<sup>†</sup>, Tianman Chen and Tianyi Chen

*Guangzhou Huifu Research Institute Co., Ltd., Guangzhou, P. R. China*

*7-20-4, Greenwich Village, Wangjianglu 1, Chengdu, P. R. China*

<sup>†</sup>Correspondence to: [gang137.chen@connect.polyu.hk](mailto:gang137.chen@connect.polyu.hk)

Dedicated to Prof. Albert Sun-Chi Chan on the occasion of his 70<sup>th</sup> birthday

## Abstract

This paper gives two series of formulas of the fine-structure constant  $\alpha$  which are reasonable, precise, smart and elegant. It also demonstrates there are two values of  $\alpha$ , i.e.,  $\alpha_1=1/137.035999037435$  and  $\alpha_2=1/137.035999111818$ , which are consistent with but much more accurate than those experiment measured values. The formulas consist of  $2\pi$ -e formulas and some factors related to nucleon numbers of nuclides. A brief explanation of the fine-structure constant shows  $1/\alpha \approx 137.036$  is the equal ratio factor between 112 and 168 (more precisely  $168-1/3$ ). Based on these, all 119<sup>th</sup> to 170<sup>th</sup> ideal extended elements were predicted, the speed of light in atomic units was mathematically calculated by  $c_{au}=1/(\alpha_1\alpha_2)^{1/2}=137.035999074627$ , Schrödinger equation of hydrogen atom was simplified and correlated with  $\alpha_1/\alpha_2$ , classical electron radius was calculated to be 2.81794032658(43) fm and proton charge radius was hypothetically calculated to be 0.833027202999(13) fm. In the end, it was found that the approximate rational numbers of  $2\pi$  marvelously related to nuclides, a mathematic shell model of nuclides was established and a picture of elements and ideal extended elements was depicted.

**Keywords:** formulas; the fine-structure constant; the ideal extended elements; the speed of light; Schrödinger equation of hydrogen atom; the proton charge radius;  $2\pi$ .

## 1. Introduction

The fine-structure constant (Sommerfeld constant) is a critical dimensionless constant in physics, it is a century mystery of physics, it has been one of the biggest enigmas in physics since it was introduced by Arnold Sommerfeld in 1916. Its definition, some interpretations and the latest measured values are as follows<sup>1,2</sup>:

$$\alpha = \frac{\lambda_e}{2\pi a_0}, \quad \alpha = \frac{2\pi r_e}{\lambda_e}, \quad \frac{a_0}{r_e} = \frac{1}{\alpha^2}; \quad \alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} = \frac{v_e}{c}, \quad \frac{c}{v_e} = \frac{1}{\alpha}$$

in atomic units, the speed of light  $c_{au} = \frac{1}{\alpha}$

the 2014 CODATA recommended value:  $\alpha = 1/137.035999139(31)$

the 2018 CODATA recommended value:  $\alpha = 1/137.035999084(21)$

Science 13 April 2018 reported value:  $\alpha = 1/137.035999046(27)$

The ratio of Bohr radius of hydrogen atom  $a_0$  to the classical electron radius  $r_e$  is  $1/\alpha^2$ . The ratio of the speed of light  $c$  to the line velocity of ground state electron in hydrogen atom  $v_e$  is  $1/\alpha$ , this means in atomic units  $c=1/\alpha$  and  $E=mc^2=m/\alpha^2$  or  $\alpha^2=m/E$ . In quantum electrodynamics it substantially characterizes the strength of electromagnetic interaction between elementary charged particles such as electron and proton, so it is the coupling constant of electric charges. It is one of the 25 fundamental constants (could not be calculated theoretically, could only be determined by experiments) in Standard Model of physics and should be the most important one. As it is dimensionless, it could be called the proportional ruler of the nature or the bridge of mathematics and physics. However, to our knowledge, up to now (except this work), no one knows how it comes from, no one could give reasonable explanations to it or formulas of it since it was introduced.

In 2016 Paul Davis gave the following comment<sup>3</sup>: “Physicists have long wondered where this number,  $1/137.035999$ , comes from. Is there a deep reason why  $\alpha$  has to be precisely this number for the world to function as it does? There is a long history of attempts to derive  $\alpha$  from physical theory or to concoct a mathematical formula that has this value. For a brief time in the 1920s, when it looked as if  $\alpha$  might be exactly  $1/137$ , astronomer Arthur Eddington searched for a theory that would throw up the numbers naturally, but his ideas ultimately led nowhere. Then in 1969 a young Swiss mathematician, Armand Wyler, pointed out that  $(9/16\pi^3)(\pi/5!)^{1/4}$  comes close to  $1/137.036$ , which matched the value of  $\alpha$  to the precision known at the time. However, his formula was not accompanied by any credible theory and was regarded as little more than a numerical curiosity. Several other attempts at  $\alpha$  numerology have been made since, none of which have gained traction in the physics community.”


As for the fascination of the fine-structure constant, in the middle of 1980s, Richard Feynman stated<sup>4</sup>: “It has been a mystery ever since it was discovered more than fifty years ago, and all good theoretical physicists put this number up on their wall and worry about it. Immediately you would like to know where this number for a coupling comes from: is it related to pi or perhaps to the base of natural logarithms? Nobody knows. It's one of the greatest damn mysteries of physics: a magic number that comes to us with no understanding by man. You might say the hand of God wrote that number, and we don't know how He pushed his pencil.”

This paper shows how God pushed his pencil to write the fine-structure constant and how God used it to coordinate elements.


## 2. $2\pi$ -e formula(s)

$2\pi$ -e formula, its related formulas and their preliminary applications were deduced independently by us from April to December of 2013.

**Fig. 1. Diagram of  $y=1/x$ .**



**Fig. 2. Diagram of  $y=\log(x)$ .**



$$\text{Euler-Mascheroni constant } \gamma : \sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{\infty} = \ln \infty + \gamma$$

$$\text{As for } y = 1/x \text{ (Fig. 1), } \gamma = 0.577215\dots = 0.5 + 0.077215\dots = \sum_{n=1}^{\infty-1} \delta_{2n} + \sum_{n=1}^{\infty-1} \delta_{1n} = \frac{1}{2} + \gamma_1$$

$$\gamma_1 = \sum_{n=1}^{\infty-1} \delta_{1n} = \lim_{N \rightarrow \infty} \left( \sum_{n=1}^{N-1} \frac{1}{n} - \int_1^N \frac{1}{x} dx \right) - \frac{1}{2}, \text{ Generally } \gamma_s = \lim_{N \rightarrow \infty} \left( \sum_{n=1}^{N-1} \frac{1}{n^s} - \int_1^N \frac{1}{x^s} dx \right) - \frac{1}{2}, \quad s \in \mathbb{N}$$

$$\begin{aligned} \text{As for } y = \log(x) \text{ (Fig. 2), } \delta_{1,n} &= \int_n^{n+1} \ln x dx - \frac{1}{2} \ln \frac{n+1}{n} - \ln n = (x \ln x - x) \Big|_n^{n+1} - \frac{1}{2} \ln(n+1)n \\ &= (n+1) \ln(n+1) - n \ln n - 1 - \frac{1}{2} \ln(n+1) - \frac{1}{2} \ln n = \left(n + \frac{1}{2}\right) \ln\left(1 + \frac{1}{n}\right) - 1 \end{aligned}$$

$$\gamma_{c,N} = \sum_{n=1}^N \delta_{1,n} = \sum_{n=1}^N \left[ \left(n + \frac{1}{2}\right) \ln\left(1 + \frac{1}{n}\right) - 1 \right] = \sum_{n=1}^N \ln \frac{\left(1 + \frac{1}{n}\right)^{\left(n + \frac{1}{2}\right)}}{e} = \ln \prod_{n=1}^N \frac{\left(1 + \frac{1}{n}\right)^{\left(n + \frac{1}{2}\right)}}{e}$$

$$\gamma_c = \gamma_{c,\infty} = \sum_{n=1}^{\infty} \delta_{1,n} = \lim_{N \rightarrow \infty} \left( \int_1^{N+1} \log(x) dx - \sum_{n=1}^N \log(n) - \frac{\log(N+1)}{2} \right)$$

$$= \sum_{n=1}^{\infty} \left[ \left(n + \frac{1}{2}\right) \ln\left(1 + \frac{1}{n}\right) - 1 \right] = \sum_{n=1}^{\infty} \ln \frac{\left(1 + \frac{1}{n}\right)^{\left(n + \frac{1}{2}\right)}}{e} = \ln \prod_{n=1}^{\infty} \frac{\left(1 + \frac{1}{n}\right)^{\left(n + \frac{1}{2}\right)}}{e}$$

$$\ln N! = \sum_{n=1}^N \ln n = \int_1^{N+1} \ln x dx - \sum_{n=1}^N \delta_{1,n} - \sum_{n=1}^N \delta_{2,n} = (x \ln x - x) \Big|_1^{N+1} - \ln e^{\gamma_{c,N}} - \sum_{n=1}^N \frac{\ln(n+1) - \ln n}{2}$$

$$= (N+1) \ln \frac{(N+1)}{e} + \ln \frac{e}{e^{\gamma_{c,N}}} - \frac{1}{2} \ln(N+1) = \ln \left[ \frac{e^{1-\gamma_{c,N}}}{\sqrt{N+1}} \left( \frac{N+1}{e} \right)^{(N+1)} \right]$$

$$N! = \frac{e^{1-\gamma_{c,N}}}{\sqrt{N+1}} \left( \frac{N+1}{e} \right)^{(N+1)}, \text{ compared to Stirling formula : } N! \sim \sqrt{2\pi N} \left( \frac{N}{e} \right)^N$$

$$(N+1)! = (N+1)N! \sim \sqrt{2\pi(N+1)} \left( \frac{N+1}{e} \right)^{N+1}, \quad N! \sim \frac{\sqrt{2\pi}}{\sqrt{N+1}} \left( \frac{N+1}{e} \right)^{N+1}$$

$$\text{Compared to previous formula, gives } \sqrt{2\pi} \sim e^{1-\gamma_{c,N}} \text{ or } 2\pi = \left( \frac{e}{e^{\gamma_c}} \right)^2$$

$$2\pi - e \text{ formula(s): } 2\pi = \left( \frac{e}{e^{\gamma_c}} \right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots, \quad (2\pi)_k = \left( \frac{e}{e^{\gamma_{c,k}}} \right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}$$

$$\gamma_c = 0.0810614668, \quad e^{\gamma_c} = 1.0844375$$

$2\pi$ -e formula is an expanding form of Stirling formula. To our knowledge, it was first deduced by us. If it was new, it could be named Chen's  $2\pi$ -e formula.

### 3. Some Formulas Related to $2\pi$ -e Formula

The following formulas which correlate each other and has similar form could be called Chen's natural group formulas, and the form is called natural group.

$$\begin{aligned}
1 &= 4\gamma_c + \frac{4\gamma_1}{1(1+1)} + \frac{4\gamma_2}{2(2+1)} + \frac{4\gamma_3}{3(3+1)} + \dots \\
&= |B| \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{|B_{2n}|(\pi/2)^{2n}}{(2n)!} = \sum_{n=1}^{\infty} \frac{|B_{2n}|\pi^{2n}}{(2n)!} = -|B| \frac{3\pi}{2} + \sum_{n=1}^{\infty} \frac{|B_{2n}|(3\pi/2)^{2n}}{(2n)!} \\
N &\sim -\frac{3}{2}|B| + \sum_{n=1}^N \frac{|B_{2n}|(2\pi)^{2n}}{2(2n)!} \\
e &= 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots \\
2\pi &= \left(\frac{e}{e^{\gamma_c}}\right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots
\end{aligned}$$

$B, B_{2n}$ : the Bernoulli numbers such as  $-\frac{1}{2}, -\frac{1}{6}, -\frac{1}{30}, \frac{1}{42}, -\frac{1}{30}, \dots$

$$\gamma_c = \lim_{N \rightarrow \infty} \left( \int_1^{N+1} \log(x) dx - \sum_{n=1}^N \log(n) - \frac{\log(N+1)}{2} \right) = 0.0810614668$$

$$\gamma_s = \lim_{N \rightarrow \infty} \left( \sum_{n=1}^{N-1} \frac{1}{n^s} - \int_1^N \frac{1}{x^s} dx \right) - \frac{1}{2}, \quad s \in \mathbb{N}$$

$\gamma_1 = 0.077215, \gamma_2 = 0.144934, \gamma_4 = 0.24899, \gamma_8 = 0.36122, \gamma_{16} = 0.433349, \dots, \gamma_{\infty} = 0.5$

$\gamma_c, \gamma_1, \gamma_2, \gamma_3, \dots$  are called Chen's natural group constants (analogue to Bernoulli numbers).

The following are some other formulas related to  $2\pi$ -e Formula.

$$\begin{aligned}
\sqrt{2\pi} &= e^{1-\gamma_c}, \quad e = \sqrt{2\pi} e^{\gamma_c} = \sqrt{2\pi} \left( 1 + \sum_{n=1}^{\infty} \frac{\gamma_c^n}{n!} \right) \\
\gamma_c &= \sum_{n=1}^{\infty} \left[ \left( n + \frac{1}{2} \right) \ln \left( 1 + \frac{1}{n} \right) - 1 \right] = \sum_{n=1}^{\infty} \frac{(2^{2n}-1)|B_{2n}|\pi^{2n} - 2(2n)!}{2(2n+1)!} = \frac{1}{4} - \sum_{s=1}^{\infty} \frac{\gamma_s}{s(s+1)} \\
\gamma_g &= \sum_{n=1}^{\infty} \left( n + \frac{1}{2} \right) \ln \left( 1 + \frac{1}{n} \right) - \int_1^{\infty} \left( x + \frac{1}{2} \right) \ln \left( 1 + \frac{1}{x} \right) dx \\
\gamma_{cg} &= \frac{1}{2} \lim_{N \rightarrow \infty} \left[ \sum_{n=1}^N \frac{(2^{2n}-1)|B_{2n}|\pi^{2n}}{(2n+1)!} - \ln N \right] \\
\frac{\pi}{2} &= \left( \frac{e}{e^{\gamma_g}} \right)^2, \quad e = \sqrt{\frac{\pi}{2}} e^{\gamma_g} = \sqrt{\frac{\pi}{2}} \left( 1 + \sum_{n=1}^{\infty} \frac{\gamma_g^n}{n!} \right); \quad \frac{\pi}{2} = \left( \frac{e^{\gamma/2}}{e^{\gamma_{cg}}} \right)^2, \quad \gamma = \ln \frac{\pi}{2} + 2\gamma_{cg} \\
\gamma_c &= \gamma_g - \ln 2 = 1 - \frac{\gamma}{2} + \gamma_{cg} - \ln 2, \quad \gamma_{cg} = \frac{1}{2} + \sum_{s=2}^{\infty} \frac{\gamma_s}{s(s+1)} - \ln 2 \\
\gamma_c &= 0.0810614668, \quad \gamma_g = 0.7742086474, \quad \gamma_{cg} = 0.0628164798 \\
\frac{\pi}{2} &= \sum_{n=1}^{\infty} \frac{|B_{2n}|\pi^{2n}}{2n(2n)!}; \quad \sum_{n=1}^{\infty} [\zeta(2n) - 1] = \frac{3}{4}, \quad \zeta(2n) = \sum_{k=1}^{\infty} \frac{1}{k^{2n}} \\
\sum_{n=1}^{\infty} \frac{1}{n} &= \sum_{n=1}^{\infty} \frac{|B_{2n}|(2\pi)^{2n}}{2n(2n)!} = \sum_{n=1}^{\infty} \frac{|B_{2n}|(2n2^{2n}+1)\pi^{2n}}{2n(2n+1)!}
\end{aligned}$$

#### 4. Some Applications of $2\pi$ -e Formula and its Related Formulas

(1).  $2\pi$ -e formula is basically an algebraic expanding of Stirling formula, but it is more meaningful, it exhibits the relationship between  $2\pi$  and e. In  $2\pi$ -e formula,  $\gamma_c$  is a real constant with geometric definition like Euler-Mascheroni constant  $\gamma$ . With  $2\pi$ -e formula and its related formulas,  $2\pi$  can be calculated from e and vice versa. So it is the real  $2\pi$ -e relationship formula.

$$2\pi = \left(\frac{e}{e^{\gamma_c}}\right)^2 = e^2 \prod_{n=1}^{\infty} \frac{e^2}{\left(1 + \frac{1}{n}\right)^{2n+1}} = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots$$

$$e = \sqrt{2\pi} e^{\gamma_c} = \sqrt{2\pi} \left(1 + \sum_{n=1}^{\infty} \frac{\gamma_c^n}{n!}\right), \quad \gamma_c = \sum_{n=1}^{\infty} \frac{(2^{2n}-1) |B_{2n}| \pi^{2n} - 2(2n)!}{2(2n+1)!}$$

(2).  $2\pi$ -e formula demonstrates  $2\pi$  is a natural constant rather than  $\pi$ .  $\pi/2$  is somewhat fundamental but not as complete as  $2\pi$ .  $\pi$  is neither fundamental nor complete. In 2001 mathematician Bob Palais said “ $\pi$  is wrong”<sup>5</sup>.  $2\pi$ -e formula and the Taylor expansion of e have similar form (natural group form), this should give a conclusive proof that  $2\pi$  is a real natural constant and  $\pi$  is not.

$$2\pi = \left(\frac{e}{e^{\gamma_c}}\right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots \Rightarrow 2\pi \text{ or } \sqrt{2\pi} \text{ is a natural constant}$$

$$\frac{\pi}{2} = \left(\frac{e}{e^{\gamma_s}}\right)^2 = \left(\frac{e^{\gamma/2}}{e^{\gamma_{cg}}}\right)^2 \Rightarrow \frac{\pi}{2} \text{ or } \sqrt{\frac{\pi}{2}} \text{ is almost a natural constant}$$

$$\pi = \left(\frac{e}{e^{\gamma_c} \sqrt{2}}\right)^2 = \left(\frac{e^{\sqrt{2}}}{e^{\gamma_s}}\right)^2 \Rightarrow \pi \text{ or } \sqrt{\pi} \text{ is not a natural constant}$$

**Table 1** lists some points of view of Piist who support  $\pi$  is a natural constant, Tauist who support  $2\pi$  is a natural constant and this work which supports the later.

**Table 1.** Comparison of points of view of Piist, Tauist and this work.

	Piist	Tauist	This work
Circumference of a circle	$\pi d$	$2\pi R$	$2\pi R$
Area of a circle	$\pi R^2$		$(1/2)(2\pi R)R$
Volume of sphere	$(4/3) \pi R^3$	$(2/3)(2\pi)R^3$	$(2\pi R^2/3)2R$
Volume of n-dimension sphere	$\frac{\pi^{n/2}}{\Gamma(n/2+1)} R^n$	$\frac{(2\pi)^{n/2}}{2^{n/2} \Gamma(n/2+1)} R^n$	$\frac{2\pi R^2}{n} V_{n-2}$
Euler's identity	$e^{i\pi} + 1 = 0$	$e^{2\pi i} = 1$	$e^{2\pi i} = 1$
Gauss integral	$\int_{-\infty}^{+\infty} e^{-x^2} dx = \sqrt{\pi}$		$\int_{-\infty}^{+\infty} e^{-x^2} dx = \frac{e}{e^{\gamma_c}} \frac{1}{\sqrt{2}}$

(3). As  $2\pi$  is a square number, the frequent appearing of its square root in some

important equations such as Gaussian distribution (normal distribution) and Maxwell–Boltzmann distribution becomes reasonable and understandable. And the distributions can be transformed as follows.

$$\text{Standard Normal Distribution: } f(x, 0, 1) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} = e^{-\frac{x^2+2(1-\gamma_e)}{2}}$$

$$\text{Maxwell–Boltzmann Distribution: } f(v) = \frac{2}{\sqrt{2\pi}} v^2 \left(\frac{m}{kT}\right)^{\frac{3}{2}} e^{-\frac{mv^2}{2kT}} = 2\left(\frac{m}{kT}\right)^{\frac{3}{2}} v^2 e^{-\left(\frac{m}{kT} \frac{v^2}{2} + 1 - \gamma_e\right)}$$

(4). Euler’s identity (Euler’s equation)  $e^{i\pi}+1=0$  is called God formula and the most beautiful formula in mathematics. However, as  $2\pi$  is the real natural constant and  $\pi$  is not,  $e^{2\pi i}=1$  should be more beautiful.

(5).  $\gamma=\ln(2\pi)+\gamma_{cg}$  may help to prove  $\gamma$  is an irrational number or even a transcendental number.

(6). The natural group formulas help us to establish “Chen’s Periodic Table of Elements and Natural Group Theory”<sup>6</sup> (2014-2017).

(7). The mathematic expression of chirality is  $\pm 2\pi$ . This concept is helpful for us to establish “Chirality and Poetry Model of Atomic Nuclei”<sup>7</sup> (2017/12-2018/3).

(8). Based on the above theories, Chen’s theory of the fine-structure constant was deduced (2018/4-6)<sup>8</sup> and has been revised, modified and improved (2018/7-2020/1).

## 5. Original Inspiration for Formulas of the Fine-structure Constant

1. According to  $\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} = \frac{\lambda_e}{2\pi a_0} = \frac{2\pi r_e}{\lambda_e} \approx \frac{1}{137.036}$ , the formulas of  $\alpha$  should relate to  $2\pi$ .

$$2. \frac{137.036}{2\pi} = \frac{137.036}{6.28318} = 21.81, \quad 137.036 = 21.81 \times 2\pi$$

$$3. \text{According to } 2\pi\text{-e formula: } 2\pi = \left(\frac{e}{e^{\gamma_e}}\right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots$$

$2\pi$  is a square number, suppose  $21.81 = x^2$ ,  $x = 4.670 \approx 14/3$

$$\text{so: } \frac{1}{\alpha} \approx \left(\frac{14}{3}\right)^2 2\pi \text{ or } \alpha \approx \left(\frac{3}{14}\right)^2 \frac{1}{2\pi} \quad (\text{Discover: about 2 am on 2018/4/12})$$

4. Apply with  $2\pi\text{-e}$  formula (in the afternoon of 2018/4/12, a meeting in the morning)

$$\alpha = \left(\frac{3}{14}\right)^2 \frac{1}{(2\pi)_{112}} = \left(\frac{3}{14}\right)^2 \frac{1}{e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} = 137.035781520, \text{ closest to the real value.}$$

As 112 is one of the most important stable numbers and the 112th element  ${}_{112}^{285}\text{Cn}^*$  is the natural end of elements according to our Chen's Chirality and Poetry Model of Atomic Nuclei<sup>6</sup>.

$$\text{So: } \textit{Eureka!} \quad \text{Subsequently transformed to: } \alpha = \frac{6^2}{7(2\pi)_{112}} \frac{1}{112} = 137.035781520,$$

$$\text{Finally modified to: } \alpha = \frac{6^2}{7(2\pi)_{112}} \frac{1}{112 + \frac{1}{75^2}} = 137.035999037435$$

## 6. Logical Deduction of Chen's Formulas of the Fine-structure Constant

Physicist Richard Feynman noticed a hydrogen-like atom with Z protons and only one electron, according to Bohr model, the line velocity of the nth rank electron  $v_{e/z/n}$  satisfies:

$$\frac{v_{e/z/n}}{c} = \frac{Ze^2}{n4\pi\epsilon_0\hbar c} = \frac{Z}{n}\alpha, \text{ as } v_{e/z/n} \leq c, \alpha = \frac{v_{e/z/n}}{c} \frac{n}{Z} \approx \frac{1}{Z_{\max\text{-ideal}}} = \frac{1}{Fy} = \frac{1}{137}$$

The 137th hydrogen-like element Fy (Feynmanium) is an ideal (imaginative) element,

in reality, the above formula should be modified to:  $\alpha = f(Z_{\text{real}}) \frac{1}{Z_{\max\text{-real}}}$

According to Chen's Chirality and Peotry Model of Atomic Nuclei<sup>6</sup>,

$$Z_{\max\text{-real}} = 112 = 2 \cdot 56, \text{ so } \alpha = f(Z_{\text{real}}) \frac{1}{Z_{\max\text{-real}}} = f(Z_{\text{real}}) \frac{1}{112}$$

Compared to  $\alpha = \frac{\lambda_e}{2\pi a_0}$ , the formula should have a  $2\pi$  factor:

$$\alpha = f(Z_{\text{real}}) \frac{1}{Z_{\max\text{-real}}} = \frac{n}{m(2\pi)_k} \frac{1}{Z_{\max\text{-real}}} = \frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112} = 1/136.8$$

Apply with  $2\pi$ -e formula:  $2\pi = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots$

the formula is transformed to:

$$\alpha = \frac{n}{m(2\pi)_k} \frac{1}{Z_{\max\text{-real}}} = \frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112} = 1/137.035782$$

Above deduction on 2018/4/12, only  $(2\pi)_{112}$  gives the closest value to  $\alpha$ , this coincidence of one part per infinity proves the formula itself is correct.

Added an calibration factor ( $\delta=1/75^2$ ) on 2018/4/20, the accurate formula is:

$$\alpha_1 = \frac{\lambda_e}{2\pi a_0} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112 + \frac{1}{75^2}}$$

Discover: 2018/4/12; Revise: 2018/4/20 (add  $1/75^2$  factor)

By the same procedure but compared to  $\alpha = \frac{2\pi r_e}{\lambda_e}$ , the other formula is:

$$\alpha_2 = \frac{2\pi r_e}{\lambda_e} = \frac{m(2\pi)_k}{n} \frac{1}{Z_{\max\text{-real}}} = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{279}{278}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$

Discover: 2018/4/24; Revise: 2018/9/18-20 ( $280 \rightarrow 278$ ,  $-\frac{1}{39^2} + \frac{1}{780^2} \rightarrow -\frac{1}{3 \cdot 29 \cdot 64}$ )

Another amazing coincidence is  $6^2$  and  $10^2$  are square numbers in accordance with  $2\pi = \left(\frac{e}{e^{\gamma_c}}\right)^2$


This also demonstrates that  $\alpha$  has two values with two kinds of formulas.

As  $f(Z_{\text{real}}) = \frac{n}{m(2\pi)_k}$  or  $f(Z_{\text{real}}) = \frac{m(2\pi)_k}{n}$ , m n k  $\delta$  should relate to nucleon numbers of nuclides.

## 7. The Two Most Important Formulas

The above two formulas for  $\alpha_1$  and  $\alpha_2$  were our first gained formulas and are the most important formulas among their serial formulas which will be given followed in this paper. Calculation to give the values of  $\alpha_1$  and  $\alpha_2$  is shown in **Fig. 3** and **Table 2**.

**Fig. 3.** Calculation diagram of  $\alpha_1$  and  $\alpha_2$  (2018/4-6).



**Table 2.** Calculation of  $\alpha_1$  and  $\alpha_2$  (2018/4-6).

k	$(2\pi)_k$	$1/\alpha_1$	k	$(2\pi)_k$	$1/\alpha_2$
	7.389056099	160.917477134		7.389056099	116.596364743
1	6.824768754	148.628533230	1	6.824768754	126.236816375
2	6.640803185	144.622165589	2	6.640803185	129.733867427
3	6.549956514	142.643723845	3	6.549956514	131.533251879
4	6.49586908	141.465817857	4	6.49586908	132.628454999
5	6.46000004	140.684668634	5	6.46000004	133.364872233
6	6.434476503	140.128821836	6	6.434476503	133.893888578
7	6.415388754	139.713132398	7	6.415388754	134.292263980
8	6.400576029	139.390543654	8	6.400576029	134.603053878
9	6.388747203	139.132937708	9	6.388747203	134.852272701
10	6.379083388	138.922480953	10	6.379083388	135.056563407
14	6.353377324	138.362659116	14	6.353377324	135.603008624
28	6.319398093	137.622665802	28	6.319398093	136.332142298
56	6.301583891	137.234711452	56	6.301583891	136.717545138
110	6.29262658	137.039640822	112	6.292459356	136.915795771
111	6.292542221	137.037803660	224	6.28784124	137.016353814
<b>112</b>	<b>6.292459356</b>	<b>137.035999037435</b>	276	6.286966940	137.035408057
113	6.292377945	137.034226098	277	6.286953333	137.035704647
114	6.292297952	137.032484014	<b>278</b>	<b>6.286939823</b>	<b>137.035999111818</b>
			279	6.286926410	137.036291474
			280	6.286913093	137.036581756




In these two formulas (deduced from the modification of  $Z_{\max}$ ), there are some factors which are essentially related to nucleon numbers of some nuclides especially some important stable numbers (stipulated by Chen's Chirality and Poetry Model of Atomic Nuclei<sup>7</sup>) such as 28, 42, 56, 83, 84, 112, 126, 166, 167, 168 *et al.* And these numbers correlate with each others. This kind of relationship is shown in the follows.

A brief illustration of the relationships between the fine-structure constant and nuclides:




Above nuclides indicate that 136–138, which can be called the fine-structure constant numbers, definitely relate to 112 and 166–168 (double of 56 and 83–84, the most stable numbers in nuclides).

$$\alpha_1 = \frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112 + \frac{1}{75^2}} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$




$$\alpha_2 = \frac{13 \cdot (2\pi)_{278}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{9 \cdot 31}{278}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$



The value of the front part of each above formula is almost equal to  $1/(3/2)^{1/2}$  (because 112 is the element natural proton end and 168 is the element natural neutron end as shown in  $^{112}\text{Cn}_{168+5}$ ), so the formulas can be transformed to the follows.


$$\alpha_1 = \alpha_{1-(3/2)} = \frac{1}{\left(\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{2^2 \cdot 3 \cdot 5^3 \cdot 13 \cdot 23 - \frac{30}{64}}\right)^{1/2}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$

2019/4/25 Relations to nuclides :



$$\alpha_2 = \alpha_{2-(3/2)} = \frac{1}{\left(\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{2 \cdot 7 \cdot 11 \cdot 19 \cdot 29 + \frac{36}{75^2}}\right)^{1/2}} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$

2019/4/25 Relations to nuclides:



## 8. The Integrated Fine-structure Constant

Multiplication of  $\alpha_1$  and  $\alpha_2$  should almost divide out the  $2\pi$  factors and give  $3/2$  and  $112 \times 112$  factors, this means  $\alpha_1 \alpha_2$  is almost equal to  $112 \times 168$ , so we define  $\alpha_c = (\alpha_1 \alpha_2)^{1/2}$  as the integrated fine-structure constant or Chen's fine-structure constant.

$$\begin{aligned} \frac{1}{\alpha_c^2} &= \frac{1}{\alpha_1 \alpha_2} = \frac{2\pi a_0}{\lambda_e} \frac{\lambda_e}{2\pi r_e} = \frac{a_0}{r_e} = \left(\frac{c}{v_e}\right)^2 \\ &= 112 \times \left(168 - \frac{1}{3} + \frac{1}{12 \cdot 47} - \frac{1}{6 \cdot 29 \cdot 53 \cdot 59 - 79/47}\right) \quad 2018/6/8-9, 9/18-19, 2019/4/19 \\ &= 136 \left(138 + \frac{1}{2} - \frac{1}{10 \cdot 29} + \frac{1}{12 \cdot 53 \cdot (6 \cdot 53 - 1) - 27/47}\right) \quad 2019/4/17-19 \\ &= 137 \left(137 + \frac{1}{13} - \frac{1}{7 \cdot 29} + \frac{1}{32 \cdot 33 \cdot 89 + 16/49}\right) \quad 2019/4/17-19 \\ &= 112 \cdot 167.668437878408 = 18778.865042381 \\ &\begin{matrix} 27 & 29 & 47,49 & 53 & 54,56,58 & 59 & 58,60,61 & 63,65 & 79 & 87 \\ 13 & Al_{14} & Si_{15} & Ti_{25,27} & Cr_{29} & Fe_{28,30,32} & Co_{32} & Ni_{30,32,33} & Cu_{34,36} & Br_{44} & Sr_{49} \end{matrix} \\ &\begin{matrix} 100,102 & 112 & 113 & 135-138 & 136,138 & 3-47 & 158,160 & 159 & 166,168 \\ 44 & Ru_{56,58} & Cd_{64} & In_{64} & Ba_{79-82} & Ce_{78,80} & Pr_{82} & Gd_{94,96} & Tb_{94} & Er_{98,100} \end{matrix} \\ &\begin{matrix} 174 & 188 & 197 & 203 & 223 & 226 & 227 & 262 & 285 & 293 \\ 70 & Yb_{104} & Os_{112} & Au_{118} & Tl_{122} & Fr_{136}^* & Ra_{138}^* & Ar_{138}^* & Lr_{159}^* & Cn_{173}^* & Lv_{177}^{ie} \end{matrix} \\ \alpha_c^2 &= \alpha_1 \alpha_2 = \left[ \frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112 + \frac{1}{75^2}} \right] \left[ \frac{13 \cdot (2\pi)_{278}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} \right] \\ &= \frac{13 \cdot 3^2}{7 \cdot 5^2} \frac{e^2}{(2 \cdot 3 \cdot 19)^{227}} \frac{e^2}{(115)^{229}} \dots \frac{e^2}{(9 \cdot 31)^{557}} \frac{1}{112^2 - \frac{1}{30^2 \cdot 5} + \frac{1}{60^2 \cdot 15} - \frac{1}{120^2 \cdot 15 \cdot 29}} \\ &= 1/18778.865042381 \quad 2019/12/14 \\ &\begin{matrix} 27 & 31 & 39 & 55 & 54,56,57,58 & 63,65 & 69,71 & 79,81 & 87 \\ 13 & Al_{14} & P_{16} & K_{20} & Mn_{30} & Fe_{28,30,31,32} & Cu_{34,36} & Ga_{38,40} & Br_{44,46} & Sm_{49} \end{matrix} \\ &\begin{matrix} 89 & 93 & 112-120-124 & 135-138 & 139 & 136,138 & 144,145 & 157 \\ 39 & Y_{50} & Nb_{52} & Sn_{62-70-74} & Ba_{79-82} & La_{82} & Ce_{78,80} & Nd_{83,84} & Gd_{93} \end{matrix} \\ &\begin{matrix} 200 & 209 & 223 & 237 & 278+7 & 284 \\ 80 & Hg_{120} & Bi_{126}^* & Fr_{136}^* & Np_{144}^* & Cn_{166+7}^* & Nh_{9,19}^{ie} \end{matrix} \\ \alpha_c^2 &= \alpha_1 \alpha_2 = \frac{1}{\left(\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{7 \cdot 19 \cdot 29 \cdot 37 - \frac{25}{44}}\right)} \frac{1}{112 + \frac{1}{75^2}} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} \\ &= 1/18778.865042381 \quad 2019/12/14 \\ &\begin{matrix} 39 & 47,50 & 55 & 63,65 & 85,87 & 87,88 & 99,100,102,104 & 112 \\ 19 & K_{20} & Ti_{25,28} & Mn_{30} & Cu_{34,36} & Rb_{48,50} & Sr_{49,50} & Ru_{55,56,58,60} & Cd_{64} \end{matrix} \\ &\begin{matrix} 112,114,115,116,120,124 & 5-37,11-17 & 223 & 226 \\ 50 & Sn_{62,64,65,66,70,74} & Re_{110,112} & Fr_{136}^* & Ra_{138}^* \end{matrix} \end{aligned}$$

## 9. A Brief Explanation of the Fine-structure Constant

According to Chen's Chirality and Poetry Model of Atomic Nuclei<sup>7</sup>, the ratio of neutron number  $N$  to proton number  $Z$  in nuclides increases from  $1/1$  to  $3/2$  (eventually slightly above  $3/2$ ) along with the increasing of atomic number, for example, from  ${}_{14}\text{Si}_{14}$ ,  ${}_{26}\text{Fe}_{30}$ ,  ${}_{29}\text{Cu}_{34,36}$ ,  ${}_{56}\text{Ba}_{82}$ ,  ${}_{84}\text{Po}_{125}$  to  ${}_{112}\text{Cn}_{168+5}^*$ . In this process,  $(3/2)^{1/2}$  will act as a transition foothold. As for nuclide  ${}_{112}\text{Cn}_{168+5}$  with  $Z=112$ ,  $N=168+5$  and  $168/112=3/2$ , 137 is just right their  $(3/2)^{1/2}$  times intermediate stage. This should be why 137 exists and what's the real meaning of 137.



electromagnetic wave or light, it should be reasonable to suppose the speed of light to be the integrated fine-structure constant, i.e.,  $c_{au}=1/\alpha_c=1/(\alpha_1\alpha_2)^{1/2}=137.035999074627$ . It means we've theoretically/mathematically calculated the speed of light, the formula is intrinsically consistent with Maxwell's formula, and the value is much accurate.

In atomic units ( $e = m_e = \hbar = 1$  and  $\varepsilon_0 = \frac{1}{4\pi}$ ),  $v_{e/au} = \alpha c_{au} = \frac{e^2}{4\pi\varepsilon_0\hbar} = 1$ , so  $c_{au} = \frac{1}{\alpha}$

There are two  $\alpha$  ( $\alpha_1$  and  $\alpha_2$ ), but there shouldn't be two  $c$  or  $c_{au}$ ,

so it should be:  $c_{au} = \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1\alpha_2}}$  ( $au$ : atomic units)

Compared to Maxwell Formula  $c = \frac{1}{\sqrt{\mu_0\varepsilon_0}}$ ,  $c_{au} = \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1\alpha_2}}$  should be reasonable.

$$c_{au} = \frac{1}{\sqrt{\mu_{0/au}\varepsilon_{0/au}}}, \mu_{0/au}\varepsilon_{0/au} = \alpha_1\alpha_2, \mu_{0/au} = 4\pi\alpha_1\alpha_2 \quad (2019/11/30)$$

So the theoretical formula of the speed of light in atomic units is as follows:

$$\begin{aligned} c_{au} &= \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1\alpha_2}} = \frac{1}{\sqrt{\left(\frac{6^2}{7 \cdot (2\pi)_{112}} \frac{1}{112 + \frac{1}{75^2}}\right) \left(\frac{13 \cdot (2\pi)_{278}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}}\right)}} \\ &= \frac{5}{3} \sqrt{\frac{7 \cdot (2\pi)_{112}}{13 \cdot (2\pi)_{278}} \left(112^2 - \frac{1}{30^2 \cdot 5} + \frac{1}{60^2 \cdot 15} - \frac{1}{120^2 \cdot 15 \cdot 29}\right)} \\ &= \sqrt{\frac{5 \cdot 17 - \frac{10}{11 \cdot 11 \cdot 23} \cdot (2\pi)_{12389}}{7 \cdot \frac{1}{7 \cdot 19} \cdot (2\pi)_{28186}} \left(112^2 - \frac{2 \cdot 7^2 \cdot 43 \cdot 67 + \frac{5}{7}}{5^2 \times 10^{10}}\right)} \\ &= \frac{5}{3} \sqrt{\left(\frac{2^3 \cdot 17}{11 \cdot 23} + \frac{2 \cdot 17}{11 \cdot 23 \cdot 97}\right) \frac{(2\pi)_{34450}}{(2\pi)_{28186}} \left(112^2 - \frac{2^5 \cdot 3 \cdot 7 \cdot 13}{10^{10}}\right)} \\ &= \sqrt{\left(\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{7 \cdot 19 \cdot 29 \cdot 37 - \frac{25}{44}}\right) \left(112^2 - \frac{1}{30^2 \cdot 5} + \frac{1}{60^2 \cdot 15} - \frac{1}{120^2 \cdot 15 \cdot 29}\right)} \\ &= \sqrt{\frac{3}{2} \left(112 - \frac{1}{3^2} + \frac{1}{12^2 \cdot 13 - \frac{30 \cdot 19}{100} - \frac{1}{125 \cdot 100}}\right)} = \sqrt{\frac{3}{2} - \frac{1}{3 \cdot 112 + 1} + \frac{1}{14 \cdot 53 \cdot 193 - \frac{33}{2 \cdot 47}}} \times 112 \\ &= \sqrt{112 \times \left(168 - \frac{1}{3} + \frac{1}{12 \cdot 47} - \frac{1}{6 \cdot 29 \cdot 53 \cdot 59 - 79/47}\right)} \\ &= \sqrt{137.035999037435 \times 137.035999111818} = 137.035999074627 \end{aligned}$$

Note:  $112/278 \approx 27/67$ ,  $12389/28186 \approx 11/25$ ,  $34450/28186 \approx 11/9 \approx 66/29$

Discover: 2019/12/16; Revise and Supplement: 2020/1/5-8, 2/24

## 12. The Special 29 and 75 Factors

In the above formulas some factors especially 29 and 75 appear several times. This feature should be analyzed and explained. Accompanying N/Z ratio from 1/1 to slightly above 3/2 along with the increasing of atomic number,  ${}_{29}\text{Cu}_{34,36}$  is the critical point of N/Z ratio approaching  $(3/2)^{1/2}$  and  ${}_{75}\text{Re}_{110,112}$  is the critical point of N/Z ratio approaching 3/2 (**Table 3**, **Fig. 4** and **Fig. 5**), so 29 and 75 are important factors and hence frequently appear in the formulas.

**Table 3. N/Z ratios of the Elements (2019/4/23).**

Z	N	N/Z	Z	N	N/Z	Z	N	N/Z	Z	N	N/Z				
H	1	0	0	Ga	31	38.80	1.25	Pm	61	84	1.38	Pa*	91	140	1.54
He	2	2.00	1.00	Ge	32	40.71	1.27	Sm	62	88.45	1.43	U*	92	146	1.59
Li	3	3.92	1.31	As	33	42	1.27	Eu	63	89.04	1.41	Np*	93	144	1.55
Be	4	5	1.25	Se	34	45.05	1.33	Gd	64	93.33	1.46	Pu*	94	150	1.60
B	5	5.80	1.16	Br	35	44.98	1.29	Tb	65	94	1.45	Am*	95	148	1.56
<b>C</b>	<b>6</b>	<b>6.01</b>	<b>1.00</b>	Kr	36	47.89	1.33	Dy	66	96.57	1.46	Cm*	96	151	1.57
N	7	7.00	1.00	Rb	37	48.56	1.31	Ho	67	98	1.46	Bk*	97	150	1.55
O	8	8.00	1.00	Sr	38	49.71	1.31	Er	68	99.33	1.46	Cf*	98	153	1.56
F	9	10	1.11	Y	39	50	1.28	Tm	69	100	1.45	Es*	99	153	1.55
Ne	10	10.19	1.02	Zr	40	51.32	1.28	Yb	70	103.11	1.47	Fm*	100	157	1.57
Na	11	12	1.09	Nb	41	52	1.27	Lu	71	104.03	1.47	Md*	101	157	1.55
Mg	12	12.32	1.03	Mo	42	54.04	1.29	Hf	72	106.54	1.48	No*	102	157	1.54
Al	13	14	1.08	Td	43	55	1.28	Ta	73	108	1.48	Lr*	103	159	1.54
Si	14	14.11	1.01	Ru	44	57.16	1.30	W	74	109.89	1.49	Rf*	104	161	1.55
P	15	16	1.07	Rh	45	58	1.29	<b>Re</b>	<b>75</b>	<b>111.25</b>	<b>1.48</b>	Db*	105	163	1.55
S	16	16.09	1.01	Pd	46	60.51	1.32	Os	76	114.27	1.50	Sg*	106	165	1.56
Cl	17	18.48	1.09	Ag	47	60.96	1.30	Ir	77	115.25	1.50	Bh*	107	163	1.52
Ar	18	21.99	1.22	Cd	48	64.52	1.34	Pt	78	117.12	1.50	Hs*	108	169	1.56
K	19	20.13	1.06	In	49	65.91	1.35	Au	79	118	1.49	Mt*	109	167	1.53
Ca	20	20.12	1.01	Sn	50	68.81	1.38	Hg	80	120.62	1.51	Ds*	110	171	1.55
Sc	21	24	1.14	Sb	51	70.86	1.39	Tl	81	123.41	1.52	Rg*	111	169	1.52
Ti	22	25.92	1.18	Te	52	75.70	1.46	Pb	82	125.24	1.53	<b>Cn*</b>	<b>112</b>	<b>173</b>	<b>1.54</b>
V	23	28	1.22	I	53	74	1.40	Bi*	83	126	1.52	Nh*	113	171	1.51
Cr	24	28.06	1.17	Xe	54	77.39	1.43	Po*	84	125	1.49	Fl*	114	175	1.54
Mn	25	30	1.20	Cs	55	78	1.42	At*	85	125	1.47	Mc*	115	173	1.50
Fe	26	29.91	1.15	Ba	56	81.42	1.45	Rn*	86	136	1.58	Lv*	116	177	1.53
Co	27	32	1.19	La	57	82	1.44	Fr*	87	136	1.56	Ts*	117	177	1.51
Ni	28	30.76	1.10	Ce	58	82.21	1.42	Ra*	88	138	1.57	Og*	118	176	1.49
<b>Cu</b>	<b>29</b>	<b>34.62</b>	<b>1.19</b>	Pr	59	82	1.39	Ac*	89	138	1.55				
Zn	30	35.45	1.18	Nd	60	84.41	1.41	Th*	90	142	1.58				


Z: atomic number, N: average neutron number or neutron number of the most stable isotope.

1. N/Z from 1/1 ( ${}_6\text{C}$ ) to slightly above 3/2 (such as  ${}_{112}\text{Cn}$  which is the natural end of elements demonstrated by Chen's Chirality and Poetry Model of Atomic Nuclei<sup>7</sup>).
2. For  ${}_{29}\text{Cu}$ , N/Z ratio 1.19 is near to  $(3/2)^{1/2}=1.22$ , slightly less is because of stability effect.
3. For  ${}_{75}\text{Re}$ , N/Z ratio 1.48 is near to  $3/2=1.50$ , slightly less is because of stability effect.
4. From  ${}_6\text{C}$  to  ${}_{112}\text{Cn}$ , the middle of N/Z 1.5 range is at  $(76.5-5)/(112-5)=0.668 \approx 2/3$  position.


**Fig. 4** and **Fig. 5** shows that stability effect of nucleon number 64 makes the neutron numbers of  ${}_{29}\text{Cu}$ 's isotopes are relatively less (34 and 36) than normal so that its N/Z ratio is a little less than  $(3/2)^{1/2}$  which is otherwise it should be. Also the

stability effect of nucleon numbers 110 and 112 make the neutron numbers of  $^{75}\text{Re}$ 's nuclides are relatively less (110 and 112) than normal so that its N/Z ratio is a little less than  $3/2$  which otherwise it should be.

**Fig. 4. Complete Graph of N/Z Ratios of Elements (2019/4/23-24).**



**Fig. 5. Partially Amplified Graph of N/Z Ratios of Elements (2019/4/24).**




The general trend of N/Z ratio of elements is from 1/1 ( ${}^6\text{C}_6$ ) to slightly above 3/2 ( ${}^{112}\text{Cn}_{173}$ ) definitely. However, the increasing process is not smooth, the N/Z ratio rising fluctuates consecutively. According to Chen's Chirality and Poetry Model of Atomic Nuclei<sup>7</sup>, there are some stable numbers (magic numbers) which can bring about this kind of fluctuation (**Table 4** and **Fig. 6**).

**Table 4. Effect of Stable Numbers on N/Z ratio's fluctuation (2019/4/22).**

Element	Z	N(Average)	Z(3/2) <sup>1/2</sup>	N-Z(3/2) <sup>1/2</sup>	Stable Number
K	19	20.13	23.27	-3.17	20
Ca	20	20.12	24.49	-4.41	20+20
Sc	21	24	25.72	-1.74	
Ti	22	25.92	26.94	-1.07	22+26=48
V	23	28.00	28.17	-0.23	28
Cr	24	28.06	29.39	-1.39	28
Mn	25	30	30.62	-0.68	
Fe	26	29.91	31.84	-1.99	26+30=56
Co	27	32.00	33.07	-1.14	
Ni	28	30.76	34.29	-3.60	28+30=58、28+32=60
<b>Cu</b>	<b>29</b>	<b>34.62</b>	<b>35.52</b>	<b>-0.97</b>	<b>64</b>
Zn	30	35.45	36.74	-1.36	30+34=64、30+36=66
Ga	31	38.80	37.97	0.75	
Ge	32	40.71	39.19	1.44	32+40=72
As	33	42.00	40.42	1.50	
Se	34	45.05	41.64	3.30	34+46=80
Br	35	44.98	42.87	2.03	
Kr	36	47.89	44.09	3.71	36+48=84

**Fig. 6. Effect of Stable Numbers on N/Z ratio's fluctuation (2019/4/22-23)**



### 13. $\alpha_1/\alpha_2$ in Schrödinger Equation of Hydrogen Atom

Stationary Schrodinger Equation  $-\frac{\hbar^2}{2m}\nabla^2\psi + U\psi = E\psi$ , applied to hydron atom:

$$\nabla^2\psi + \frac{2m_e}{\hbar^2}(E + \frac{e^2}{4\pi\epsilon_0 r})\psi = 0, \quad E = -\frac{m_e e^4}{2n^2(4\pi\epsilon_0)^2\hbar^2}, \text{ do substitution and simplification:}$$

$$\frac{2m_e}{\hbar^2}(\frac{m_e e^4}{2n^2(4\pi\epsilon_0)^2\hbar^2} - \frac{e^2}{4\pi\epsilon_0 r})\psi = \nabla^2\psi, \quad [\frac{1}{n^2}(\frac{m_e e^2}{4\pi\epsilon_0\hbar^2})^2 - \frac{2}{r} \frac{m_e e^2}{4\pi\epsilon_0\hbar^2}]\psi = \nabla^2\psi,$$

$$[\frac{1}{n^2}(\frac{e^2}{4\pi\epsilon_0\hbar c} \frac{m_e c}{\hbar})^2 - \frac{2}{r} \frac{e^2}{4\pi\epsilon_0\hbar c} \frac{m_e c}{\hbar}]\psi = \nabla^2\psi,$$

$$\text{As } \sqrt{\alpha_1\alpha_2} = \frac{v_e}{c} = \frac{e^2}{4\pi\epsilon_0\hbar c}, \lambda_e = \frac{h}{m_e c} \text{ and } \alpha_1 = \frac{\lambda_e}{2\pi a_0}:$$

$$[\frac{1}{n^2}(\sqrt{\alpha_1\alpha_2} \frac{2\pi}{\lambda_e})^2 - \frac{2}{r} \sqrt{\alpha_1\alpha_2} \frac{2\pi}{\lambda_e}]\psi = \nabla^2\psi,$$

$$[\frac{1}{n^2(\lambda_e/2\pi/\sqrt{\alpha_1\alpha_2})^2} - \frac{2}{(\lambda_e/2\pi/\sqrt{\alpha_1\alpha_2})r}]\psi = \nabla^2\psi,$$

$$[\frac{1}{n^2 a_0^2 (\alpha_1/\alpha_2)} - \frac{2}{a_0 r \sqrt{\alpha_1/\alpha_2}}]\psi = \nabla^2\psi$$

$$\text{As } \alpha_1/\alpha_2 \approx 1, \text{ simplified to: } [\frac{1}{n^2 a_0^2} - \frac{2}{a_0 r}]\psi = \nabla^2\psi \text{ (factor 2 seems not beautiful)}$$

In atomic units (*au*:  $e = m_e = \hbar = 1$  and  $\epsilon_0 = \frac{1}{4\pi}$ ),

$$a_{0/au} = \frac{4\pi\epsilon_0\hbar^2}{m_e e^2} = 1, \quad v_{e/au} = \frac{e^2}{4\pi\epsilon_0\hbar} = 1, \quad c_{au} = \frac{v_{e/au}}{\alpha_c} = \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1\alpha_2}}$$

$$[\frac{1}{n^2(\alpha_1/\alpha_2)} - \frac{2}{r_{au}\sqrt{\alpha_1/\alpha_2}}]\psi = \nabla_{au}^2\psi, \text{ or } (\frac{c_{au}^2}{\alpha_1^2 n^2} - \frac{2c_{au}}{\alpha_1 r_{au}})\psi = \nabla_{au}^2\psi$$

the above equation could be called Schrodinger-Chen equation of hydrogen atom, the later form of the equation shows factor 2 is still reasonable and beautiful.

$$\text{As } \alpha_1/\alpha_2 \approx 1, \text{ simplified to: } [\frac{1}{n^2} - \frac{2}{r_{au}}]\psi = \nabla_{au}^2\psi$$

Discover: 2018/4-6; Revise: 2019/12/13 (add *au* form)

$$\alpha_1/\alpha_2 = \frac{137.035999111818}{137.035999037435} = 1.0000000005428 = 1 + \frac{23 \cdot 59}{25 \cdot 10^{11}} = (1 + \frac{23 \cdot 59}{50 \cdot 10^{11}})^2$$

$$\sqrt{\alpha_1/\alpha_2} = 1 + \frac{23 \cdot 59}{50 \cdot 10^{11}} = 1.0000000002714$$

Relations to nuclides:  ${}_{11}^{23}\text{Na}_{12}$   ${}_{23}^{50,51}\text{V}_{27,28}$   ${}_{25}^{55}\text{Mn}_{30}$   ${}_{44}^{99,100}\text{Ru}_{55,56}$   ${}_{46}^{105}\text{Pd}_{59}$   ${}_{56}^{137}\text{Ba}_{81}$   
 ${}_{50}^{118+1}\text{Sn}_{69}$   ${}_{59}^{141}\text{Pr}_{82}$   ${}_{69}^{169}\text{Tm}_{100}$   ${}_{75}^{185,187}\text{Re}_{110,112}$   ${}_{88}^{169}\text{Ra}_{137}^*$

2019/8/28-29

Solution of Schrödinger equation of hydrogen atom gives some quantum numbers such as  $n$ ,  $l$  and  $m_l$  which determine the electron shell structure and the chemical



properties of atoms. That means Schrödinger equation of hydrogen atom is the base of chemical periodicity of elements. On the other hand, from above analysis, we have already demonstrated the formulas of the fine-structure constant  $\alpha$  are derived from Chen's Chirality and Poetry Model of Atomic Nuclei<sup>7</sup> and hence mainly connected to the stability of atomic nuclei. So, a question is whether and how  $\alpha$  is connected to Schrödinger Equation of hydrogen atom. This question should reveal the connection of the theory of electron shell of atoms and the theory of nuclei of elements. The above deduction provides the answer. The fine-structure constant  $\alpha$  relates to Schrödinger Equation of hydrogen atom in  $\alpha_1/\alpha_2$  way which is subtle and negligible but could show the equation is really reasonable and beautiful.

#### 14. The Two Kinds of General Formulas of the Fine-structure Constant

Based on the above two formulas of  $\alpha_1$  and  $\alpha_2$ , it should be reasonable to assume there are two kinds of serial formulas of  $\alpha_1$  and  $\alpha_2$  which are listed in follows. Among these formulas, the above two first discovered formulas are the most fundamental and important. Some formulas both with a big  $m$  and an extra large  $k$  should be more important referring to the trend of the approximate values of  $\alpha$ .

Approximate formulas:

$$\alpha_{1-m'} = \frac{n}{m \cdot (2\pi)_k} \frac{1}{112} = \frac{n}{m \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}} \frac{1}{112} \approx 1/137.036$$

$$\alpha_{2-m'} = \frac{m \cdot (2\pi)_k}{n} \frac{1}{112} = \frac{m \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}}{n} \frac{1}{112} \approx 1/137.036$$

Accurate Formulas:

$$\alpha_{1-m} = \frac{n}{m \cdot (2\pi)_k} \frac{1}{112 + \delta_1} = \frac{n}{m \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}} \frac{1}{112}$$

$$= 1/137.035999037435$$

$$\alpha_{2-m} = \frac{m \cdot (2\pi)_k}{n} \frac{1}{112 - \delta_2} = \frac{m \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{k+1}{k}\right)^{2k+1}}}{n} \frac{1}{112 - \delta_2}$$


$$= 1/137.035999111818$$

Discover: 2019/6/27; Revise: 2019/7/2-3


**Table 5. Parameters and Results of Approximate Formulas of  $\alpha_1$  (2019/7/2).**

<b>m</b>	<b>n</b>	<b>k</b>	<b><math>\alpha_{1-m'}</math></b>	<b>m</b>	<b>n</b>	<b>k</b>	<b><math>\alpha_{1-m'}</math></b>
1	6	1	122.265854937	24	124	27	137.016359405
2	11	2	135.230901223	<b>25</b>	129	34	137.030171763
3	<b>16</b>	4	136.413250690	26	134	46	137.027100696
<b>4</b>	21	7	136.861626741	27	139	66	137.033636049
5	26	13	136.950569252	28	144	112	137.035781520
6	31	27	137.016359405	29	149	<b>321</b>	137.035917078
<b>7</b>	<b>36</b>	<b>112</b>	<b>137.035781520</b>	30	155	27	137.016359405
8	42	7	136.861626741	31	<b>160</b>	32	137.035453560
<b>9</b>	47	9	137.018237882	<b>32</b>	165	40	137.034309209
10	52	13	136.950569252	33	170	52	137.034083409
11	57	18	137.001388822	34	<b>175</b>	72	137.034617877
12	62	27	137.016359405	35	180	112	137.035781520
13	67	46	137.027100696	<b>36</b>	185	<b>236</b>	137.035810961
14	72	112	137.035781520	43	221	<b>200</b>	137.035845637
15	78	13	136.950569252	50	257	<b>181</b>	137.035307038
<b>16</b>	83	16	136.992590996	59	303	<b>2645</b>	137.035986189
17	<b>88</b>	20	137.028423583	<b>81</b>	<b>416</b>	<b>1605</b>	137.035992406
18	93	27	137.016359405	<b>96</b>	493	<b>5806</b>	137.035998789
19	<b>98</b>	37	137.034579883	103	<b>529</b>	<b>1310</b>	137.035994308
<b>20</b>	103	58	137.030572071	<b>133</b>	<b>683</b>	<b>12389</b>	<b>137.035999034</b>
21	108	112	137.035781520	140	719	<b>1923</b>	137.035994882
<b>22</b>	<b>113</b>	<b>782</b>	<b>137.035967638</b>	155	<b>796</b>	<b>3988</b>	137.035997989
23	119	22	137.001596764	<b>170</b>	<b>873</b>	<b>34450</b>	<b>137.035999031</b>


**Fig. 7. Results of Approximate Formulas of  $\alpha_1$  (2019/7/2).**




$$\alpha_{1-1} = \frac{6}{1 \cdot e^2 \left(\frac{2}{1}\right)^2} \frac{1}{112 + \frac{17}{2} - \frac{1}{40} + \frac{1}{6 \cdot 23 \cdot 25 - \frac{36}{55}}} = 1/137.035999037434$$




$$\alpha_{1-2} = \frac{11}{2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5}} \frac{1}{112 + \frac{3}{2} - \frac{1}{200} + \frac{1}{5 \cdot (3 \cdot 42 + 1) \cdot (6 \cdot 37 - 1) + \frac{2}{7}}} = 1/137.035999037435$$




$$\alpha_{1-3} = \frac{4^2}{3 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \frac{e^2}{\left(\frac{5}{4}\right)^9}} \frac{1}{112 + 1 - \frac{1}{2} + \frac{1}{88} - \frac{1}{13 \cdot (2 \cdot 9 \cdot 5 \cdot 13 + 1) - \frac{2}{73}}} = 1/137.035999037435$$




$$\alpha_{1-4} = \frac{21}{2^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{8}{7}\right)^{15}}} \frac{1}{112 + \frac{1}{7} - \frac{1}{8 \cdot 19 \cdot 41 - \frac{75}{98}}} = 1/137.035999037435$$




$$\alpha_{1-5} = \frac{26}{5 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14}{13}\right)^{27}}} \frac{1}{112 + \frac{1}{14} - \frac{1}{9 \cdot 71} + \frac{1}{67 \cdot (75 \cdot 100 - 1) + \frac{1}{10}}} = 1/137.035999037435$$




$$\alpha_{1-6} = \frac{31}{6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{28}{27}\right)^{55}}} \frac{1}{112 + \frac{1}{2 \cdot 31} - \frac{1}{3 \cdot 11 \cdot 13 \cdot 31 - \frac{43}{4 \cdot 27}}} = 1/137.035999037435$$




$$\alpha_1 = \alpha_{1-7} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$




$$\alpha_{1-9} = \frac{47}{3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{10}{9}\right)^{19}} 112 + \frac{1}{4 \cdot 17} - \frac{1}{2(8 \cdot 9 \cdot 37 - 1) + \frac{3 \cdot 17}{157}}} = 1/137.035999037436$$




$$\alpha_{1-11} = \frac{3 \cdot 19}{11 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{19}{18}\right)^{37}} 112 + \frac{1}{35} - \frac{1}{88 \cdot 41 - \frac{5 \cdot 53}{22 \cdot 13}}} = 1/137.035999037435$$




$$\alpha_{1-13} = \frac{67}{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{47}{2 \cdot 23}\right)^{3 \cdot 31}} 112 + \frac{1}{137} - \frac{1}{6(2 \cdot 27 \cdot 59 + 1) + \frac{9}{50}}} = 1/137.035999037435$$




$$\alpha_{1-16} = \frac{83}{4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{17}{16}\right)^{33}} 112 + \frac{1}{28} - \frac{1}{6 \cdot (18 \cdot 41 + 1) + \frac{173}{2 \cdot (2 \cdot 75 - 1)}}} = 1/137.035999037435$$




$$\alpha_{1-17} = \frac{2^2 \cdot 22}{17 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{21}{20}\right)^{41}} 112 + \frac{1}{137} - \frac{1}{2 \cdot 19 \cdot 23 \cdot 59 - \frac{30}{100}}} = 1/137.035999037435$$



$$\alpha_{1-19} = \frac{2 \cdot 7^2}{19 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{38}{37}\right)^{75}} 112 + \frac{1}{2 \cdot (8 \cdot 54 - 1) + \frac{54}{19^2}}} = 1/137.035999037440$$




$$\alpha_{1-20} = \frac{103}{2^2 \cdot 5 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{59}{2 \cdot 29}\right)^{9 \cdot 13}} 112 + \frac{1}{32 \cdot 45 \cdot 79 + \frac{22}{3 \cdot 17}}} = 1/137.035999037435$$




$$\alpha_{1-22} = \frac{113}{22 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{27 \cdot 29}{2 \cdot 17 \cdot 23}\right)^{5 \cdot (2 \cdot 157 - 1)}}} \frac{1}{112 + \frac{1}{2 \cdot [2 \cdot 3 \cdot 17 \cdot (10 \cdot 19 + 1) + 1] + \frac{29}{49}}}$$


$$= 1/137.035999037435$$




$$\alpha_{1-23} = \frac{7 \cdot 17}{23 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{23}{22}\right)^{45}}} \frac{1}{112 + \frac{1}{35} - \frac{1}{4 \cdot 13 \cdot 43 - \frac{2 \cdot 29}{16 \cdot 17 - 1}}} = 1/137.035999037435$$




$$\alpha_{1-25} = \frac{3 \cdot 43}{5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{35}{34}\right)^{3 \cdot 23}}} \frac{1}{11 \cdot 19 - \frac{1}{13^2(16 \cdot 17 - 1) + \frac{11}{25}}} = 1/137.035999037435$$




$$\alpha_{1-27} = \frac{139}{27 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{67}{66}\right)^{7 \cdot 19}}} \frac{1}{11 \cdot 47 + \frac{18}{23} + \frac{1}{6 \cdot 23 \cdot 137}} = 1/137.035999037435$$




$$\alpha_{1-29} = \frac{149}{29 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14 \cdot 23}{3 \cdot 107}\right)^{643}}} \frac{1}{6 \cdot 8 \cdot (12 \cdot 26 - 1) + \frac{11}{18}} = 1/137.035999037434$$




$$\alpha_{1-31} = \frac{4^2 \cdot 10}{31 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{33}{32}\right)^{5 \cdot 13}}} \frac{1}{12 \cdot 11 \cdot 17 - \frac{4 \cdot 49}{5 \cdot 41}} = 1/137.035999037434$$




$$\alpha_{1-32} = \frac{15 \cdot 11}{2 \cdot 4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{41}{40}\right)^{81}}} \frac{1}{25 \cdot 29 - \frac{5 \cdot 83}{19 \cdot 23}} = 1/137.035999037435$$




$$\alpha_{1-33} = \frac{170}{33 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{53}{4 \cdot 13}\right)^{105}} 112 + \frac{1}{22 \cdot 29 + \frac{4 \cdot 73}{5 \cdot 83}}} = 1/137.035999037436$$




$$\alpha_{1-34} = \frac{7 \cdot 5^2}{34 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{73}{72}\right)^{5 \cdot 29}} 112 + \frac{1}{15 \cdot 59 + \frac{13}{15} + \frac{1}{3 \cdot (2 \cdot 15 \cdot 17 - 1)}}} = 1/137.035999037435$$




$$\alpha_{1-36} = \frac{5 \cdot 37}{6^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 79}{4 \cdot 59}\right)^{11 \cdot 43}} 112 + \frac{1}{5 \cdot (31 \cdot 42 - 1) + \frac{3 \cdot 31}{14 \cdot 13}}} = 1/137.035999037436$$




$$\alpha_{1-43} = \frac{13 \cdot 17}{43 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 67}{200}\right)^{401}} 112 + \frac{1}{8 \cdot (12 \cdot 83 + 1) + \frac{4}{3 \cdot 13}}} = 1/137.035999037436$$




$$\alpha_{1-50} = \frac{2 \cdot 257}{100 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14 \cdot 13}{181}\right)^{311^2}} 112 + \frac{1}{29 \cdot 61 + \frac{157}{16 \cdot 11}}} = 1/137.035999037436$$



$$\alpha_{1-59} = \frac{3 \cdot 101}{59 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 27 \cdot 49}{5 \cdot 23^2}\right)^{11 \cdot 13 \cdot 37}} 112 + \frac{1}{48 \cdot 64 \cdot 31 - \frac{17}{81}}} = 1/137.035999037435$$




$$\alpha_{1-81} = \frac{4^2 \cdot 26}{9^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{22 \cdot 73}{15 \cdot 107}\right)^{13^2 \cdot 19}} 112 + \frac{1}{2 \cdot 81 \cdot 17 \cdot 67 + \frac{35}{88}}} = 1/137.035999037435$$




$$\alpha_{1-96} = \frac{17 \cdot 29}{4^2 \cdot 6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{16 \cdot 3 \cdot 11^2 - 1}{27 \cdot 5 \cdot 43 + 1}\right)^{79 \cdot 147}}} \frac{1}{112 + \frac{163 \cdot (8 \cdot 21 \cdot 37 + 1)}{50 \cdot 10^{11}}}$$

$$= 1/137.035999037435$$




$$\alpha_{1-103} = \frac{23^2}{103 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 19 \cdot 23}{7 \cdot 11 \cdot 17 + 1}\right)^{2621}}} \frac{1}{112 + \frac{1}{6 \cdot (12 \cdot (8 \cdot (64 \cdot 7 + 1) + 1) + 1) + \frac{3}{4}}}$$

$$= 1/137.035999037435$$




$$\alpha_{1-133} = \frac{683}{133 \cdot (2\pi)_{12389}} \frac{1}{112 + \frac{14651}{50 \cdot 10^{11}}} = \frac{6^2 \cdot 19 - 1}{7 \cdot 19 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{59 \cdot 210}{13 \cdot (17 \cdot 56 + 1)}\right)^{71 \cdot (12 \cdot 29 + 1)}}} \frac{1}{112 + \frac{7^2 \cdot 13 \cdot 23}{50 \cdot 10^{11}}}$$

$$= 1/137.035999037435$$




$$\alpha_{1-140} = \frac{6^2 \cdot 20 - 1}{140 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{4 \cdot 13 \cdot 37}{3 \cdot (64 \cdot 10 + 1)}\right)^{3847}}} \frac{1}{112 + \frac{1}{4 \cdot 9 \cdot (2 \cdot 3 \cdot 29 \cdot 47 + 1) + \frac{29}{54}}}$$

$$= 1/137.035999037435$$




$$\alpha_{1-155} = \frac{2^2 \cdot 199}{5 \cdot 31 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left[\frac{19 \cdot 210 - 1}{3^2 \cdot (2 \cdot 13 \cdot 17 + 1) + 1}\right]^{7977}}} \frac{1}{112 + \frac{1}{5 \cdot 17 \cdot 31 \cdot (2 \cdot 13 \cdot 17 + 1) - \frac{15}{43}}}$$

$$= 1/137.035999037435$$



$$\alpha_{1-170} = \frac{873}{170 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{34451}{34450}\right)^{68901}}} \frac{1}{112 + \frac{4171}{8 \times 10^{11}}}$$


$$= \frac{3^2 \cdot 97}{170 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left[\frac{47(12 \cdot 61 + 1)}{2 \cdot 25 \cdot 13 \cdot 53}\right]^{3 \cdot 7 \cdot 17 \cdot 193}}} \frac{1}{112 + \frac{43 \cdot 97}{8 \cdot 10^{11}}} = 1/137.035999037435$$



**Table 6. Parameters and Results of Approximate Formulas of  $\alpha_2$  (2019/7/3).**


<b>m</b>	<b>n</b>	<b>k</b>	<b><math>\alpha_{2-m'}</math></b>	<b>m</b>	<b>n</b>	<b>k</b>	<b><math>\alpha_{2-m'}</math></b>
1	<b>8</b>	4	137.933814383	22	170	32	137.047480404
2	16	4	137.933814383	23	177	<b>161</b>	137.036664793
3	24	4	137.933814383	<b>24</b>	185	62	137.040748949
<b>4</b>	31	20	137.054511358	<b>25</b>	193	39	137.039552569
5	39	11	137.120466691	26	200	278	137.036218856
6	47	8	137.070996332	27	<b>208</b>	80	137.038980680
7	<b>54</b>	48	137.042951195	28	216	48	137.042951195
8	62	20	137.054511358	<b>29</b>	<b>223</b>	<b>655</b>	<b>137.036002235</b>
<b>9</b>	70	14	137.109928583	30	231	104	137.037530964
10	77	<b>104</b>	137.037530964	31	239	58	137.040063944
11	85	32	137.047480404	<b>32</b>	247	41	137.044550585
12	93	20	137.054511358	33	254	<b>138</b>	137.036795730
<b>13</b>	<b>100</b>	<b>278</b>	<b>137.036218856</b>	34	262	70	137.038016730
14	108	48	137.042951195	35	270	48	137.042951195
15	<b>116</b>	28	137.059466839	<b>36</b>	277	<b>190</b>	137.036562950
16	124	20	137.054511358	37	285	85	137.037566566
17	131	70	137.038016730	38	293	56	137.041569603
18	139	37	137.048943854	39	300	278	137.036218856
19	147	26	137.063298933	<b>125</b>	<b>961</b>	<b>4293</b>	137.035999678
20	154	104	137.037530964	<b>253</b>	<b>1945</b>	<b>28186</b>	<b>137.035999128</b>
21	162	48	137.042951195	269	2068	<b>41654</b>	137.035999118

**Fig. 8. Results of Approximate Formulas of  $\alpha_2$  (2019/7/3).**







$$\alpha_{2-1} = \frac{e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \frac{e^2}{\left(\frac{5}{4}\right)^9}}{2 \cdot 2^2} \frac{1}{112 - 1 + \frac{1}{3} - \frac{1}{16} + \frac{1}{41 \cdot (12 \cdot 13 + 1) + \frac{13}{41}}} = 1/137.035999111816$$




$$\alpha_{2-4} = \frac{2^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{21}{20}\right)^{41}}}{31} \frac{1}{112 - \frac{1}{66} + \frac{1}{71 \cdot (14 \cdot 43 - 1) - \frac{56}{95}}} = 1/137.035999111818$$




$$\alpha_{2-5} = \frac{5 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{12}{11}\right)^{23}}}{39} \frac{1}{112 - \frac{1}{14} + \frac{1}{10 \cdot 41} - \frac{1}{23 \cdot (14 \cdot 11 \cdot 79 + 1) + \frac{11}{16}}} = 1/137.035999111818$$




$$\alpha_{2-6} = \frac{6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{9}{8}\right)^{17}}}{47} \frac{1}{112 - \frac{1}{2 \cdot 17} + \frac{1}{2 \cdot (36 \cdot 17 + 1) - \frac{4}{47}}} = 1/137.035999111818$$




$$\alpha_{2-7} = \frac{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{48}{47}\right)^{95}}}{6 \cdot 3^2} \frac{1}{112 + \frac{1}{2 \cdot 13 \cdot 17} - \frac{1}{2 \cdot 29 \cdot (24 \cdot 89 + 1) + \frac{11}{2 \cdot 17}}} = 1/137.035999111818$$




$$\alpha_{2-9} = \frac{3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{15}{14}\right)^{29}}}{70} \frac{1}{112 - \frac{1}{16} + \frac{1}{11 \cdot 43} - \frac{1}{70 \cdot 17 \cdot (3 \cdot 64 - 1) - \frac{41}{70}}} = 1/137.035999111818$$




$$\alpha_{2-10} = \frac{10 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{5 \cdot 21}{8 \cdot 13}\right)^{11 \cdot 19}}}{77} \frac{1}{112 - \frac{1}{3 \cdot 14 \cdot 19} + \frac{1}{14 \cdot (4 \cdot 27 \cdot (2 \cdot 15 \cdot 19 + 1) - 1)}} = 1/137.035999111818$$




$$\alpha_{2-11} = \frac{11 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{33}{32}\right)^{65}}}{85} \frac{1}{112 - \frac{1}{106} + \frac{1}{30 \cdot (4 \cdot 163 + 1) - \frac{35}{52}}} = 1/137.035999111818$$




$$\alpha_2 = \alpha_{2-13} = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{9 \cdot 31}{2 \cdot 139}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.035999111818$$




$$\alpha_{2-15} = \frac{15 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{29}{28}\right)^{57}}}{2^2 \cdot 29} \frac{1}{112 - \frac{1}{4 \cdot 13} + \frac{1}{12 \cdot (36 \cdot 43 + 1) - \frac{1}{16}}} = 1/137.035999111818$$



$$\alpha_{2-17} = \frac{17 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{71}{70}\right)^{3 \cdot 47}}}{131} \frac{1}{112 - \frac{1}{6 \cdot 101} + \frac{1}{23 \cdot (30 \cdot 35^2 - 1) + \frac{6}{23}}} = 1/137.035999111818$$



$$\alpha_{2-18} = \frac{18 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{38}{37}\right)^{75}}}{139} \frac{1}{112 - \frac{1}{2 \cdot 47} + \frac{1}{2 \cdot 31 \cdot (16 \cdot 17 - 1) + \frac{83}{137}}} = 1/137.035999111818$$



$$\alpha_{2-19} = \frac{19 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{27}{26}\right)^{53}}}{3 \cdot 49} \frac{1}{112 - \frac{1}{44} + \frac{1}{16 \cdot (4 \cdot 37 + 1) - \frac{23}{6 \cdot 47 + 1}}} = 1/137.035999111818$$

$$\alpha_{2-23} = \frac{23 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 81}{7 \cdot 23}\right)^{17 \cdot 19}}}{3 \cdot 59} \frac{1}{112 - \frac{1}{2 \cdot (40 \cdot 23 - 1) + \frac{9}{32 \cdot 10}}} = 1/137.035999111818$$

$$\alpha_{2-24} = \frac{2^2 \cdot 6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{63}{2 \cdot 31}\right)^{125}}}{5 \cdot 37} \frac{1}{112 - \frac{1}{257} + \frac{1}{10 \cdot (12 \cdot 13 \cdot 83 + 1) + \frac{23}{81}}} = 1/137.035999111818$$

$$\alpha_{2-25} = \frac{5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{40}{3 \cdot 13}\right)^{79}}}{193} \frac{1}{112 - \frac{1}{8 \cdot 43} + \frac{1}{18 \cdot 23 \cdot (32 \cdot 27 - 1) - \frac{3}{7}}} = 1/137.035999111818$$

$$\alpha_{2-27} = \frac{3 \cdot 3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{81}{80}\right)^{7 \cdot 23}}}{4^2 \cdot 13} \frac{1}{112 - \frac{1}{10 \cdot 41} + \frac{1}{2 \cdot 27 \cdot 43 \cdot (3 \cdot 64 + 1) - \frac{19}{26}}} = 1/137.035999111818$$

$$\alpha_{2-29} = \frac{29 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{16 \cdot 41}{5 \cdot 131}\right)^{3 \cdot 19 \cdot 23}}}{223} \frac{1}{112 - \frac{1}{29 \cdot 59 \cdot (12 \cdot 19 + 1) + \frac{19}{29}}} = 1/137.035999111818$$

$$\alpha_{2-31} = \frac{31 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{16 \cdot 31}{5 \cdot 131}\right)^{3 \cdot 19 \cdot 23}}}{223} \frac{1}{112 - \frac{1}{29 \cdot 59 \cdot (12 \cdot 19 + 1) + \frac{19}{29}}} = 1/137.035999111818$$

$$\alpha_{2-31} = \frac{31 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{59}{58}\right)^{9 \cdot 13}}}{7 \cdot 34 + 1} \frac{1}{112 - \frac{1}{7 \cdot 43 + \frac{9}{5 \cdot 113}}} = 1/137.035999111819$$

<sup>63,65</sup>Cu<sub>34,36</sub> <sup>69,71</sup>Ga<sub>38,40</sub> <sup>76,77,78</sup>Se<sub>42,43,44</sub> <sup>89</sup>Y<sub>50</sub> <sup>99</sup>Tc\*<sub>43,56</sub> <sup>136,138,140,141</sup>Ce<sub>58,78,80,82,84</sub> <sup>146,147,148,152</sup>Sm<sub>62,84,85,86,90</sub>  
<sup>222</sup>Rn\*<sub>136</sub> <sup>284</sup>Nh<sup>ie</sup><sub>113</sub> <sup>310,24,13</sup>Ch<sup>ie</sup><sub>124,125</sub> <sup>186,187</sup>Ch<sup>ie</sup><sub>148</sub> <sup>22,17</sup>Ch<sup>ie</sup><sub>155=5,31,156=12,13</sub> <sup>8,49,394</sup>Ch<sup>ie</sup><sub>237,7,34</sub>

$$\alpha_{2-32} = \frac{2 \cdot 4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{42}{41}\right)^{83}}}{13 \cdot 19} \frac{1}{112 - \frac{1}{11 \cdot 13} + \frac{1}{6 \cdot 37 \cdot (5 \cdot 210 - 1) + \frac{10}{11}}} = 1/137.035999111818$$

= 1/137.035999111818

<sup>39,41</sup>K<sub>19,20,22</sub> <sup>70,72,73,74,76</sup>Ge<sub>32,38,40,41,42,44</sub> <sup>85,87</sup>Rb<sub>37,48,50</sub> <sup>209</sup>Bi\*<sub>83,126</sub> <sup>210</sup>At\*<sub>85,125</sub> <sup>235</sup>U\*<sub>92,113</sub> <sup>18,17,28,11</sup>Ch<sup>ie</sup><sub>123=3,41</sub>  
<sup>3,11,2,28,13</sup>Ch<sup>ie</sup><sub>143=11,13</sub> <sup>410</sup>Ch<sup>ie</sup><sub>220,13,17</sub> <sup>247=13,19</sup>Ch<sup>ie</sup><sub>163</sub> <sup>32,13</sup>Ch<sup>ie</sup><sub>164=4,41</sub> ? <sup>420</sup>Ch<sup>ie</sup><sub>165=11,15</sub> <sup>418,420</sup>Ch<sup>ie</sup><sub>166,168</sub> <sup>252</sup>Ch<sup>ie</sup><sub>166,168</sub>

$$\alpha_{2-33} = \frac{33 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{139}{138}\right)^{277}}}{2 \cdot (2 \cdot 8^2 - 1)} \frac{1}{112 - \frac{1}{32 \cdot 48 - \frac{36}{35 \cdot 13}}} = 1/137.035999111818$$

<sup>75</sup>As<sub>33,42</sub> <sup>84</sup>Kr<sub>36,48</sub> <sup>112</sup>Cd<sub>48,64</sub> <sup>226</sup>Ra\*<sub>88,138</sub> <sup>326,326,328</sup>Ch<sup>ie</sup><sub>128,129</sub> <sup>16,33,197,199</sup>Fy<sup>ie</sup><sub>136,137,138</sub> <sup>344,346,348</sup>Fy<sup>ie</sup><sub>208,209,210</sub> <sup>12,29</sup>Ch<sup>ie</sup><sub>139</sub> <sup>209=1,119</sup>Ch<sup>ie</sup><sub>209=1,119</sub>

$$\alpha_{2-36} = \frac{6^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{191}{190}\right)^{3 \cdot 127}}}{2 \cdot 138 + 1} \frac{1}{112 - \frac{1}{10 \cdot 7 \cdot 31 + \frac{13}{25 \cdot 23}}} = 1/137.035999111818$$

<sup>27</sup>Al<sub>13,14</sub> <sup>50,51</sup>V<sub>23,27,28</sub> <sup>53</sup>Cr<sub>24,29</sub> <sup>55</sup>Mn<sub>25,30</sub> <sup>54,56,57</sup>Fe<sub>26,28,30,31</sub> <sup>69,71</sup>Ga<sub>31,38,40</sub> <sup>89</sup>Rb<sub>39,50</sub> <sup>95</sup>Mo<sub>42,53</sub> <sup>108</sup>Pd<sub>46,62</sub>  
<sup>112,115,119,122</sup>Sn<sub>50,62,65,69,72</sub> <sup>127</sup>I<sub>53,74</sub> <sup>124,126,128,130</sup>Te<sub>52,72,74,76,78</sub> <sup>185,187</sup>Re<sub>75,110,112</sub> <sup>190</sup>Pt<sub>78,112</sub> <sup>8,36</sup>Mc<sup>ie</sup><sub>115=5,25</sub> <sup>173</sup>Mc<sup>ie</sup><sub>173</sub>

<sup>310,24,13</sup>Ch<sup>ie</sup><sub>124,125</sub> <sup>186,187</sup>Ch<sup>ie</sup><sub>186,187</sub> <sup>6,53,320</sup>Ch<sup>ie</sup><sub>127</sub> <sup>191,193</sup>Ch<sup>ie</sup><sub>191,193</sub> <sup>344,346,348</sup>Fy<sup>ie</sup><sub>136,137,138</sub> <sup>208,209,210</sup>Fy<sup>ie</sup><sub>208,209,210</sub> <sup>378,380,382</sup>Ch<sup>ie</sup><sub>150,151,152</sub> <sup>228,229,230</sup>Ch<sup>ie</sup><sub>228,229,230</sub>

$$\alpha_{2-37} = \frac{37 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 43}{5 \cdot 17}\right)^{9 \cdot 19}}}{3 \cdot 5 \cdot 19} \frac{1}{112 - \frac{1}{4 \cdot 3 \cdot 5 \cdot 13} + \frac{1}{5 \cdot 37^2 \cdot 149}}} = 1/137.035999111818$$

<sup>27</sup>Al<sub>13,14</sub> <sup>39</sup>K<sub>19,20</sub> <sup>85,87</sup>Rb<sub>37,48,50</sub> <sup>89</sup>Y<sub>39,50</sub> <sup>126</sup>Te<sub>52,74</sub> <sup>144,145,146,147,148,150</sup>Nd<sub>60,84,85,86,88,90</sub> <sup>210</sup>Po\*<sub>85,125</sub> <sup>222</sup>Rn\*<sub>86,136</sub>


<sup>284</sup>Nh<sup>ie</sup><sub>113</sub> <sup>172</sup>Tl<sup>ie</sup><sub>114=2,57</sub> <sup>2,149,300</sup>Ch<sup>ie</sup><sub>119=7,17,120</sub> <sup>179,180</sup>Ch<sup>ie</sup><sub>179,180</sub> <sup>22,17</sup>Ch<sup>ie</sup><sub>148=4,37,149</sub> <sup>226,225</sup>Ch<sup>ie</sup><sub>226,225</sub>

$$\alpha_{2-38} = \frac{2 \cdot 19 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{3 \cdot 19}{56}\right)^{113}}}{6 \cdot 7^2 - 1} \frac{1}{112 - \frac{1}{3 \cdot 73} + \frac{1}{30(8 \cdot 27 \cdot 17 + 1) - \frac{12}{13}}} = 1/137.035999111816$$

<sup>19</sup>F<sub>9,10</sub> <sup>39</sup>K<sub>19,20</sub> <sup>56</sup>Fe<sub>26,30</sub> <sup>64,66,68</sup>Zn<sub>30,34,36,38</sub> <sup>87,88</sup>Sr<sub>38,49,50</sub> <sup>125</sup>Te<sub>52,73</sub> <sup>180,181</sup>Ta<sub>73,107,108</sub> <sup>284</sup>Nh<sup>ie</sup><sub>113</sub> <sup>172</sup>Tl<sup>ie</sup><sub>114=6,19</sub> <sup>175</sup>Tl<sup>ie</sup><sub>175</sub>

$$\alpha_{2-125} = \frac{5 \cdot 5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{4294}{4293}\right)^{8587}}}{31^2} \frac{1}{112 - \frac{1}{2159481}}$$


$$= \frac{5 \cdot 5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{2 \cdot 19 \cdot 113}{81 \cdot 53}\right)^{31 \cdot (12 \cdot 23 + 1)}}}{31^2} \frac{1}{112 - \frac{1}{3 \cdot 101 \cdot (8 \cdot 81 \cdot 11 - 1)}}} = 1/137.035999111818$$



$$\alpha_{2-253} = \frac{253 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{28187}{28186}\right)^{56373}}}{1945} \frac{1}{112 - \frac{10411}{8 \times 10^{11}}}$$

$$11 \cdot 23 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left[\frac{71(36 \cdot 11 + 1)}{2 \cdot 17(36 \cdot 23 + 1)}\right]^{319 \cdot 23 \cdot 43}}$$

$$= \frac{1}{5(4 \cdot 97 + 1)} \frac{1}{112 - \frac{29(360 - 1)}{8 \times 10^{11}}} = 1/137.035999111818$$




$$\alpha_{2-269} = \frac{269 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{41655}{41654}\right)^{83309}}}{2068} \frac{1}{112 - 5.317 \times 10^{-9}}$$

$$(4 \cdot 67 + 1) \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{15 \cdot (6 \cdot (16 \cdot 29 - 1) - 1)}{2 \cdot 59 \cdot (6 \cdot 59 - 1)}\right)^{227 \cdot (6 \cdot 61 + 1)}}$$

$$= \frac{1}{4 \cdot 11 \cdot 47} \frac{1}{112 - \frac{13 \cdot (24 \cdot 17 + 1)}{10^{12}}}$$

$$= 1/137.035999111818$$



In above formulas, there are many amazing coincidences. As  $136=8 \times 17$  and  $138=6 \times 23$ , 17 and 23 both appear in  $\alpha_{1-1}$ ,  $\alpha_{1-17}$ ,  $\alpha_{1-22}$ ,  $\alpha_{1-23}$ ,  $\alpha_{1-25}$ ,  $\alpha_{1-59}$ ,  $\alpha_{1-103}$ ,  $\alpha_{1-133}$ ,  $\alpha_{2-17}$  and  $\alpha_{2-23}$ , 17 frequently appears in  $\alpha_1$  and 23 frequently appears in  $\alpha_2$ . 157 and 257 in  $\alpha_{1-50}$  should relate to  ${}_{100}\text{Fm}_{157}^*$ , 173 in  $\alpha_{1-16}$  should relate to  ${}_{112}\text{Cn}_{173}^*$ , and so on. As the factors in formulas of  $\alpha$  are reasonably assumed to relate to nuclides, some ideal extended elements such as  ${}_{136,137,138}\text{Fy}_{208,209,210}$  and  ${}_{169}\text{Ch}_{257}$  are predicted.

## 15. Radius of Electron and Proton

The classical electron radius  $r_e$  has been calculated very accurately. However, the proton charge radius  $r_p$  hasn't yet been determined precisely. Recent two experiments

measured  $r_p$  and had given the best results up to now which was  $r_p=0.833(19) \text{ fm}^9$  and  $r_p=0.831(19) \text{ fm}^{10}$ , and hence CODATA revised its recommended data of  $r_p$  to  $0.8414(19) \text{ fm}$ . Here we give our calculation results of  $r_e$  and  $r_p$ . And it seems there is  $\alpha_p$  similar to  $\alpha$ .  $\alpha_p$  could be called “the second fine-structure constant”.

Ratio of Bohr radius of hydrogen atom to classical electron radius:

$$\frac{a_0}{r_e} = \frac{1}{\alpha_c^2} = \frac{1}{\alpha_1 \alpha_2} = 112 \times \left( 168 - \frac{1}{3} + \frac{1}{2^2 \cdot 3 \cdot 47} - \frac{1}{2 \cdot 3 \cdot 29 \cdot 53 \cdot 59 - 79 / 47} \right) = 18788.865042381$$

$$r_e = \alpha_c^2 a_0 = \alpha_1 \alpha_2 a_0 = \frac{5.29177210903(80) \times 10^{-11} \text{ m}}{18788.865042381} = 2.81794032658(43) \text{ fm}$$

Comparable to CODATA recommended value  $r_e = 2.8179403262(13) \text{ fm}$  but more precise.

Ratio of Bohr radius of hydrogen atom to the proton charge radius should have the similar form, and is assumed to have the following hypothetical formulas:

$$\frac{a_0}{r_p} = \frac{1}{\alpha_{p/c}^2} = \frac{1}{\alpha_{p/1} \alpha_{p/2}} = 225 \cdot \left( 282 + \frac{1}{3} - \frac{1}{12 \cdot 47} + \frac{1}{6 \cdot 29 \cdot 53 \cdot 59 - 79 / 47} \right) = 63524.60147736$$


$$= 247 \cdot \left( 257 + \frac{1}{5} - \frac{1}{5 \cdot 13} + \frac{1}{30 \cdot (28 \cdot (2 \cdot 100 - 1) + 1) + \frac{8}{45}} \right)$$

$$= \left( 252 + \frac{1}{24} - \frac{1}{2 \cdot 17 \cdot 37} + \frac{1}{11 \cdot 13 \cdot 19 \cdot (2 \cdot 11 \cdot 19 + 1) + \frac{11}{20}} \right)^2 = 252.040872632515^2$$


$$r_p = \alpha_{p/c}^2 a_0 = \alpha_{p/1} \alpha_{p/2} a_0 = \frac{5.29177210903(80) \times 10^{-11} \text{ m}}{63524.60147736} = 0.833027202999(13) \text{ fm}$$

$\alpha_{p/c} \approx \alpha_{p/1} \approx \alpha_{p/2} \approx 252.04$ ,  $\alpha_p$  could be called the second fine-structure constant.

2019/12/19-23




$$\alpha_{p/1} = \frac{5 \cdot 3^2}{8 \cdot (2\pi)_{58}} \frac{1}{225 + \frac{1}{4 \cdot 112} - \frac{1}{5 \cdot 19 \cdot 83 \cdot 103 + \frac{19}{20}}} = 1 / 252.040872632515$$



$$\alpha_{p/2} = \frac{23 \cdot (2\pi)_{227}}{2 \cdot 9^2} \frac{1}{225 - \frac{1}{3 \cdot 16 \cdot 29 - \frac{71}{2 \cdot 67}}} = 1 / 252.040872632514$$

2020/1/2



$$\alpha_{p/2} = \frac{22 \cdot (2\pi)_{164}}{5 \cdot 31} \frac{1}{225 - \frac{1}{7 \cdot 137 - \frac{1}{197}}} = 1/252.040872632512 \quad 2020/1/3$$

<sup>47,48,49,50</sup><sub>22</sub>Ti <sup>56,57</sup><sub>26</sub>Fe <sup>69,71</sup><sub>31</sub>Ga <sup>89</sup><sub>39</sub>Y <sup>99,100</sup><sub>44</sub>Ru <sup>134,136,137,138</sup><sub>56</sub>Ba <sup>5-31</sup><sub>64</sub>Gd <sup>164</sup><sub>66</sub>Dy <sup>196,198</sup><sub>78</sub>Pt <sup>118,120</sup>  
<sup>197</sup><sub>79</sub>Au <sup>206,207,208</sup><sub>82</sub>Pb <sup>223</sup><sub>87</sub>Ra\* <sup>226</sup><sub>88</sub>Ra\* <sup>310,312</sup><sub>138</sub>Ch<sup>ie</sup> <sup>326,326/328</sup><sub>128,129</sub>Ch<sup>ie</sup> <sup>12-31</sup><sub>147</sub>Ch<sup>ie</sup> <sup>155=5-31,156=12-13</sup><sub>147</sub>Ch<sup>ie</sup> <sup>8-49,2197</sup><sub>164</sub>Ch<sup>ie</sup> <sup>32-13</sup><sub>164</sub>Ch<sup>ie</sup>

$$\alpha_{p/2} = \frac{21 \cdot (2\pi)_{126}}{2^2 \cdot 37} \frac{1}{225 - \frac{1}{16 \cdot 29} + \frac{1}{20 \cdot 13^2 \cdot 179 + \frac{8}{17}}} = 1/252.040872632515 \quad 2020/1/3$$

<sup>45</sup><sub>21</sub>Sc <sup>63,65</sup><sub>29</sub>Cu <sup>85,87</sup><sub>37</sub>Rb <sup>126</sup><sub>52</sub>Te <sup>148</sup><sub>60</sub>Nd <sup>169</sup><sub>69</sub>Tm <sup>179</sup><sub>72</sub>Hf <sup>16-13</sup><sub>82</sub>Pb <sup>298,300</sup><sub>119,120</sub>Ch<sup>ie</sup> <sup>312,314</sup><sub>125,126</sub>Ch<sup>ie</sup> <sup>426</sup><sub>169</sub>Ch<sup>ie</sup>

## 16. Direct Relationships of 2π with Nuclides

In Chen's formulas of the fine-structure constant, there are 2π-e formulas, in which k gets certain numbers and relate to nucleon numbers of some nuclides. So in the end of this paper we feel curious about whether 2π directly relate to nuclides.

$$2\pi = 6.2831853 \dots \approx \frac{4 \cdot 157}{100} = 6.28 \approx \frac{3 \cdot 7 \cdot 44 \cdot 68}{100^2} = 6.2832 \quad \begin{matrix} 7 \\ 3 \end{matrix} \text{Li} \quad \begin{matrix} 100 \\ 44 \end{matrix} \text{Ru} \quad \begin{matrix} 157 \\ 64 \end{matrix} \text{Gd} \quad \begin{matrix} 168 \\ 68 \end{matrix} \text{Er} \quad \begin{matrix} 257 \\ 100 \end{matrix} \text{Fm}^* \quad \begin{matrix} 400 \\ 157 \end{matrix} \text{Ch}^{\text{ie}}_{243}$$

$$2\pi \approx \frac{4 \cdot 157}{100} = \frac{157}{25} = 6.28 \quad \begin{matrix} 55 \\ 25 \end{matrix} \text{Mn} \quad \begin{matrix} 100 \\ 44 \end{matrix} \text{Ru} \quad \begin{matrix} 157 \\ 64 \end{matrix} \text{Gd} \quad \begin{matrix} 118,119,120 \\ 50 \end{matrix} \text{Sn} \quad \begin{matrix} 168 \\ 68 \end{matrix} \text{Er} \quad \begin{matrix} 169 \\ 69 \end{matrix} \text{Tm} \quad \begin{matrix} 185,187 \\ 75 \end{matrix} \text{Re} \quad \begin{matrix} 200 \\ 80 \end{matrix} \text{Hg} \quad \begin{matrix} 257 \\ 100 \end{matrix} \text{Fm}^* \quad \begin{matrix} 258 \\ 101 \end{matrix} \text{Md}^* \quad \begin{matrix} 259 \\ 102 \end{matrix} \text{No}^* \quad \begin{matrix} 312,2157 \\ 125,126 \end{matrix} \text{Ch}^{\text{ie}}_{117,188} \quad \begin{matrix} 400 \\ 157 \end{matrix} \text{Ch}^{\text{ie}}_{243}$$

$$2\pi \approx \frac{16 \cdot 3 \cdot 7 \cdot 11 \cdot 17}{100^2} = \frac{48 \cdot 7 \cdot 11 \cdot 17}{100^2} = \frac{3 \cdot 7 \cdot 44 \cdot 68}{100^2} = \frac{3 \cdot 112 \cdot 11 \cdot 17}{100^2} = \frac{2 \cdot 168 \cdot 11 \cdot 17}{100^2} = \frac{2 \cdot 3 \cdot 7 \cdot 11 \cdot 136}{100^2} = \dots = 6.2832$$

<sup>7</sup><sub>3</sub>Li <sup>20-22</sup><sub>10</sub>Ne <sup>23</sup><sub>11</sub>Na <sup>45</sup><sub>21</sub>Sc <sup>46,47,49,50</sup><sub>22</sub>Ti <sup>24,25,27,28</sup><sub>22</sub>Ti <sup>61</sup><sub>28</sub>Ni <sup>55</sup><sub>25</sub>Mn <sup>54,56</sup><sub>26</sub>Fe <sup>78,80</sup><sub>34</sub>Se <sup>98,100</sup><sub>42</sub>Mo <sup>100</sup><sub>44</sub>Ru <sup>112</sup><sub>48</sub>Cd <sup>136,137,138</sup><sub>56</sub>Ba <sup>80,81,82</sup><sub>68</sub>Er <sup>185,1117</sup><sub>75</sub>Re <sup>209</sup><sub>84</sub>Po\* <sup>222</sup><sub>86</sub>Rn\* <sup>223</sup><sub>87</sub>Fa\* <sup>226</sup><sub>88</sub>Ra\* <sup>227</sup><sub>89</sub>Ac\* <sup>278+7</sup><sub>112</sub>Cn <sup>344,346,348</sup><sub>117</sub>Fy <sup>208,209,210</sup>

$$2\pi \approx \frac{44}{7} = \frac{2 \cdot 22}{7} = 6.2857 \dots \quad \begin{matrix} 50 \\ 22 \end{matrix} \text{Ti} \quad \begin{matrix} 61 \\ 28 \end{matrix} \text{Ni} \quad \begin{matrix} 100 \\ 44 \end{matrix} \text{Ru} \quad \begin{matrix} 136,137,138 \\ 56 \end{matrix} \text{Ba} \quad \begin{matrix} 226 \\ 88 \end{matrix} \text{Ra}^* \quad \begin{matrix} 294 \\ 118 \end{matrix} \text{Og}^{\text{ie}} \quad \begin{matrix} 8-44 \\ 140 \end{matrix} \text{Ch}^{\text{ie}}_{212} \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}^{\text{ie}}_{226}$$

$$2\pi \approx \frac{201}{32} = \frac{3 \cdot 67}{32} = 6.2812 \dots \quad \begin{matrix} 32 \\ 16 \end{matrix} \text{S} \quad \begin{matrix} 59 \\ 16 \end{matrix} \text{Co} \quad \begin{matrix} 67 \\ 32 \end{matrix} \text{Zn} \quad \begin{matrix} 112 \\ 48 \end{matrix} \text{Cd} \quad \begin{matrix} 117 \\ 50 \end{matrix} \text{Sn} \quad \begin{matrix} 128,134 \\ 54 \end{matrix} \text{Xe} \quad \begin{matrix} 134 \\ 56 \end{matrix} \text{Ba} \quad \begin{matrix} 165 \\ 67 \end{matrix} \text{Ho} \quad \begin{matrix} 201 \\ 80 \end{matrix} \text{Hg} \quad \begin{matrix} 332 \\ 121 \end{matrix} \text{Ch}^{\text{ie}}_{131} \quad \begin{matrix} 402 \\ 158 \end{matrix} \text{Ch}^{\text{ie}}_{244}$$

$$2\pi \approx \frac{245}{39} = \frac{5 \cdot 7^2}{3 \cdot 13} = 6.2820 \dots \quad \begin{matrix} 7 \\ 3 \end{matrix} \text{Li} \quad \begin{matrix} 27 \\ 13 \end{matrix} \text{Al} \quad \begin{matrix} 54,56 \\ 26 \end{matrix} \text{Fe} \quad \begin{matrix} 89 \\ 39 \end{matrix} \text{Y} \quad \begin{matrix} 79,81 \\ 35 \end{matrix} \text{Br} \quad \begin{matrix} 113,115 \\ 49 \end{matrix} \text{In} \quad \begin{matrix} 24-13,314 \\ 125,126 \end{matrix} \text{Ch}^{\text{ie}}_{187,188}$$

$$2\pi \approx \frac{289}{46} = \frac{17^2}{2 \cdot 23} = 6.2826 \dots \quad \begin{matrix} 3-17 \\ 23 \end{matrix} \text{V} \quad \begin{matrix} 78,80 \\ 34 \end{matrix} \text{Se} \quad \begin{matrix} 6-17 \\ 46 \end{matrix} \text{Pd} \quad \begin{matrix} 168 \\ 68 \end{matrix} \text{Er} \quad \begin{matrix} 169 \\ 69 \end{matrix} \text{Tm} \quad \begin{matrix} 136,137,138 \\ 56 \end{matrix} \text{Ba} \quad \begin{matrix} 11-17 \\ 75 \end{matrix} \text{Re} \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}^{\text{ie}}_{226}$$

<sup>223</sup><sub>87</sub>Fa\* <sup>226</sup><sub>88</sub>Ra\* <sup>227</sup><sub>89</sub>Ac\* <sup>238</sup><sub>92</sub>U\* <sup>17-17</sup><sub>114</sub>Fl<sup>ie</sup> <sup>344,346,348</sup><sub>117</sub>Fy <sup>208,209,210</sup> <sup>22-17</sup><sub>148=4 \cdot 37</sub>Ch<sup>ie</sup><sub>226</sub>

$$2\pi \approx \frac{333}{53} = \frac{9 \cdot 37}{53} = 6.2830 \dots \quad \begin{matrix} 85,87 \\ 37 \end{matrix} \text{Rb} \quad \begin{matrix} 3 \cdot 37=111 \\ 48 \end{matrix} \text{Cd} \quad \begin{matrix} 127 \\ 53 \end{matrix} \text{I} \quad \begin{matrix} 180,184,189 \\ 74 \end{matrix} \text{W} \quad \begin{matrix} 222 \\ 86 \end{matrix} \text{Rn}^* \quad \begin{matrix} 269 \\ 106 \end{matrix} \text{Sg}^* \quad \begin{matrix} 280 \\ 111 \end{matrix} \text{Rg}^* \quad \begin{matrix} 22-17 \\ 148 \end{matrix} \text{Ch}^{\text{ie}}_{226}$$

$$2\pi \approx \frac{377}{60} = \frac{13 \cdot 29}{4 \cdot 3 \cdot 5} = 6.2833 \dots \quad \begin{matrix} 24,25,26 \\ 12 \end{matrix} \text{Mg} \quad \begin{matrix} 28,29,30 \\ 14 \end{matrix} \text{Si} \quad \begin{matrix} 31 \\ 15 \end{matrix} \text{P} \quad \begin{matrix} 54,56 \\ 26 \end{matrix} \text{Fe} \quad \begin{matrix} 63,65 \\ 29 \end{matrix} \text{Cu} \quad \begin{matrix} 116,120 \\ 50 \end{matrix} \text{Sn} \quad \begin{matrix} 140,142 \\ 58 \end{matrix} \text{Ce} \quad \begin{matrix} 144,145,146,148,150 \\ 60 \end{matrix} \text{Nd} \quad \begin{matrix} 200 \\ 80 \end{matrix} \text{Hg} \quad \begin{matrix} 223 \\ 87 \end{matrix} \text{Fa}^* \quad \begin{matrix} 24-13,314 \\ 125,126 \end{matrix} \text{Ch}^{\text{ie}}_{187,188}$$

$$2\pi \approx \frac{465}{74} = \frac{30 \cdot 31}{4 \cdot 37} = 6.2837 \dots \quad \begin{matrix} 31 \\ 15 \end{matrix} \text{P} \quad \begin{matrix} 67 \\ 30 \end{matrix} \text{Zn} \quad \begin{matrix} 69,71 \\ 31 \end{matrix} \text{Ga} \quad \begin{matrix} 6-31 \\ 74 \end{matrix} \text{W} \quad \begin{matrix} 85,67 \\ 37 \end{matrix} \text{Rb} \quad \begin{matrix} 4-37 \\ 60 \end{matrix} \text{Nd} \quad \begin{matrix} 157 \\ 64 \end{matrix} \text{Gd} \quad \begin{matrix} 243 \\ 95 \end{matrix} \text{Am}^* \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}^{\text{ie}}_{226}$$

$$2\pi \approx \frac{509}{81} = \frac{2 \cdot 3 \cdot 5 \cdot 17 - 1}{9^2} = 6.2839 \dots \quad \begin{matrix} 19 \\ 9 \end{matrix} \text{F} \quad \begin{matrix} 35,37 \\ 17 \end{matrix} \text{Cl} \quad \begin{matrix} 64,70 \\ 30 \end{matrix} \text{Zn} \quad \begin{matrix} 80,82 \\ 34 \end{matrix} \text{Se} \quad \begin{matrix} 136,137,138 \\ 56 \end{matrix} \text{Ba} \quad \begin{matrix} 203,205 \\ 81 \end{matrix} \text{Tl} \quad \begin{matrix} 210 \\ 85 \end{matrix} \text{At}^* \quad \begin{matrix} 3-81 \\ 95 \end{matrix} \text{Am}^* \quad \begin{matrix} 344,346,348 \\ 136,137,138 \end{matrix} \text{Fy}^* \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}^{\text{ie}}_{226} \quad \begin{matrix} 400 \\ 157 \end{matrix} \text{Ch}^{\text{ie}}_{243}$$

$$2\pi \approx \frac{622}{99} = \frac{4 \cdot (24 \cdot 13 - 1)}{9 \cdot 22} = 6.2828 \dots \quad \begin{matrix} 23 \\ 11 \end{matrix} \text{Na} \quad \begin{matrix} 27 \\ 12 \end{matrix} \text{Al} \quad \begin{matrix} 46,48,49 \\ 22 \end{matrix} \text{Ti} \quad \begin{matrix} 50,52,54 \\ 24 \end{matrix} \text{Cr} \quad \begin{matrix} 54,56,58 \\ 26 \end{matrix} \text{Fe} \quad \begin{matrix} 99 \\ 44 \end{matrix} \text{Ru} \quad \begin{matrix} 167 \\ 68 \end{matrix} \text{Er} \quad \begin{matrix} 252 \\ 99 \end{matrix} \text{Es}^* \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}^{\text{ie}}_{226}$$

$$2\pi \approx \frac{2 \cdot 355}{113} = \frac{4 \cdot 5 \cdot 71}{2 \cdot 113} = 6.2831858 \dots \quad \begin{matrix} 71 \\ 31 \end{matrix} \text{Ga} \quad \begin{matrix} 112,113 \\ 48 \end{matrix} \text{Cd} \quad \begin{matrix} 113,115 \\ 49 \end{matrix} \text{In} \quad \begin{matrix} 120,122 \\ 50 \end{matrix} \text{Sn} \quad \begin{matrix} 2-71 \\ 60 \end{matrix} \text{Nd} \quad \begin{matrix} 171 \\ 70 \end{matrix} \text{Yb} \quad \begin{matrix} 175 \\ 71 \end{matrix} \text{Lu} \quad \begin{matrix} 186 \\ 74 \end{matrix} \text{W} \quad \begin{matrix} 187 \\ 75 \end{matrix} \text{Re} \quad \begin{matrix} 188,189 \\ 76 \end{matrix} \text{Os} \quad \begin{matrix} 226 \\ 88 \end{matrix} \text{Ra}^* \quad \begin{matrix} 232 \\ 90 \end{matrix} \text{Th}^* \quad \begin{matrix} 4-71 \\ 113 \end{matrix} \text{Nh}^{\text{ie}} \quad \begin{matrix} 358/360 \\ 142=2 \cdot 71 \end{matrix} \text{Ch}^{\text{ie}} \quad \begin{matrix} 22-17 \\ 148=4 \cdot 37 \end{matrix} \text{Ch}^{\text{ie}}_{226} \quad \begin{matrix} 6-71 \\ 169 \end{matrix} \text{Ch}^{\text{ie}}_{257} \quad 2020/1/8-10$$

The approximate rational numbers of  $2\pi$  (could be called  $2\pi$  formulas) relate to nuclides marvelously. This means  $2\pi$  (along with  $2\pi$ -e formula) plays important roles in atomic nuclei, and acts as a rational number rather than an irrational number in the world of atomic nuclei.

### 17. Correlations among $\alpha$ , $2\pi$ and nuclides

Some Chen's formulas of the fine-structure constant and  $2\pi$  formulas correlate with each others with the same factors and all together relate to the same nuclides. For example,  $\alpha_{1-50}$  and  $2\pi \approx 4 \times 157/100$  have the same 157 and 100 factors,  $\alpha_{1-50}$  and  $2\pi \approx 3 \times 7 \times 44 \times 68/100^2$  have the same 100, 7, 11 and 16 factors, and they relate to the same corresponding nuclides. They also have common factors with  $\alpha_{1-7}$  and  $\alpha_{2-13}$  which should relate to  $2\pi \approx 5 \times 7^2/3/13$  and  $2\pi \approx 13 \times 29/4/3/5$ .

$$\alpha_{1-9} = \frac{47}{3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{10}{9}\right)^{19}} 112 + \frac{1}{4 \cdot 17}} \frac{1}{2(8 \cdot 9 \cdot 37 - 1) + \frac{3 \cdot 17}{157}}$$

$$\alpha_{1-50} = \frac{2 \cdot 257}{100 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14 \cdot 13}{181}\right)^{3 \cdot 11^2}} 112 + \frac{1}{29 \cdot 61 + \frac{157}{16 \cdot 11}}}$$

$$2\pi \approx \frac{4 \cdot 157}{100} = \frac{157}{25} = 6.28 \quad 2\pi \approx \frac{3 \cdot 7 \cdot 44 \cdot 68}{100^2} = \frac{3 \cdot 7 \cdot 11 \cdot 17}{25^2} = 6.2832$$

$$\alpha_{1-7} = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}} 112 + \frac{1}{75^2}} \quad \alpha_{2-13} = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{279}{278}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}}$$

$$2\pi \approx \frac{245}{39} = \frac{5 \cdot 7^2}{3 \cdot 13} = 6.2820 \dots \quad 2\pi \approx \frac{377}{60} = \frac{13 \cdot 29}{4 \cdot 3 \cdot 5} = 6.2833 \dots$$

$\alpha_{1-22}$  relates to  $2\pi \approx 2 \times 22/7$ ,  $2\pi \approx 17^2/7/23$  and  $2\pi \approx 2 \times 355/113$  as follows. And  $2\pi \approx 17^2/7/23$  also relates to  $\alpha_{1-1}$ ,  $\alpha_{1-17}$ ,  $\alpha_{1-22}$ ,  $\alpha_{1-23}$ ,  $\alpha_{1-25}$ ,  $\alpha_{1-59}$ ,  $\alpha_{1-103}$ ,  $\alpha_{1-133}$ ,  $\alpha_{2-17}$  and  $\alpha_{2-23}$ , in which both 17 and 23 factors appear.

$$\alpha_{1-22} = \frac{113}{22 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{27 \cdot 29}{2 \cdot 17 \cdot 23}\right)^{5 \cdot (2 \cdot 157 - 1)}} 112 + \frac{1}{2 \cdot [2 \cdot 3 \cdot 17 \cdot (10 \cdot 19 + 1) + 1] + \frac{29}{49}}}$$

$$2\pi \approx \frac{2 \cdot 22}{7} = 6.2857 \dots, \quad 2\pi \approx \frac{17^2}{2 \cdot 23} = 6.2826 \dots, \quad 2\pi \approx \frac{2 \cdot 355}{113} = \frac{4 \cdot 5 \cdot 71}{2 \cdot 113} = 6.2831858 \dots$$



$\alpha_{1-13}$  and  $\alpha_{1-43}$  relate to  $2\pi \approx 3 \times 67/32$ ,  $2\pi \approx 5 \times 7^2/39$ ,  $2\pi \approx 17^2/46$  and others as follows.

$$\alpha_{1-13} = \frac{67}{13 \cdot e^2 \frac{e^2}{\binom{2}{1}^3} \frac{e^2}{\binom{3}{2}^5} \dots \frac{e^2}{\left(\frac{47}{2 \cdot 23}\right)^{3 \cdot 31}}} \frac{1}{112 + \frac{1}{137} - \frac{1}{6(2 \cdot 27 \cdot 59 + 1) + \frac{9}{50}}}$$

$$\alpha_{1-43} = \frac{13 \cdot 17}{43 \cdot e^2 \frac{e^2}{\binom{2}{1}^3} \frac{e^2}{\binom{3}{2}^5} \dots \frac{e^2}{\left(\frac{3 \cdot 67}{200}\right)^{401}}} \frac{1}{112 + \frac{1}{8 \cdot (12 \cdot 83 + 1) + \frac{4}{3 \cdot 13}}}$$

$$2\pi \approx \frac{3 \cdot 67}{32}, 2\pi \approx \frac{5 \cdot 7^2}{3 \cdot 13}, 2\pi \approx \frac{289}{46} = \frac{17^2}{2 \cdot 23}, 2\pi \approx \frac{13 \cdot 29}{60}, 2\pi \approx \frac{30 \cdot 31}{4 \cdot 37}$$

$\alpha_{1-11}$ ,  $\alpha_{1-36}$ ,  $\alpha_{2-24}$ ,  $\alpha_{2-23}$ ,  $\alpha_{2-37}$  and  $\alpha_{2-125}$  relate to  $2\pi \approx 9 \times 37/53$ ,  $2\pi \approx 15 \times 31/2/37$  and  $2\pi \approx (30 \times 17 - 1)/81$  as follows.

$$\alpha_{1-11} = \frac{57}{11 \cdot e^2 \frac{e^2}{\binom{2}{1}^3} \frac{e^2}{\binom{3}{2}^5} \dots \frac{e^2}{\left(\frac{19}{18}\right)^{37}}} \frac{1}{112 + \frac{1}{35} - \frac{1}{88 \cdot 41 - \frac{5 \cdot 53}{22 \cdot 13}}}$$

$$\alpha_{1-36} = \frac{5 \cdot 37}{6^2 \cdot e^2 \frac{e^2}{\binom{2}{1}^3} \frac{e^2}{\binom{3}{2}^5} \dots \frac{e^2}{\left(\frac{3 \cdot 79}{4 \cdot 59}\right)^{11 \cdot 43}}} \frac{1}{112 + \frac{1}{5 \cdot (31 \cdot 42 - 1) + \frac{3 \cdot 31}{14 \cdot 13}}}$$

$$\alpha_{2-24} = \frac{2^2 \cdot 6 \cdot e^2 \frac{e^2}{\binom{2}{1}^3} \frac{e^2}{\binom{3}{2}^5} \dots \frac{e^2}{\left(\frac{63}{2 \cdot 31}\right)^{125}}}{5 \cdot 37} \frac{1}{112 - \frac{1}{257} + \frac{1}{10 \cdot (12 \cdot 13 \cdot 83 + 1) + \frac{23}{81}}}$$

$$\alpha_{2-23} = \frac{23 \cdot e^2 \frac{e^2}{\binom{2}{1}^3} \frac{e^2}{\binom{3}{2}^5} \dots \frac{e^2}{\left(\frac{2 \cdot 81}{7 \cdot 23}\right)^{17 \cdot 19}}}{3 \cdot 59} \frac{1}{112 - \frac{1}{2 \cdot (40 \cdot 23 - 1) + \frac{9}{32 \cdot 10}}}$$


$$\alpha_{2-37} = \frac{37 \cdot e^2 \frac{e^2}{\binom{2}{1}^3} \frac{e^2}{\binom{3}{2}^5} \dots \frac{e^2}{\left(\frac{2 \cdot 43}{5 \cdot 17}\right)^{9 \cdot 19}}}{3 \cdot 5 \cdot 19} \frac{1}{112 - \frac{1}{4 \cdot 3 \cdot 5 \cdot 13} + \frac{1}{5 \cdot 37^2 \cdot 149}}$$

$$\alpha_{2-125} = \frac{5 \cdot 5^2 \cdot e^2 \frac{e^2}{\binom{2}{1}^3} \frac{e^2}{\binom{3}{2}^5} \dots \frac{e^2}{\left(\frac{2 \cdot 19 \cdot 113}{81 \cdot 53}\right)^{31 \cdot (12 \cdot 23 + 1)}}}{31^2} \frac{1}{112 - \frac{1}{101 \cdot (20 \cdot (12 \cdot 89 + 1) + 1)}}$$

$$2\pi \approx \frac{9 \cdot 37}{53} = 6.2830 \dots \quad 2\pi \approx \frac{15 \cdot 31}{2 \cdot 37} = 6.2837 \dots \quad 2\pi \approx \frac{30 \cdot 17 - 1}{81} = 6.2839 \dots$$

## 18. Chen's Mathematic Shell Model of Nuclides

In overall, there are multi-correlations among  $\alpha$ ,  $2\pi$  and nuclides. It seems there should be a mathematical shell model of nuclides, in which the core is  $2\pi$  formulas and the middle layer is  $2\pi$ -e formulas and the outer layer is Chen's formulas of the fine-structure constant (**Fig. 9**). The nucleon numbers, stability and abundance of nuclides are regulated by these formulas, especially by their integer factors.



Chen's Mathematic Shell Model of Nuclides

Dr. Gang Chen (2020/1/12-13)

**Fig. 9**

## 19. Ideal Extended Elements

In the deduction of Chen's formulas of the fine-structure constant, it was reasonably assumed the factors in them related to nucleon numbers of nuclides, and it seems this assumption is quite correct. So by somewhat correlation and decoding methodology, all 119<sup>th</sup> to 170<sup>th</sup> ideal extended elements were predicted (**Table 7**). In addition, nuclides can even relate to naked  $2\pi$ 's approximate rational numbers ( $2\pi$  formulas). Some typical examples of correlations of ideal extended elements with formulas of  $\alpha$  and  $2\pi$  are listed as follows.

**Example 1:** Correlations of 100, 121, 125, 126, 157, 257, 169, *et al.*

$$\alpha_{1-9} = \frac{47}{3^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{10}{9}\right)^{19}}} \frac{1}{112 + \frac{1}{4 \cdot 17} - \frac{1}{2(8 \cdot 9 \cdot 37 - 1) + \frac{3 \cdot 17}{157}}}$$

$$\alpha_{1-50} = \frac{2 \cdot 257}{100 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{14 \cdot 13}{181}\right)^{3 \cdot 11^2}}} \frac{1}{112 + \frac{1}{29 \cdot 61 + \frac{157}{16 \cdot 11}}}$$

$$\alpha_{2-24} = \frac{2^2 \cdot 6 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{63}{62}\right)^{125}}}{5 \cdot 37} \frac{1}{112 - \frac{1}{257} + \frac{1}{10 \cdot (12 \cdot 13 \cdot 83 + 1) + \frac{23}{81}}}$$

$$2\pi \approx \frac{4 \cdot 157}{100} \quad 2\pi \approx \frac{16 \cdot 3 \cdot 7 \cdot 11 \cdot 17}{100} \quad 2\pi \approx \frac{4 \cdot 11}{7} \quad 2\pi \approx \frac{17^2}{2 \cdot 23} \quad 2\pi \approx \frac{30 \cdot 31}{4 \cdot 37} \quad 2\pi \approx \frac{4 \cdot 5 \cdot 71}{2 \cdot 113}$$

${}^{100}_{44}Ru_{56}$   ${}^{168,169}_{68,69}Tm_{100}$   ${}^{257}_{100}Fm_{157}^*$   ${}^{302}_{121=11^2}Ch_{181}^{ie}$   ${}^{24-13,2-157}_{125,126}Ch_{117,4-47}^{ie}$   ${}^{2-11-17}_{148=4-37}Ch_{226}^{ie}$   ${}^{400,402}_{157,158}Ch_{3-81,4-61}^{ie}$   ${}^{6-71}_{169}Ch_{257}^{ie}$

**Example 2:** Correlations of 83, 126, 84, 125, 209, 112, 173, 285, 115 and 137

$$\alpha_{1-16} = \frac{83}{4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{17}{16}\right)^{33}} 112 + \frac{1}{28} - \frac{1}{6 \cdot (18 \cdot 41 + 1) + \frac{173}{2 \cdot (2 \cdot 75 - 1)}}$$

$$\alpha_{1-25} = \frac{3 \cdot 43}{5^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{35}{34}\right)^{3 \cdot 23}} 112 + \frac{1}{11 \cdot 19} - \frac{1}{13^2(2 \cdot 136 - 1) + \frac{11}{25}}$$

$$\alpha_{1-32} = \frac{15 \cdot 11}{2 \cdot 2^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{41}{40}\right)^{81}} 112 + \frac{1}{25 \cdot 29} - \frac{5 \cdot 83}{19 \cdot 23}$$

$$\alpha_{2-10} = \frac{10 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{5 \cdot 21}{8 \cdot 13}\right)^{11 \cdot 19}}}{77} \frac{1}{112 - \frac{1}{3 \cdot 14 \cdot 19} + \frac{1}{14 \cdot (4 \cdot 27 \cdot (2 \cdot 15 \cdot 19 + 1) - 1)}}$$

$$\alpha_{2-32} = \frac{2 \cdot 4^2 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{42}{41}\right)^{83}}}{13 \cdot 19} \frac{1}{112 - \frac{1}{11 \cdot 13} + \frac{1}{6 \cdot 37 \cdot (5 \cdot 210 - 1) + \frac{10}{11}}}$$

$$\alpha_{1-13} = \frac{67}{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{47}{2 \cdot 23}\right)^{3 \cdot 31}} 112 + \frac{1}{137} - \frac{1}{6(2 \cdot 27 \cdot 59 + 1) + \frac{9}{50}}$$

$$\alpha_{1-17} = \frac{2^2 \cdot 22}{17 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{21}{20}\right)^{41}} 112 + \frac{1}{137} - \frac{1}{2 \cdot 19 \cdot 23 \cdot 59 - \frac{30}{100}}$$

$$\alpha_{1-27} = \frac{139}{27 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{67}{66}\right)^{7 \cdot 19}} 112 + \frac{1}{11 \cdot 47 + \frac{18}{23} + \frac{1}{6 \cdot 23 \cdot 137}}$$

$$\alpha_{2-18} = \frac{18 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{38}{37}\right)^{75}}}{139} \frac{1}{112 - \frac{1}{2 \cdot 47} + \frac{1}{2 \cdot 31 \cdot (16 \cdot 17 - 1) + \frac{83}{137}}}$$

$$2\pi \approx \frac{16 \cdot 3 \cdot 7 \cdot 11 \cdot 17}{100} \approx \frac{4 \cdot 11}{7} \approx \frac{17^2}{2 \cdot 23} \quad \begin{matrix} 8-17,137,6-23 \\ 56 \end{matrix} Ba_{80,81,82} \quad \begin{matrix} 11-19 \\ 83 \end{matrix} D_{126}^* \quad \begin{matrix} 209 \\ 84 \end{matrix} Po_{125}^* \quad \begin{matrix} 210 \\ 85 \end{matrix} At_{125}^* \quad \begin{matrix} 15-19 \\ 112 \end{matrix} Cn_{173}^* \quad \begin{matrix} 2-173 \\ 137 \end{matrix} Fy_{209}^{ie}$$

**Table 7.** Correlations of Ideal Extended Elements (IEE) with Formulas of  $\alpha$  and  $2\pi$ .

IEE	Page	$\alpha$	$2\pi$
<sup>113</sup> Nh <sub>171</sub>	10 19 21 28 29 31	$\alpha_c^2 \alpha_{1-5,7} \alpha_{2-22,23,31,37,38,253}$	$2\pi \approx 4 \times 355/226$
<sup>114</sup> Fl <sub>175</sub>	19 23 28 31	$\alpha_{1-11,133,155} \alpha_{2-37,38}$	$2\pi \approx 17^2/46$
<sup>115</sup> Mc <sub>173</sub>	20 21 25 31	$\alpha_{1-1,16,23} \alpha_{2-5}$	$2\pi \approx 17^2/46$
<sup>116</sup> Lv <sub>177</sub> <sup>117</sup> Ts <sub>177</sub>	10 20 22 27 31	$\alpha_{1-13,59} \alpha_{2-23} 1/\alpha_c^2$	$2\pi \approx 622/99$
<sup>118</sup> Og <sub>176</sub>	20 22 23 27	$\alpha_{1-17,20,50,59,133} \alpha_{2-19,269}$	$2\pi \approx 44/7$
<sup>119-122</sup> Ch <sub>179-182</sub>	21-23 28 31 37 39	$\alpha_{1-23,29,50,170} \alpha_{2-37} \alpha_{p/2} c_{au}$	$2\pi \approx 44/7$ et al.
<sup>123</sup> Ch <sub>183/185</sub>	19 20 21 25 28	$\alpha_{1-4,11,16,17,31} \alpha_{2-1,4,5,9,32}$	$2\pi \approx 333/53 \approx 465/74$ et al



The relationships between elements and ideal extended elements (the frontier of elements) and an overall picture of them were depicted as above.


## 21. Some Supplements

### Supplement 1:

$$2\pi \approx \frac{16 \cdot 3 \cdot 7 \cdot 11 \cdot 17}{100^2} = \frac{3 \cdot 7 \cdot 44 \cdot 68}{100^2} = \frac{3 \cdot 112 \cdot 11 \cdot 17}{100^2} = \frac{2 \cdot 168 \cdot 11 \cdot 17}{100^2} = \frac{2 \cdot 3 \cdot 7 \cdot 11 \cdot 136}{100^2} = \dots = 6.2832$$

Refer to Section 16; Supplements:  ${}_{16}^{32,33,34}O_{16,17,18}$  and some of the follows


$$2\pi \approx \frac{9 \cdot 7 \cdot 127 \cdot (4 \cdot 13 \cdot 151 + 1)}{10^7} = \frac{63 \cdot 127 \cdot (52 \cdot 151 + 1)}{10 \cdot 100^3} = \frac{63 \cdot 127 \cdot (2 \cdot 3 \cdot 7 \cdot 11 \cdot 17 - 1)}{10 \cdot (8 \cdot 125)^2} = \dots = 6.2831853$$



2020/2/11-12


### Supplement 2:

$$2\pi \approx \frac{509}{81} = \frac{4 \cdot 127 + 1}{9^2} \quad 2\pi \approx \frac{201}{32} = \frac{3 \cdot 67}{32} \quad 2\pi \approx \frac{333}{53} = \frac{9 \cdot 37}{53}$$



$$2\pi \approx \frac{622}{99} = \frac{4 \cdot (310 + 1)}{9 \cdot 22} \quad 2\pi \approx \frac{465}{74} = \frac{30 \cdot 31}{4 \cdot 37} \quad 2\pi \approx \frac{44}{7} = \frac{2 \cdot 22}{7}$$

$$2\pi \approx \frac{245}{39} = \frac{5 \cdot 7^2}{3 \cdot 13} \quad 2\pi \approx \frac{289}{46} = \frac{17^2}{2 \cdot 23} \quad 2\pi \approx \frac{377}{60} = \frac{13 \cdot 29}{4 \cdot 3 \cdot 5} \quad 2\pi \approx \frac{628}{100} = \frac{4 \cdot 157}{100}$$



2020/2/12

### Supplement 3:

**Table 8.** Relationships of factors in  $\alpha_{1-7}$  and  $\alpha_{2-13}$  with primordial nuclides (2020/2/16-17).

Nuclides	${}_3Li_4$	${}_{29}Cu_{34}$	${}_{31}Ga_{40}$	${}_{64}Gd_{92}$	${}_{75}Re_{112}$
A	7	65=5×13	71	156=12×13	187=11×17
PN before	5	70	78	209	252
PN all	285	285	285	285	285
Ratios	1/57	14/57	26/95	11/15	84/95

- 3, 29, 31, 64, 75 and 112 are factors in  $\alpha_1$  and  $\alpha_2$ .
- PN: primordial nuclides; PN all: usually regarded as 286.
- Nucleon number 285 of  ${}_{112}Cn_{173}$  would relate to PN all, or PN all should be 285 rather than 286.
- ${}^{235}U_{143}$  should not be a primordial nuclide, its relative stability (but not much stable) should come from relative stable nucleon numbers 92=96-4 and 143=11×13, so number of PN would become 285 from 286.



**Supplement 6: Other formulas of the speed of light  $c_{au}$**

$$c_{au} = \frac{1}{\sqrt{\alpha_{1-9/11}\alpha_{2-20/25}}}$$

$$= \sqrt{\frac{(11 + \frac{1}{84} - \frac{1}{11 \cdot 17 \cdot 53 + \frac{2 \cdot 191}{3 \cdot 157}})(112 + \frac{1}{75^2}) \cdot 25 \cdot (112 - \frac{1}{3 \cdot 29 \cdot 64})}{9 \cdot (20 + \frac{1}{2} - \frac{1}{14} + \frac{1}{251} - \frac{1}{8 \cdot (12 \cdot 43 \cdot 227 + 1) - \frac{8}{37}})}}$$

$$= \frac{5}{3} \sqrt{\frac{(11 + \frac{1}{84} - \frac{1}{11 \cdot 17 \cdot 53 + \frac{2 \cdot 191}{3 \cdot 157}})(112^2 - \frac{7 \cdot 19}{2^2 \cdot 3^2 \cdot 25^2 \cdot 29} - \frac{1}{2^6 \cdot 3^3 \cdot 25^2 \cdot 29})}{20 + \frac{1}{2} - \frac{1}{14} + \frac{1}{251} - \frac{1}{8 \cdot (12 \cdot 43 \cdot 227 + 1) - \frac{8}{37}}}}$$

$$= \frac{5}{3} \sqrt{\frac{(11 + \frac{1}{84} - \frac{1}{11 \cdot 17 \cdot 53 + \frac{2 \cdot 191}{3 \cdot 157}})(112^2 - \frac{1}{6 \cdot 17 \cdot 47 + \frac{2}{3}})}{20 + \frac{1}{2} - \frac{1}{14} + \frac{1}{251} - \frac{1}{8 \cdot (12 \cdot 43 \cdot 227 + 1) - \frac{8}{37}}}}$$

$$= \sqrt{137.035999037435 \times 137.035999111818} = 137.035999074627$$

2020 / 2 / 21 - 22

$$c_{au} = \frac{25 \cdot 112}{3} \sqrt{\frac{37}{11 \cdot 12 \cdot 13} - \frac{1}{2 \cdot 17 \cdot 41 \cdot 163 + \frac{47}{6 \cdot 31}}} = 137.035999074628$$

<sup>23</sup> <sub>11</sub> Na <sub>12</sub>	<sup>24,25</sup> <sub>12</sub> Mg <sub>12,13</sub>	<sup>35,37</sup> <sub>17</sub> Cl <sub>18,20</sub>	<sup>55</sup> <sub>25</sub> Mn <sub>30</sub>	<sup>69,71</sup> <sub>31</sub> Ga <sub>38,40</sub>	<sup>74,77,78,82</sup> <sub>34</sub> Se <sub>40,43,44,48</sub>	<sup>84</sup> <sub>37</sub> Rb <sub>47</sub>	<sup>85,87</sup> <sub>37</sub> Rb <sub>48,50</sub>	<sup>93</sup> <sub>41</sub> Nb <sub>52</sub>
<sup>100</sup> <sub>44</sub> Ru <sub>56</sub>	<sup>107,109</sup> <sub>47</sub> Ag <sub>60,62</sub>	<sup>112</sup> <sub>48</sub> Cd <sub>64</sub>	<sup>112,114-120,122,124</sup> <sub>50</sub> Sn <sub>62,64-70,72,74</sub>	<sup>144,147,148,150,154</sup> <sub>62</sub> Sm <sub>82,85,86,88,92</sub>	<sup>157,158</sup> <sub>64</sub> Gd <sub>93,94</sub>			
<sup>163</sup> <sub>66</sub> Dy <sub>97</sub>	<sup>168</sup> <sub>68</sub> Er <sub>100</sub>	<sup>169</sup> <sub>69</sub> Tm <sub>100</sub>	<sup>5,37,11,17</sup> <sub>75</sub> Re <sub>110,112</sub>	<sup>204,206-1613</sup> <sub>82</sub> Pb <sub>122,124-126</sub>	<sup>237</sup> <sub>93</sub> Np <sub>122</sub>	<sup>247</sup> <sub>97</sub> Bk <sub>150</sub>	<sup>257</sup> <sub>100</sub> Fm <sub>157</sub>	<sup>268</sup> <sub>105</sub> Db <sub>163</sub>
<sup>269</sup> <sub>106</sub> Sg <sub>163</sub>	<sup>270</sup> <sub>107</sub> Bh <sub>163</sub>	<sup>285</sup> <sub>112</sub> Cn <sub>173</sub>	<sup>310</sup> <sub>124</sub> Ch <sup>ie</sup> <sub>186</sub>	<sup>334</sup> <sub>132</sub> Ch <sup>ie</sup> <sub>202</sub>	<sup>311<sup>2</sup>,2813</sup> <sub>143</sub> Ch <sup>ie</sup> <sub>220,13-17</sub>	<sup>378</sup> <sub>150</sub> Ch <sup>ie</sup> <sub>228</sub>	<sup>394</sup> <sub>156</sub> Ch <sup>ie</sup> <sub>14-17</sub>	<sup>410</sup> <sub>163</sub> Ch <sup>ie</sup> <sub>13-19</sub>

Note:  $112 \times 5/3 \approx 187 = 11 \times 17$ ,  $112 \times 25/3 \approx 5 \times 11 \times 17$  2020 / 2 / 24

$$c_{au} = \frac{25 \cdot 112}{3} \sqrt{\frac{1}{47} + \frac{1}{40 \cdot 89} - \frac{1}{16 \cdot 9 \cdot (2 \cdot 21 \cdot 31 \cdot 43 + 1) + \frac{3}{4}}} = 137.035999074627$$

<sup>45</sup> <sub>21</sub> Ti <sub>24</sub>	<sup>47</sup> <sub>22</sub> Ti <sub>25</sub>	<sup>55</sup> <sub>25</sub> Mn <sub>30</sub>	<sup>64,66,70</sup> <sub>30</sub> Zn <sub>34,36,40</sub>	<sup>69,71</sup> <sub>31</sub> Ga <sub>38,40</sub>	<sup>72</sup> <sub>32</sub> Ge <sub>40</sub>	<sup>78,80,83,84,86</sup> <sub>36</sub> Kr <sub>42,44,47,48,50</sub>	<sup>89</sup> <sub>39</sub> Y <sub>50</sub>	<sup>90,94,96</sup> <sub>40</sub> Zr <sub>50,54,56</sub>
<sup>92,94-98,100</sup> <sub>42</sub> Mo <sub>50,52-56,58</sub>	<sup>98,99</sup> <sub>43</sub> Tc <sup>*</sup> <sub>55,56</sub>	<sup>107,109</sup> <sub>47</sub> Ag <sub>60,62</sub>	<sup>112</sup> <sub>48</sub> Cd <sub>64</sub>	<sup>144,147,148,150,152</sup> <sub>62</sub> Sm <sub>82,85,86,88,90</sub>	<sup>151,153</sup> <sub>63</sub> Eu <sub>88,90</sub>	<sup>178</sup> <sub>72</sub> Hf <sub>106</sub>		
<sup>185,187</sup> <sub>75</sub> Re <sub>110,112</sub>	<sup>222</sup> <sub>86</sub> Rn <sup>*</sup> <sub>136</sub>	<sup>227</sup> <sub>89</sub> Ac <sup>*</sup> <sub>138</sub>	<sup>237</sup> <sub>93</sub> Np <sup>*</sup> <sub>144</sub>	<sup>244</sup> <sub>94</sub> Pu <sup>*</sup> <sub>150</sub>	<sup>326,328</sup> <sub>129</sub> Ch <sup>ie</sup> <sub>197/199</sub>	<sup>366</sup> <sub>144</sub> Ch <sup>ie</sup> <sub>222</sub>	<sup>9-42</sup> <sub>150</sub> Ch <sup>ie</sup> <sub>228</sub>	

$$c_{au} = \frac{25 \cdot 112}{3} \sqrt{\frac{1}{46} - \frac{1}{55 \cdot 100} + \frac{1}{9 \cdot 25 \cdot 13 \cdot (20 \cdot 293 + 1) - \frac{4}{23}}} = 137.035999074627$$

<sup>50,51</sup> <sub>23</sub> Mn <sub>27,28</sub>	<sup>55</sup> <sub>25</sub> Mn <sub>30</sub>	<sup>89</sup> <sub>39</sub> Y <sub>50</sub>	<sup>99,100</sup> <sub>44</sub> Ru <sub>55,56</sub>	<sup>106,110</sup> <sub>46</sub> Pd <sub>60,64</sub>	<sup>117</sup> <sub>50</sub> Sn <sub>67</sub>	<sup>133</sup> <sub>55</sub> Cs <sub>78</sub>	<sup>169</sup> <sub>69</sub> Tm <sub>100</sub>	<sup>185,187</sup> <sub>75</sub> Re <sub>110,112</sub>	<sup>195</sup> <sub>78</sub> Pd <sub>117</sub>
<sup>5,47,238</sup> <sub>92</sub> U <sup>*</sup> <sub>113,146</sub>	<sup>257</sup> <sub>100</sub> Fm <sup>*</sup> <sub>157</sub>	<sup>285</sup> <sub>112</sub> Cn <sup>*</sup> <sub>173</sub>	<sup>293</sup> <sub>116</sub> Lv <sup>ie</sup> <sub>177</sub>	<sup>294</sup> <sub>117</sub> Ts <sup>ie</sup> <sub>177</sub>	<sup>400</sup> <sub>157</sub> Ch <sup>ie</sup> <sub>243</sub>	<sup>426</sup> <sub>169</sub> Ch <sup>ie</sup> <sub>257</sub>			

**Supplement 7: Comparison of formulas of 1, N, e, 2π, π/2, φ, α, α<sub>c</sub>, c<sub>au</sub> and α<sub>p/c</sub>**

$$1 = 4\gamma_c + \frac{4\gamma_1}{1(1+1)} + \frac{4\gamma_2}{2(2+1)} + \frac{4\gamma_3}{3(3+1)} + \dots$$

$$= |B| \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{|B_{2n}| (\pi/2)^{2n}}{(2n)!} = \sum_{n=1}^{\infty} \frac{|B_{2n}| \pi^{2n}}{(2n)!} = -|B| \frac{3\pi}{2} + \sum_{n=1}^{\infty} \frac{|B_{2n}| (3\pi/2)^{2n}}{(2n)!}$$

$$N \sim -\frac{3}{2}|B| + \sum_{n=1}^N \frac{|B_{2n}| (2\pi)^{2n}}{2(2n)!}$$

$$e = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots$$

$$2\pi = \left(\frac{e}{e^{\gamma_c}}\right)^2 = e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \frac{e^2}{\left(\frac{4}{3}\right)^7} \dots, \quad \frac{\pi}{2} = \left(\frac{e}{e^{\gamma_s}}\right)^2 = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$$

$$2\pi \approx \frac{4 \cdot 157}{100} \approx \frac{9 \cdot 37}{53} \approx \frac{4 \cdot 5 \cdot 71}{15^2 + 1} \approx \dots, \quad \frac{\pi}{2} \approx \frac{157}{25} \approx \frac{9(9+1/4)}{53} \approx \frac{5 \cdot 71}{15^2 + 1} \approx \dots$$

$$\varphi_1 = \frac{\sqrt{5}-1}{2} = 0.618\dots, \quad \varphi_2 = -\frac{\sqrt{5}+1}{2} = -1.618\dots$$

$$\sqrt{\frac{\sqrt{5}+1}{2} + 2} - \frac{\sqrt{5}+1}{2} = \frac{e^{-\frac{2\pi}{5}}}{1 + \frac{e^{-2\pi}}{1 + \frac{e^{-4\pi}}{1 + \frac{e^{-6\pi}}{1 + \dots}}}} \quad (\text{Ramanujan Formula})$$

$$\alpha_1 = \frac{6^2}{7 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{113}{112}\right)^{225}}} \frac{1}{112 + \frac{1}{75^2}} = 1/137.035999037435$$

$$\alpha_2 = \frac{13 \cdot e^2 \frac{e^2}{\left(\frac{2}{1}\right)^3} \frac{e^2}{\left(\frac{3}{2}\right)^5} \dots \frac{e^2}{\left(\frac{9 \cdot 31}{278}\right)^{557}}}{10^2} \frac{1}{112 - \frac{1}{3 \cdot 29 \cdot 64}} = 1/137.0359991118181$$

$$\frac{1}{\alpha_c^2} = \frac{1}{\alpha_1 \alpha_2} = 112 \times \left(168 - \frac{1}{3} + \frac{1}{12 \cdot 47} - \frac{1}{6 \cdot 29 \cdot 53 \cdot 59 - 79/47}\right)$$

$$= 137.035999074627^2 = 18778.865042381$$

$$c_{au} = \frac{1}{\alpha_c} = \frac{1}{\sqrt{\alpha_1 \alpha_2}} = \frac{5}{3} \sqrt{\frac{7 (2\pi)_{112}}{13 (2\pi)_{278}} \left(112^2 - \frac{1}{30^2 \cdot 5} + \frac{1}{60^2 \cdot 15} - \frac{1}{120^2 \cdot 15 \cdot 29}\right)}$$

$$= \frac{25 \cdot 112}{3} \sqrt{\frac{1}{47} + \frac{1}{40 \cdot 89} - \frac{1}{16 \cdot 9 \cdot (2 \cdot 21 \cdot 31 \cdot 43 + 1)} + \frac{3}{4}}$$

$$= \frac{25 \cdot 112}{3} \sqrt{\frac{1}{46} - \frac{1}{55 \cdot 100} + \frac{1}{9 \cdot 25 \cdot 13 \cdot (20 \cdot 293 + 1)} - \frac{4}{23}} = \dots = 137.035999074627$$


$$\frac{1}{\alpha_{p/c}^2} = \frac{1}{\alpha_{p/1} \alpha_{p/2}} = 252.040872632515^2 = 63524.60147736 \quad (\text{Supposed})$$



The relations of the above formulas are sophisticated. In general, some formulas such as 1, N, e and  $2\pi$  have similar form (called the natural group form), some formulas such as  $\phi$ ,  $\alpha$ ,  $\alpha_c$  and  $c_{\text{au}}$  can be divided into rational parts and irrational parts for each which may imply they have the same reasonability. In addition,  $2\pi$ ,  $\pi/2$ ,  $\phi$ ,  $\alpha$ ,  $\alpha_c$ ,  $c_{\text{au}}$  and  $\alpha_{\text{p/c}}$  are all proportional constants, so they should have some similar or the same regularities.

**Supplement 8: Comparison of pictures of elements and  $\phi$**

With the hints of the above formulas, it is not strange that the gold section ( $\phi \approx 0.618$ ) appears in the elements, it should appear in some places with some forms.



**Fig. 11**

Imagine a one-dimension creature lives in the line 0-1, he is familiar with 0-0.618 line space and can reach 0.618-1 line space, if he is enough smart, he may feel there should be an ideal extended line space from 0 to -1.618, but he couldn't reach all or can only get the margin of it. The same situation is suitable for us, we live in the space of elements, we mainly utilize the stable elements and can use some radioactive elements before the 112th element Cn, moreover, there should be a space for ideal extended elements from the 119th to the 170th, a few of which we can synthesize, many of which we can't, but this space should exist. This situation is also suitable for our lives in the earth, the solar system and the universe, or even in the matter, dark matter and dark energy, except that the proportion ratios should be different.

## 22. Discussion and Conclusion

Regarding the fine-structure constant, Richard Feynman said: “is it related to  $\pi$  or perhaps to the base of natural logarithms?”<sup>4</sup> Our answer is that it relate to  $2\pi$ -e formula. He also deduced that the maximum element should be the 137<sup>th</sup> element Fynmanium (Fy) based on the analyses of the electron line velocity of his ideal hydrogen-like atoms. Our answer is that the natural end of the elements is the 112<sup>th</sup> element Copernicium (Cn<sup>\*</sup>), but the elements could have some ideal extensions, and above all, the fine-structure constant does relate to elements.

So, based on the analyses of ideal and real natural maximum element, Chen’s Chirality and Poetry Model of Atomic Nuclei<sup>7</sup> and  $2\pi$ -e formula<sup>6,7,8</sup>, we deduced two series of Chen’s formulas of the fine-structure constant which gave two values  $\alpha_1=1/137.035999037435$  and  $\alpha_2=1/137.035999111818$ . The factors in the formulas are much coincident to nucleon numbers of some nuclides, this means the formulas should be correct (too many coincidences mean too few possibilities to be wrong, or too many coincidences imply science). And we indicate the reason of  $\alpha \approx 1/137.036$  is that it’s almost the equal ratio factor between 112 and 168 (more precisely  $168-1/3$ ) which are the key stable numbers (magic numbers) in Chen’s Chirality and Poetry Model of Atomic Nuclei<sup>7</sup>.

With Chen’s formulas of the fine-structure constant, we predicted the nucleon numbers of all 119<sup>th</sup> to 170<sup>th</sup> ideal extended elements; we theoretically or mathematically calculated the speed of light in atomic units, i.e.,  $c_{au}=1/\alpha_c=1/(\alpha_1\alpha_2)^{1/2}=137.035999074627$ ; we deduced a concise Schrödinger-Chen equation of hydrogen atom which included  $\alpha_1/\alpha_2$  factor; in analogy to  $\alpha$  and its formulas,  $\alpha_p$  (the second fine-structure constant) and its formulas were hypothesized, and the proton charge radius  $r_p$  was supposed to be 0.833027203 fm; in the end we discovered that the approximate rational numbers of  $2\pi$  marvelously and directly related to nuclides. Based on these, a mathematic shell model of elements was established and a picture of elements and ideal extended elements was depicted.

In their relations to nuclides,  $2\pi$  formulas can only be certain approximate rational numbers and  $2\pi$ -e formulas in Chen’s formulas of the fine-structure constant

can only take certain  $k$  values. So we also believe the two values of the fine-structure constant should be rational numbers with definite digits rather than irrational numbers with infinite digits, and actually the fine-structure constant is transformed to nucleon numbers of 136, 137 and 138 in the world of nuclides.

In a recent paper<sup>11</sup>, physicist Nicolas Gisin commented that in 1920s there once was a debate between mathematicians David Hilbert and Luitzen Egbertus Jan Brouwer. Hilbert was promoting formalized mathematics, in which every real number with its infinite series of digits is a completed individual object. On the other side the Luitzen Egbertus Jan Brouwer was defending the view that each point on the line should be represented as a never-ending process that develops in time, a view known as intuitionistic mathematics. Hilbert and his supporters clearly won the debate. In consequence, formalized mathematics has been adopted as the language of physics. In the end of his paper, Nicolas Gisin said: “Physics can be as successful if built on intuitionistic mathematics, even if this breaks its marriage to determinism. Contrary to usual expectations, I bet that the next physical theory will not be even more abstract than quantum field theory, but might well be closer to human experience.”

In this paper we adopted mathematical language like intuitionistic mathematics, but we go ahead even more. The formulas of  $2\pi$ ,  $2\pi-e$  and the fine-structure constant consist of integer factors and relate to nucleon numbers of nuclides, and hence correlate with each others. So in this paper we may use super-intuitionistic mathematics or decoding methodology with features of multi-correlations of integer factors or rational numbers which relate to nucleon numbers of nuclides, and it seems it is the real language in the world of nuclides. As we know an atomic nucleus is a  $N$ -body system and chaos should be its real state, so it seems  $N$ -body chaos returns to integers. In overall, Leopold Kronecker’s famous saying “God made the integers, all else is the work of man” should be correct in the world of nuclides or even in other fields of the real world. It seems an irrational number can only be a rational number to play roles in the real world.

“God is a pure mathematician!” declared British astronomer Sir James Jeans(1877-1946). The physical Universe does seem to be organised around elegant

mathematical relationships<sup>3</sup>. The fine-structure constant may be the most important number in physics. As it is dimensionless, it could be called the proportional ruler of the nature or the bridge of mathematics and physics. And we have successfully given reasonable and precise formulas of it. In some sense, we explain the bridge between mathematics and physics, or we may realize the unification of mathematics and physics. It seems we prove the saying “God is a pure mathematician”. At least, it seems that good mathematics means good physics, and some pure mathematical numbers do have scientific meanings.

## References

1. Wikipedia. The fine-structure constant.
2. R. H. Parker *et al.*, *Science* **360**, 191-195 (2018).
3. P. Davies, *Cosmos* 67 - Feb-Mar 2016.
4. R. P. Feynman, *QED: The Strange Theory of Light and Matter*, Princeton University Press, ISBN 0-691-08388-6, p. 129 (1985).
5. B. Palais, *the Mathematical Intelligencer* 03, 2001, Vol. 23, pp. 7-8.
6. G. Chen and T. Chen, Copyright Registration, Chen’s Periodic Table of Elements and Natural Group Theory, GuoZuoDengZi-2018-L-00472808.
7. G. Chen, T. Chen and T. Chen, Copyright Registration, Chirality and Poetry Model of Atomic Nuclei, GuoZuoDengZi-2018-L-00421847.
8. G. Chen, T. Chen and T. Chen, Copyright Registration, Chen’s Theory of the Fine-structure Constant, GuoZuoDengZi-2018-L-00547467.
9. N. Bezginov *et al.*, *Science* November 06, 2019, Vol. 365, pp. 1007-1012.
10. W. Xiong *et al.*, *Nature* November 06 2019, Vol. 575, pp. 147-150.
11. N. Gisin, *Nature Physics*, <http://www.nature.com/articles/s41567-019-0748-5>.

## Acknowledgements

The main author Dr. Gang Chen studied in the Department of Chemistry at Sichuan University from 1983 to 1987 (Bachelor’s degree), in Institute of Chemistry of the Academy of Sciences of China from 1987-1990 (Master’s degree under supervision of Prof. Rongben Zhang),

in the Hong Kong Polytechnic University from 1999 to 2004 (Ph. D and research assistant under supervision of Prof. Albert Sun Chi Chan) and in Kyoto University from 2004 to 2005 (postdoctoral research under supervision of Prof. Tamio Hayashi).

Leeman Chemical (HK) Co., Ltd. and Jiangshu Leeman Chemical Co., Ltd. gave Dr. Gang Chen a full time employment from May 2005 to July 2017.

Yichang Huifu Silicon Material Co., Ltd. has been giving Dr. Gang Chen a part-time employment since Dec. 2018. Special thanks to Dr. Yuelin Wang of the company for his appreciation and financial support.

Thank Mr. Qiyu Yang who is a famous Chinese traditional poet and an amateur researcher in mathematics. He gave me his book “The Story of Infinity” in 2010, and inspired by this, I started to write my new theory about infinity sets and one of my discoveries was  $2\pi$ -e formula (in 2013).

Thank Ramanujan, the famous Indian mathematician. His story, the picture of his Goddess and the film about him gave me some inspirations in this work.

The main author Dr. Gang Chen is deeply grateful to the above universities, institutions and companies, and to all teachers, classmates, friends and colleagues for their appreciation, encouragement and help to this work.

## Appendix I: Research History

Items	Page	Discover/Create	Revise/Supplement
$2\pi$ -e formula	3	2013/4-12	
Formulas related to $2\pi$ -e formula	4	2013/4-12	
Preliminary applications of $2\pi$ -e formula and its related formulas	6	2013/4-12	
Chen’s Periodic Table of Elements and Natural Group Theory	6	2014-2017/12	
Chirality and Poetry Model of Atomic Nuclei	6	2017/12-2018/3	
Chen’s theory of the fine-structure constant	6	2018/4-6	2018/7-2020/1
Original Inspiration for Formulas of the Fine-structure Constant	6	2018/4/12	
Logical deduction of Chen’s formulas of the fine-structure constant	6 7	2018/4/12-24	
$\alpha_1$ ( $\alpha_{1.7}$ )	6 7	2018/4/12	2018/4/20 (+1/75 <sup>2</sup> )

$\alpha_2 (\alpha_{2-13})$	7	2018/4/24	2018/9/18-20 (280→278 <i>et al.</i> )
Calculation tables and diagrams of $\alpha$	8	2018/4/12-24	2018/9/18-20
$\alpha_{1-(3/2)}$	9	2019/4/25	
$\alpha_{2-(3/2)}$	9	2019/4/25	
$\alpha_c^2$	10	2018/6/8-9, 9/18-19; 2019/4/17-19	
$1/\alpha_c^2$	10	2019/12/14	
112/137≈137/168 <i>et al.</i>	11	2018/4-6	
$^{136,137,138}\text{Fy}_{208,209,210}$	11	2019/12-2020/1	
$^{125,126}\text{Ch}$ , $^{144-149}\text{Ch}$ , $^{153,154}\text{Ch}$ , $^{155,156}\text{Ch}$ , $^{157}\text{Ch}$ , $^{163}\text{Ch}$ , $^{164-168}\text{Ch}$ , $^{169}\text{Ch}$	11	2019/12-2020/1	
$c_{\text{au}}$ formulas	12	2019/12/16	2020/1/5-8
The special 29 and 75 factors	12-15	2019/4/22-24	
$\alpha_1/\alpha_2$ in Schrödinger Equation of Hydrogen Atom	16	2018/4-6	2019/12/13
$\alpha_1/\alpha_2$	16	2019/8/28-29	
The two kinds of general formulas of $\alpha$	17	2019/6/27	2019/7/2-3
Parameters and results of $\alpha_{1-m'}$	18	2019/7/2	
$\alpha_{1-1}$	19	2019/7/2	
$\alpha_{1-2}$	19	2019/6/26	
$\alpha_{1-3}$	19	2019/5/26	2019/6/26
$\alpha_{1-4}$	19	2019/6/27	
$\alpha_{1-5}$	19	2019/6/27	
$\alpha_{1-6}$	19	2019/6/27	
$\alpha_{1-7} (\alpha_1)$	19	2018/4/12	2018/4/20 (+1/75 <sup>2</sup> )
$\alpha_{1-9}$	20	2019/6/28	
$\alpha_{1-11}$	20	2019/6/29	
$\alpha_{1-13}$	20	2019/6/29	
$\alpha_{1-16}$	20	2019/6/29-30	
$\alpha_{1-17}$	20	2019/6/30	2019/10/29
$\alpha_{1-19}$	20	2019/7/1	
$\alpha_{1-20}$	20	2018/4-6	2019/6/26
$\alpha_{1-22}$	21	2019/5/25	2019/12/12
$\alpha_{1-23}$	21	2019/7/4	
$\alpha_{1-25}$	21	2019/7/4	
$\alpha_{1-27}$	21	2019/5/25-26	
$\alpha_{1-29}$	21	2019/5/25	
$\alpha_{1-31}$	21	2019/7/4	
$\alpha_{1-32}$	21	2019/7/5	
$\alpha_{1-33}$	22	2019/7/5	
$\alpha_{1-34}$	22	2019/5/24	

$\alpha_{1-36}$	22	2019/5/24	
$\alpha_{1-43}$	22	2019/12/15	
$\alpha_{1-50}$	22	2018/4-6	2019/6/26
$\alpha_{1-59}$	22	2019/5/25	
$\alpha_{1-81}$	22	2019/5/25-26	
$\alpha_{1-96}$	23	2019/5/25-26	2019/12/29
$\alpha_{1-103}$	23	2019/12/15	
$\alpha_{1-133}$	23	2019/5/26	2019/12/29
$\alpha_{1-140}$	23	2019/5/26	
$\alpha_{1-155}$	23	2019/12/15	
$\alpha_{1-170}$	23	2019/7/2	2019/7/9
<hr/>			
Parameters and results of $\alpha_{2-m'}$	24	2019/7/3	
$\alpha_{2-1}$	25	2018/4-6	2019/5/26-27
$\alpha_{2-4}$	25	2019/5/28	
$\alpha_{2-5}$	25	2019/6/22	
$\alpha_{2-6}$	25	2019/6/23	
$\alpha_{2-7}$	25	2019/5/24	
$\alpha_{2-9}$	25	2019/6/21	
$\alpha_{2-10}$	26	2019/5/28	
$\alpha_{2-11}$	26	2019/6/22	
$\alpha_{2-13} (\alpha_2)$	26	2018/4/24	2018/9/18-20 (280→278 <i>et al.</i> )
$\alpha_{2-15}$	26	2019/6/19	
$\alpha_{2-17}$	26	2019/6/24	
$\alpha_{2-18}$	26	2019/6/24	
$\alpha_{2-19}$	27	2019/6/24	
$\alpha_{2-23}$	27	2019/6/23	
$\alpha_{2-24}$	27	2019/6/23	
$\alpha_{2-25}$	27	2019/6/24	
$\alpha_{2-27}$	27	2019/6/21	
$\alpha_{2-29}$	27	2019/6/20	
$\alpha_{2-31}$	28	2019/7/6	
$\alpha_{2-32}$	28	2019/7/6	
$\alpha_{2-33}$	28	2019/7/6	
$\alpha_{2-36}$	28	2019/6/25	
$\alpha_{2-37}$	28	2019/7/7	
$\alpha_{2-38}$	28	2019/7/7	
$\alpha_{2-125}$	29	2019/5/25	2019/12/20
$\alpha_{2-253}$	29	2019/7/3	2019/7/9
$\alpha_{2-269}$	29	2019/7/7	2019/12/19
<hr/>			
$a_0/r_e, r_e; a_0/r_p, r_p$	30	2019/12/19-23	
<hr/>			
$\alpha_{p/1}$	30	2020/1/2	
<hr/>			

$\alpha_{p/2}$	30 31	2020/1/2-3	
${}_{164}\text{Ch}_{252}$	31	2020/1/3	
Direct relationships of $2\pi$ with nuclides	31	2020/1/8-10	
Some correlations of formulas of $2\pi$ and $\alpha$	32 33	2020/1/11-13	
Chen's Mathematic Shell Model of Nuclides	33	2020/1/12-13	
${}_{130,131}\text{Ch}_{200,201}$	20 22 26 27	2020/1/26-28	
${}_{119,120,121}\text{Ch}_{179,180,181}$	21 28 31	2020/1/28-29	2020/2/5 (add 121)
${}_{128,129}\text{Ch}_{198,197/199}$	23 26 31	2020/1/28-29	2020/1/31
${}_{139}\text{Ch}_{209}$	21 26 28	2020/1/29	
${}_{132,133}\text{Ch}_{202,203}$	21 23	2020/1/31	
${}_{169}\text{Ch}_{257}$	11 22 27	2020/1/29-30	
${}_{157}\text{Ch}_{243}$	11 20 21 22 31	2019/1/8	
${}_{158}\text{Ch}_{243}$	11 20 22 25 30 31	2019/1/31	
${}_{169}\text{Ch}_{257}$	11 22 26 27	2020/1/29-30	
${}_{134,135}\text{Ch}_{206,205}$	20	2020/1/31	
${}_{127}\text{Ch}_{191,193}$	27	2020/1/31	2020/2/1 (add 191)
${}_{150,151,152}\text{Fy}_{228,229,230}$	19 28	2020/2/2	
${}_{170}\text{Ch}_{250}$	23 26	2020/2/3	
Chen's Picture of Elements and Ideal Extended Elements	41	2018/1-3 2020/2/2-5	2020/2/12, 16, 17, 19, 22-24
$2\pi=62831853/10^7$	37	2020/2/11-12	
${}_{124}\text{Ch}_{186}$	37	2020/2/12	
Table 8	37	2020/2/16-17	
${}_{143}\text{Ch}_{220,221}$	36	2020/2/17	
Supplement 4	38	2020/2/18-21	
${}_{122}\text{Ch}_{182}$	22	2020/2/19	
Supplement 5: $\alpha_{1-9/11}$ $\alpha_{2-20/25}$	38	2020/2/21	
${}_{127}\text{Ch}_{192}$ ${}_{167}\text{Ch}_{251}$	38	2020/2/21	
Other Formulas of $c_{\text{au}}$	39	2020/2/21-22, 24-25	
${}_{140,141}\text{Ch}_{212,215}$	39	2020/2/22	
${}_{123}\text{Ch}_{183,185}$	36	2020/2/23	
${}_{159/161,160}\text{Ch}_{245,246}$	36	2020/2/23	
${}_{165}\text{Ch}_{255}$	36	2020/2/23	
${}_{162}\text{Ch}_{246}$	36	2020/2/23	
Supplement 7	40	2020/2/25-26	
Supplement 8	41	2020/2/26	
Preparing this paper	1-46	2019/12/1-2020/2/26	