

MATTER THEORY ON EM FIELD

WU SHENG-PING

ABSTRACT. This article try to unified the four basic forces by Maxwell equations, the only experimental theory. Self-consistent Maxwell equations with the e-current coming from matter current is proposed, and is solved to electrons and the structures of particles and atomic nucleus. The static properties and decay are reasoned, all meet experimental data. The equation of general relativity sheerly with electromagnetic field is discussed as the base of this theory. In the end the conformation elementarily between this theory and QED and weak theory is discussed.

CONTENTS

1. Bound Dimensions	1
2. Inner Field of Electron	2
3. General Electromagnetic Field	3
4. Solution of Electron	3
5. Electrons	5
6. System and TSS of Electrons	6
7. Muon	8
8. Pion	8
9. Pion Neutral	9
10. Proton	9
11. Neutron	9
12. Atomic Nucleus	10
12.1. Decay Energy	11
12.2. β -stable and Neutron Hide	11
13. Basic Results for Interaction	12
14. Grand Unification	12
15. Conclusion	13
References	13

1. BOUND DIMENSIONS

A rebuilding of units and physical dimensions is needed. Time s is fundamental.

We can define:

The unit of time: s (second)

The unit of length: cs (c is the velocity of light)

Date: Jun 16, 2021.

Key words and phrases. Maxwell equations; Decay; Electron function; Recursive re-substitution.

The unit of energy: \hbar/s (\hbar is Plank constant)

The unit dielectric constant ϵ is

$$[\epsilon] = \frac{[Q]^2}{[E][L]} = \frac{[Q]^2}{\hbar c}$$

The unit of magnetic permeability μ is

$$[\mu] = \frac{[E][T]^2}{[Q]^2[L]} = \frac{\hbar}{c[Q]^2}$$

The unit of Q (charge) is defined as

$$c[\epsilon] = c[\mu] = 1$$

then

$$[Q] = \sqrt{\hbar}$$

$$\sqrt{\hbar} = (1.0546 \times 10^{-34})^{1/2} C$$

C is charge's SI unit Coulomb.

For convenience, new base units by unit-free constants are defined,

$$c = 1, \hbar = 1, [Q] = \sqrt{\hbar} = [1]$$

then the units are reduced.

Define

$$\text{UnitiveElectricalCharge} : \sigma = \sqrt{\hbar}$$

$$\sigma = 1.027 \times 10^{-17} C \approx 64e$$

$$e_{/\sigma} = e/\sigma = 1.5602 \times 10^{-2} \approx 1/64$$

It's defined that

$$\beta := m/e = 1, \quad m := |k_e| \approx m_e$$

Then all units are power σ^n . This unit system is called *bound dimension* or *bound unit*. We always take the definition latter in this article

$$\beta = 1, \quad \sigma = 1$$

We always take them as a standard unit.

Define a *measure* σ/z , $[z] = 1$:

$$\sigma = z \quad \sigma/z = 1$$

2. INNER FIELD OF ELECTRON

Try the self-consistent Maxwell equation for the inner electromagnetic (EM) field of electrons

$$(2.1) \quad \partial^l \partial_l A^\nu = i A_\mu^* \partial_\nu A^\mu / 2 + cc. = \mu J^\nu \quad m = 1$$

$$\partial^\nu \cdot A_\nu = 0$$

with definition

$$[W := \langle A | A_\mu^* \partial_\nu A^\mu \rangle] = \sigma^4, \quad [A] = \sigma^2$$

$$(A^i) := (V, \mathbf{A}), (A_i) := (V, -\mathbf{A})$$

$$(J^i) = (\rho, \mathbf{J}), (J_i) = (\rho, -\mathbf{J})$$

$$\partial := (\partial_i) := (\partial_t, \partial_{x_1}, \partial_{x_2}, \partial_{x_3})$$

$$\partial' := (\partial^i) := (\partial_t, -\partial_{x_1}, -\partial_{x_2}, -\partial_{x_3})$$

$$g_{ij} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}$$

This equation has the symmetry CPT that transfers electron to its negative.

3. GENERAL ELECTROMAGNETIC FIELD

We find

$$(x', t') := (x, t - r) \\ \partial_x^2 - \partial_t^2 = \partial_{x'}^2 =: \nabla'^2$$

The following is the energy of a piece of field A :

$$(3.1) \quad \varepsilon := \frac{1}{2} (\langle E, D \rangle + \langle H, B \rangle)$$

The time-variant part is neglected, as a convention for energy calculation. If the field has Fourier transformation then this *electromagnetic (EM) energy* becomes

$$(3.2) \quad \varepsilon = \frac{1}{2} \langle A_\nu | \partial_t^2 - \nabla^2 | A^\nu \rangle$$

under Lorentz gauge.

4. SOLUTION OF ELECTRON

The solution by *recursive re-substitution* (RRS) for the two sides of the equation is proposed. For the equation

$$\hat{P}'B = \hat{P}B$$

Its algorithm is that (It's approximate, the exact solution needs a rate on the start state in the re-substitution for the normalization condition)

$$\hat{P}' \left(\sum_{k \leq n} B_k + B_{n+1} \right) = \hat{P} \sum_{k \leq n} B_k$$

A function is initially set and is corrected by RRS of the equation 2.1. Here is the start state

$$A_i = A_r e^{-ikt}, \partial_\mu \partial^\mu A_i = 0$$

The fields' correction A_n with n degrees of A_i is called the n degrees correction.

Firstly

$$\nabla^2 \phi = -k^2 \phi$$

is solved. Exactly, it's solved in spherical coordinate


$$-k^2 = \nabla^2 = \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{r^2 \sin^2 \theta} (\partial_\varphi)^2$$

Its solution is

$$\Omega_k := \Omega_{klm} = k j_l(kr) Y_{lm}(\theta, \varphi) e^{-ikt} \\ \phi_k := k \phi_{klm} e^{-ikt} := k h_l(kr) Y_{lm}(\theta, \varphi) e^{-ikt} \\ \omega_k := k j_1(kr) Y_{11}(\theta, \varphi) e^{-ikt}$$

After normalization it's in effect

$$h = \frac{e^{\pm ir}}{r}, \quad j = \frac{\cos r, \sin r}{r}$$

FIGURE 1. The function of j_1

In fact, it has to be re-defined in the *smooth truncation* that keeps Fourier form

$$\Omega^\beta(x) = \Omega(x)e^{-\beta r}, \quad \beta \rightarrow 0^+$$

or equivalently

$$r \rightarrow |x| + |y| + |z|$$

The similar is for ϕ .

We use the following definition

$$\omega_k := n_1(kr)ke^{-ikt} = \sum_{\hat{\mathbf{k}}} \mathbf{F}(\hat{\mathbf{k}})e^{i\mathbf{k}x - ikt}, \quad \hat{\mathbf{k}} = \frac{\mathbf{k}}{|\mathbf{k}|}, \quad |\mathbf{k}| = k$$

There are calculations:

$$(\partial_t^2 - \nabla^2)u = -\nabla'^2 u = \delta(x')\delta(t') = \delta(x)\delta(t),$$

$$u := \frac{\delta(t-r)}{4\pi r} = \frac{\delta(t')}{4\pi r'},$$

$$\nabla^2 = \sum_{\mathbf{k}} -\mathbf{k}^2 e^{i\mathbf{k}x}$$

$$(4.1) \quad (k\Omega_k(x) * |k\Omega_k(-x)) = \sigma k^2 \frac{\delta^3(kx)}{\sigma^3 \delta^3(0)} \quad \sigma = 1$$

Its smooth truncation is considered.

$$\delta(0) := \delta(\sigma t)|_{t=0}$$

$$\int_I dx (\Omega(x) * \Omega(x))^n = \left(\int_I dx \Omega(x) * \Omega(x) \right)^n$$

In the frequencies of $\Omega(x) \cdot \Omega(x)$ the zero frequency is with the highest degrees of infinity. This principle is quite extensive in fact.

To avoid singularity it's calculated like

$$(e^{ir} * \nabla^2 |e^{ir}) = (\nabla_\nu e^{ir} * |\nabla_\nu e^{ir})$$

in Fourier space.

It's found that

$$\begin{aligned}
& \langle e^{-\beta r} \frac{e^{i(r+\varphi)}}{r} \sin \theta | * | e^{-\beta r} \frac{e^{i(r+\varphi)}}{r} \sin \theta \rangle \\
&= - \langle e^{-\beta r} \frac{e^{i(r-\varphi)}}{r} \sin \theta | * | e^{-\beta r} \frac{e^{i(r-\varphi)}}{r} \sin \theta \rangle \\
&= - \langle e^{-\beta r} \frac{e^{i(-r+\varphi)}}{r} \sin \theta | * | e^{-\beta r} \frac{e^{i(-r+\varphi)}}{r} \sin \theta \rangle \\
& \langle e^{-\beta r} \frac{\cos r e^{i\varphi}}{r} \sin \theta | * | e^{-\beta r} \frac{\cos r e^{i\varphi}}{r} \sin \theta \rangle \\
&= - \langle e^{-\beta r} \frac{\cos r e^{-i\varphi}}{r} \sin \theta | * | e^{-\beta r} \frac{\cos r e^{-i\varphi}}{r} \sin \theta \rangle
\end{aligned}$$

These also can be proved in Fourier space. We can find that the symmetry at the singularity is different, the convolution can't commute.

There is a problem of complex function

$$F(t) := h(t)e^{it} = \begin{cases} e^{it} & t > 0 \\ 0e^{it} & t < 0 \end{cases}$$

$$F^*(t) = F(-t)$$

We best rely on real function.

5. ELECTRONS

It's the start electron function for the RRS of the equation 2.1:

$$A_i^\nu := i\lambda\partial^\nu\omega_k(x,t)/\sqrt{2}, \quad \lambda \approx 1$$

Some states are defined as the core of the electron, which's the start function $A_i(x,t)$ for the RRS of the equation 2.1 to get the whole electron function of field A : e

$$\begin{aligned}
e_r^+ &: \omega_m(\varphi, t), & e_l^- &: \omega_m(-\varphi, -t) \\
e_l^+ &: \omega_m(-\varphi, t), & e_r^- &: \omega_m(\varphi, -t) \\
e_r^+ &\rightarrow e_l^+ : (x, y, z) \rightarrow (x, -y, -z)
\end{aligned}$$

The electron function is normalized with charge as

$$e = \langle A^\mu | i\partial_t | A_\mu \rangle / 2 + cc.$$

The MDM (Magnetic Dipole Moment) of electron is calculated as the second degree proximation

$$\begin{aligned}
\mu_z &= \langle A_{i\nu} | -i\partial_\varphi | A_i^\nu \rangle \cdot \hat{z} / 4 + cc. \\
&= \frac{e}{2m}
\end{aligned}$$

The spin is

$$S_z = \mu_z k_e / e = \hat{k}_e / 2$$

The correction in RRS of the equation 2.1 is calculated as

$$A - A_i = \frac{(A_i^* \cdot i\partial A_i / 2 + cc.) *_4 u}{1 - i\partial(A_i - A_i^*) / 2 *_4 u}$$

The function of e_r^+ is decoupled with e_l^+

$$\langle (e_r^+)^{\nu} | i\partial_t | (e_l^+)^{\nu} \rangle / 2 + \langle (e_l^+)^{\nu} | i\partial_t | (e_r^+)^{\nu} \rangle / 2 + cc. = 0$$

The following is the increment of the energy ε on the coupling of e_r^+, e_r^- , mainly between A_1 and A_3

$$\begin{aligned}\varepsilon_e &= \langle (e_r^+)^{\nu} | i\partial_t | (e_r^-)_{\nu} \rangle / 2 + \langle (e_r^-)^{\nu} | i\partial_t | (e_r^+)_{\nu} \rangle / 2 + cc. \quad m = 1 \\ &\approx -2e^3_{/\sigma} m = -\frac{1}{1.66 \times 10^{-16} s}\end{aligned}$$

Their smooth truncation are considered. Its algorithm is

$$-\frac{2}{2} \cdot \frac{2 \cdot 2^2}{2^2}$$

The following is the increment of the energy ε on the coupling of e_r^+, e_l^- , mainly between A_1 and A_7 .

$$\begin{aligned}\varepsilon_x &= \langle (e_r^+)^{\nu} | i\partial_t | (e_l^-)_{\nu} \rangle / 2 + \langle (e_l^-)^{\nu} | i\partial_t | (e_r^+)_{\nu} \rangle / 2 + cc. \quad m = 1 \\ &\approx -\frac{1}{2} e^7_{/\sigma} m = -\frac{1}{1.145 \times 10^{-8} s}\end{aligned}$$

Its algorithm is

$$-\frac{2}{.4 \cdot 4} \cdot \frac{2 \cdot 2^7}{2^6}$$

and

$$\begin{aligned}&\langle \cos^5(r - \phi) | \cos(r + \phi) \rangle, \quad \phi = t - \varphi \\ &= \langle \cos(2r) \cos^2 \phi (\cos^4 r \cos^4 \phi - \sin^4 r \sin^4 \phi) \rangle \\ &= -\frac{1}{4} \cdot \frac{1}{4} \cdot N\end{aligned}$$

These calculations of the integral product are applied:

$$(5.1) \quad \phi = Y_{11} \frac{e^{ir}}{r}, \quad -i\nabla\phi = Y_{11} \frac{e^{ir}}{r} \hat{r}$$

Normalization is considered.

6. SYSTEM AND TSS OF ELECTRONS

The movement of electron makes an EM field denoted by A , the unit of which is verified by interactions:

$$A := f *_3 \sum_i e_i = N \sum_X f(X, T) \delta(x - X, t - T) *_4 \sum_i e_i(x, t)$$

The following are naked stable particles:

<i>particle</i>	<i>electron</i>	<i>photon</i>	<i>neutino</i>
<i>notation</i>	e_r^+	γ_r	ν_r
<i>structure</i>	e_r^+	$(e_r^+ + e_r^-)$	$(e_r^+ + e_l^-)$

The following is the system of particle x with the initial state

$$A_0 := e_x \sum_c e_c,$$

There are two *base transforms*

$$\begin{aligned}e_c &:= *e, \pm e, \quad d := (r, l) \\ \Phi * e_x * *e &:= (\Phi(x, t) * e_x)^*(x, t) * e(x, t) \\ \Phi * e_x * -e &:= \Phi(-x, -t) * e_x(-x, -t) * e(-x, -t) \\ [f\Phi] * *e &:= (f\Phi)^*(x, t) * e(x, t)\end{aligned}$$

$$[f\Phi] * -e := (f\Phi)(-x, -t) * e(-x, -t)$$

The second one transfers an incoming wave to an outgoing wave, with which we can shift a particle from the left to the right in the reaction formula, with its meaning (decay energy, charge) unchanged.

e_x, Φ meet normalization (including dimensions):

$$(6.1) \quad \langle e^{-\beta r} \cdot \Phi * e_x | e^{-\beta r} \cdot \Phi * e_x \rangle = 1, \quad \langle e^{-\beta r} \cdot e_x | e^{-\beta r} \cdot e_x \rangle = 1 \quad \beta \rightarrow 0^+$$

$$\sigma[\langle \Phi * e_x * e | \Phi * e_x * e \rangle] = 1$$

The normalization is invariant under the two base transforms.

Its dense of e-current of particle x is

$$J = (e^{-\beta r} \cdot \Phi * e_{xc} * e_c | e^{-\beta r} \cdot [i\partial_t \Phi] * e_{xc} * i\partial_t e_c) / 2 + cc. \quad k_\Phi = 1 \quad \beta \rightarrow 0^+$$

The static or motive results are the same. This is compared to the latter calculation of charge conservation and the 2-degree correction of electron.

The momentum dense of system is

$$p = (e^{-\beta r} \cdot \Phi * e_{xc} * e_c | i\partial_t | e^{-\beta r} \cdot [i\partial_t \Phi] * e_{xc} * e_c) / 2 + cc. \quad k_\Phi = 1 \quad \beta \rightarrow 0^+$$

Try the action of isolated system for static mass

$$I = \langle e_{xc} * (e_c)^\mu | \frac{1}{2} \partial_\nu \partial^\nu | e_{xc} * (e_c)_\mu \rangle + \langle e_{xc} * (e_c)^\mu | i\partial_t | e_{xc} * i\partial_t (e_c)_\mu \rangle + cc.$$

and with the zero-border condition it's found that

$$\delta I = 0$$

e_{xc} here is without truncation. As its singularity is cut-off the function is quite normal, and the limit of β is go first of δ or the both can commute (that's a fact).

$$(6.2) \quad -\frac{1}{2} \partial_\nu \partial^\nu (e_{xc} * (e_c)_\mu) = i\partial_t (e_{xc} * i\partial_t (e_c)_\mu)$$

The initial A_0 is found that

$$(\partial_t^2 - \nabla^2) e_x = 0$$

Because of the normalization (6.1), the singularity is neglected.

$$e_x := k_x \Omega_{k_x}(x) \quad \sigma = 1$$

As this state A_0 is considered as a (Transient) Steady State (TSS), e_x here is a Light State (LS).

With the charge conservation, after all electrons off-couple and become isolated electrons,

$$\begin{aligned} & \langle e_x * \sum_c (e_c)^\mu | e_x * i\partial_t \sum_c (e_c)_\mu \rangle / 2 + cc. \\ & = \sum_\nu \langle \phi_{k_\nu} * (e_c)^\mu | \phi_{k_\nu} * i\partial_t (e_c)_\mu \rangle / 2 + cc. \end{aligned}$$

then it's found that

$$|k_x| = \left| \frac{n_x}{Q_x/e} \right| \quad \sigma = 1, \quad n_x := \langle \sum_c e_{xc} | \sum_c e_{xc} \rangle$$

Smooth truncation is used. It's right at least for proton and muon.

About the conditions of the parities $P(d)$, I think the coupling of d on an electron is the results of coupling of MDM or EDM. In some case if an E-field is put on to uncouple them, the gravitational mass of them would become negative (see the latter).




FIGURE 2. neutrino radiation

7. MUON

$$\mu^- : e_{\mu l} * (e_l^- - e_l^+ - e_r^-), \quad e_{\mu} = e_x(k_x = -m_{\mu})$$

μ is approximately with mass $3m/e/\sigma = 3 \times 64m$ [3.2][1] (The data in bracket is experimental by the referenced lab), spin S_e (electron spin), MDM $\mu_B m/k_{\mu}$.

The main channel of decay is

$$\mu^- \rightarrow e_l^- - \nu_l, \quad e_l^- \rightarrow -e_r^+ + \nu_r$$

$$e_{\mu} * e_l^- - e_{\mu} * \nu_l \rightarrow e_{\mu} * e_l^- - \Phi * \nu_l$$

The isolated system conserves momentum in mass-center frame.

Its main life is

$$\begin{aligned} \varepsilon_{\mu} &:= \langle (e_{\mu}^*(-x, -t) * e_{\mu}) \cdot ((e_l^+)^{\mu*}(-x, -t) * i\partial_t(e_r^-)_{\mu}) \rangle |_{t=0} + cc. \quad m = 1 \\ &= -\frac{m\varepsilon_x}{k_{\mu}} = -\frac{1}{2.2015 \times 10^{-6}s} \quad [2.1970 \times 10^{-6}s] \end{aligned}$$

Smooth truncation is applied. The charge conservation on neutrino is applied.

8. PION

The initial of pion perhaps is

$$\pi^- : e_{\pi} * (e_l^+ - e_l^+ + *e_r^-) = e_{\pi} * (e_l^+ - e_l^+) + e_{\pi} * *e_r^-$$

It's approximately with EM energy 3σ [4.2][1]. spin S_e , MDM $\mu_B m/k_{\pi^-}$.

Decay Channels:

$$\pi^- \rightarrow -e_l^+ + \nu_l, \quad e_l^+ \rightarrow -e_r^- + \nu_l$$

The mean life approximately is

$$-\varepsilon_x/2 = \frac{1}{2.3 \times 10^{-8}s} \quad [(2.603 \times 10^{-8}s)[1]]$$

The precise result is calculated with successive decays.

9. PION NEUTRAL

The initial of pion neutral is perhaps like an atom

$$\pi^0 : (e_r^+ + e_l^+, e_r^- + e_l^-)$$

It's the main decay mode as

$$\pi^0 \rightarrow \gamma_r + \gamma_l$$

The mean life is

$$-2\varepsilon_e = \frac{1}{8.3 \times 10^{-17} s} \quad [8.4 \times 10^{-17} s][1]$$

10. PROTON

The initial of proton may be like

$$p^+ : e_{pr} * (4e_r^- - 2e_l^+ + - * 3e_r^+), \quad e_p = e_x (k_x = m_p)$$

The mass is $29 \times 64m$ [29][1] that's very close to the real mass. The MDM is calculated as $3\mu_N$ (mainly by track), spin is S_e . The proton thus designed is eternal.

11. NEUTRON

Neutron is the atom of a proton and a muon

$$n = (p^+, \mu^-)$$

The muon take the first track, with the decay process

$$\Phi * \mu^- = \Phi * e_\mu * (e_l^- - e_l^+ - e_r^-) \rightarrow \Phi * e_\mu * e_l^- - \nu_l$$

Make some normalization to fit 6.1. Calculate the variation of the action of the open system, the energy of system subtracting the affection,

$$(11.1) \quad i\partial_t \Phi + \frac{1}{2} \nabla^2 \Phi = -\frac{\alpha}{r} \Phi \quad m_\mu = 1$$

$$\alpha = \frac{e^2}{4\pi\epsilon\hbar c} \approx 1/137$$

One of the terms of EM energy are neglected, and its domain is cut off O . It's resolved to

$$\Phi = N e^{-r/r_0} e^{-iE_1 t}$$

$$E_1 = E_B \frac{\sigma^2}{m_\mu^2} \cdot \frac{m_\mu}{m}$$

$$E_B = -13.6 eV$$

It's approximately the decay life of muon in the track that

$$\varepsilon_n = -\frac{\sigma^2}{m_\mu^2} \cdot \frac{E_B}{m} e^3 / \sigma \varepsilon_x = -\frac{1}{1019 s}$$

12. ATOMIC NUCLEUS

We can find the equation for the fields of Z' ones of proton: Φ_i , and the fields of n ones of muon: $\Phi_{j:j>Z'} = \phi_i$:

$$\Phi = \sum_i \Phi_i, \quad \phi = \sum_i \phi_i$$

$$\varphi_\nu := \Phi_\nu * (p/\mu)_l, \quad \varphi^\nu := \Phi_\nu * (p/\mu)^l$$

We have

$$I = \sum_\mu \langle \varphi^\mu | \frac{1}{2} \partial^\nu \partial_\nu | \varphi_\mu \rangle_4 / 2 - \sum_\nu \langle \varphi^\nu | \frac{1}{2} \partial^\nu \partial_\nu | \sum_{i \neq \nu} \varphi_i \rangle_4 / 2 + cc.$$

The third-level waves must catch independent tracks, or involve in strong effects. Strong effects (interactions) are omitted. Make the variation of this action on Φ_i, ϕ_i

$$\delta I = 0$$

to find

$$\begin{aligned} \frac{1}{2} \partial_t^2 \Phi + ik_p \partial_t \Phi + \frac{1}{2} \nabla^2 \Phi &= (Z' + 2) \frac{\alpha \sigma^2}{r} * \Phi - n \frac{\alpha \sigma^4}{r} * \phi \\ \frac{1}{2} \partial_t^2 \phi + ik_\mu \partial_t \phi + \frac{1}{2} \nabla^2 \phi &= -Z' \frac{\alpha \sigma^2}{r} * \Phi + (n - 2) \frac{\alpha \sigma^4}{r} * \phi \end{aligned}$$

If E is known (which does exist undoubtedly and invariantly) the second equation is rendered to

$$\frac{1}{2} \partial_t^2 \phi + ik_p \partial_t \phi + \frac{1}{2} \nabla^2 \phi = -Z' \frac{\alpha \sigma^2}{r} * \Phi + (n - 2) \frac{\alpha \sigma^4}{r} * \phi$$

Make

$$\begin{aligned} \zeta &= \Phi + \phi \eta \\ (Z' + 2) - \eta Z' &= -n/\eta + (n - 2) =: N \\ \eta &= \frac{(Z' - n + 4) \pm \sqrt{(Z' - n + 4)^2 + 4Z'n}}{2Z'} \end{aligned}$$

then

$$\begin{aligned} -(E^2/2 + Ek_p) \nabla^2 \zeta + \frac{1}{2} \nabla^4 \zeta + 4\pi \alpha \sigma^2 N \zeta &= 0 \\ \zeta e^{-ikt} &= \sum C_{lm} \Omega_{klm} \end{aligned}$$

So that the first approximation from $\nabla^2 (= -k^2) = -k_p^2$:

$$E = -k_p - \sqrt{8\pi \alpha \sigma^2 N}$$

$$(12.1) \quad N(Z', n) = \frac{1}{2} (Z' + n - \sqrt{(Z' + n)^2 + 8(Z' - n) + 16})$$

$$\approx -2\chi \quad \chi := \frac{Z' - n}{Z' + n}$$

$$E \approx -k_p - E_g, \quad E_g =: 10 \text{ MeV}, \quad \chi = 1/3$$

These state is TSS and LS obviously: $|k| = |E|$, hence

$$E^4 + E^3 + 8\pi \alpha \sigma^2 N = 0 \quad k_p = 1$$

Every single muon or proton has third-level wave

$$\Phi_i = \sum C'_{lm} \Omega_{Elm}, \quad \phi_i = \sum C''_{lm} \Omega_{Elm}$$


12.1. **Decay Energy.** Make some normalization to fit 6.1. The life-involved energy of proton or muon under the solved wave Φ_i is

$$\begin{aligned}\varepsilon &= \langle \Phi_i * (p, \mu)^l | i\partial_t | \Phi_i * (p, \mu)_l \rangle / 2 + cc. \quad m = 1 \\ &= \left(1 - \frac{2}{|E|}, \quad 1/3 + \frac{4}{|E|}\right) e^{11/\sigma}\end{aligned}$$

Smooth truncation is on Φ_ν . The measure change the calculation a few.

$$\begin{aligned}\Delta_E \varepsilon &= E_\Delta(2, -4) \quad m = 1 \\ E_\Delta &\approx \frac{1}{4s}, \quad \Delta E = E_g\end{aligned}$$

The decay



is with zero energy decrease unless the neglected weak crossing is considered.

12.2. **β -stable and Neutron Hide.** In the solutions when a proton combine with a muon the both have the same wave function:

$$\Phi_\mu = \phi_\mu$$

then this terms will quit from the interactional terms of the previous equations, which change to

$$(Z'_x, n_x) \rightarrow (Z'_x - 1, n_x - 1)$$

This will change the *flag energy* ΔE . If a β -decay can't happen then a dismiss of this *neutron hide* will help. Out of the hidden nucleons ($\geq 2z$), the ratio $2 : 1 = Z' : n$ between protons and muons causes the most stable state. So that if

$$-N(Z' + \max(z) + z, n + \max(z) + z) \leq -N(Z' - z, n)$$

$$(12.2) \quad \frac{Z + \kappa - h}{3(Z + \kappa) + 2\max(z) + 2z} \leq \frac{Z + \kappa - h - z}{3(Z + \kappa) - z}, \quad 8h := 3\kappa^2 + 8\kappa - 16$$

then β -decay wouldn't happen for the reversed process can happen. The sign of the coefficient (minus here) of the gross decay energy is also noticed.

It's found a critic point

$$\kappa \approx -8 : \quad h \approx 14$$

and

$$Z \approx 29$$

hence N of $\max(z) = 0$ is calculated near to $-2/3$ through parameter κ . By the following result 12.3, conditions $\chi = 1/3$ and $\max(z) = 0$ are specific for this critical point, for the other Z or κ , the both of which are incompatible.

The condition 12.2 is solved to

$$2(z + \max(z))z < 2\max(z)(Z + \kappa) - (2\max(z) + z)h$$

hence as

$$(12.3) \quad \frac{1}{4} < \chi \leq \frac{1}{3}, \quad z \leq \max(z) = ((Z + \kappa) - 1.5h)/2$$

The reaction is very weak.

It's the *Average Binding Mass Per Nucleon* according to charge number that

$$\begin{aligned}\varepsilon_M(Z) &= X \cdot \sigma \Delta_0^Z \sqrt{-8\pi\alpha N} \\ X &= (Z \leq 29) + (Z > 29) \frac{2.0}{2.5 - 10.5/(Z - 8)}\end{aligned}$$




FIGURE 3. Average Binding Mass: $\varepsilon_M(0) - \varepsilon_M(Z)$

The EM energy is gravitational mass by the discussions in the section 14, for proton

$$m_p = \langle \varphi_\nu | \frac{1}{2} \partial_\mu \partial^\mu | \varphi_\nu \rangle \approx k_p$$

Smooth truncation is on Φ_ν . By the motion Φ_ν to find its mechanical energy

$$M_p/m_p = -E \quad m_p = 1$$

13. BASIC RESULTS FOR INTERACTION

For decay

$$(13.1) \quad W(t) = \Gamma e^{-\Gamma t}$$

$$\Gamma = \langle A_\nu | i \partial_t | A^\nu \rangle_{\infty=0} / 2 + cc., \quad m = 1$$

$$\int_0^\infty W(t) dt = 1$$

Γ is the gross mechanical energy of electrons. It leads to the result between decay life and this mechanical emission. To prove this result with the condition 2.1, vary the measure from $\sigma = 1$ to set this equation, at last $m = 1$. As varying, every atomic term is affected including the parts that out of unit variation.

The distribution shape of decay can be explain as

$$A_0 e^{-\Gamma t/2 - i k_x t}, \quad 0 < t < \Delta$$

It's the real wave of the particle x near the initial time and expanded in that time span

$$\approx \sum_k \frac{C e^{-ikt}}{k - k_x - i\Gamma/2}$$

14. GRAND UNIFICATION

The General Theory of Relativity is

$$(14.1) \quad R_{ij} - \frac{1}{2} R g_{ij} = -8\pi G T_{ij} / c^4$$

Firstly the unit second is redefined as S to simplify the equation 14.1

$$R_{ij} - \frac{1}{2}Rg_{ij} = -T_{ij}$$

Then

$$R_{ij} - \frac{1}{2}Rg_{ij} = \frac{1}{\mu}(F_{i\mu}^*F_j^\mu - g_{ij}F_{\mu\nu}^*F^{\mu\nu}/4)$$

F is conjugate and antisymmetric. We observe that the co-variant curvature is

$$R_{ij} = \frac{1}{\mu}(F_{i\mu}^*F_j^\mu + g_{ij}F_{\mu\nu}^*F^{\mu\nu}/8)$$

15. CONCLUSION

Fortunately, this model explained all the effects in the known world: strong, weak and electromagnetic effects, and even subclassify them further if not being to add new ones. In this model the only field is electromagnetic field, and this stands for the philosophical that the unified world is from an unique source,all that depend on the hypothesis: in electron the movements of charge and mass are the same.

My description of particles is compatible with QED elementarily and depends on momentum quantification formula, and only contributes to it with theory of consonance state in fact. In some way, the electron function is a good promotion for the experimental models of proton and electron that went up very early.

Underlining my calculations a fact is that the electrons have the same phase (electrons consonance), which the BIG BANG theory would explain, all electrons are generated in the same time and place, the same source.

REFERENCES

- [1] K.A. Olive et al. (Particle Data Group), Chin. Phys. C, 38, 090001 (2014) and 2015 update
E-mail address: hiyaho@126.com

TIANMEN, HUBEI PROVINCE, THE PEOPLE'S REPUBLIC OF CHINA. POSTCODE: 431700