## TITLE PAGE

"A new formula for the Riemann hypothesis "

**KEYWORDS**,Non standard analysis,Non standard rearrangement,Riemann Dini's theorem.Zeta function of Riemann

**Author: Ciro Cesarano**

## INTRODUCTION

Since the monumental Riemann's article of 1859 [9] the demonstration of what will become the Riemann hypothesis,it was and still is the central problem of number theory and all mathematics,having profound implications in various fields of it,and even in quantum mechanics [2] .From Riemann onwards, a great number of famous mathematicians have tried in vain to prove it.Just to name a few Hardy,Littlewood,Hilbert,and in more recent times,Weil and others[2]. Furthermore the Riemann hypothesis is one of seven "Millennium Problems" [1].In this article the author tackles the problem of proving the Riemann hypothesis in a completely new different way from previous attempts,in fact in this article we will use some fundamental concept of non standard analysis founded by mathematical logician Abraham Robinson in the sixties of twenty century [10].Subsequently it was greatly simplified,without losing in mathematical rigor,by Jerome Keisler [6]which is the approach used in this article.

## DEFINITION AND PRELIMINARIES

We give here some concept and definition that we will use for the continuations of this article.Foundamentally Keisler approach is based on the following two principles [6][7].

## THE EXTENSION PRINCIPLE

a) The real numbers form a subset of the hyperreal numbers, and the order relation $x< y$ for the real numbers is a subset of the order relation for the hyperreal numbers
b)There is a hyperreal number that is greater than zero but less than every positive real number
c) For every real function f of one or more variables we are given a corresponding hyperreal function $f^*$ of the same number of variables , $f^*$ is called the natural extension of $f$ (in this article $f^*$ is called the extension of $f$ at non standard model of analysis) .
Furthermore with each relation X on $R$ there is corresponding relation X* on $R$* called the natural extension of X [7]

## THE TRANSFER PRINCIPLE

Every real statement that hold for one or more particular real function holds for the hyperreal natural extensions of these functions, the transfer principle is equivalent to Leibniz' principle ,which is the property that for each real bounded sentence $\phi \in L$,is true if and only if $\phi^*$ is true . L is the language of the first order predicate [7]we still give the following definitions always of Keisler [6]:

## DEFINITIONS

A hyperreal number b is said to be:
positive infinitesimal if b is positive but less than every positive real number,
negative infinitesimal if b is negative but greater than every negative real number.

A hyperreal number b is said to be:
finite if b is between two real numbers,
positive infinite if b is greater than every real number,
negative infinite if b is less than every real number.
Finally in the text we will assume that zeta function is an analytic function throughout $C - 1$,and monodromous as demonstrated by Riemann in 1859 [9]

## ABSTRACT

By a simple extension of rearrangement definition of a simply converging series, at non standard model of analysis, the author finds a new formula for $\zeta^*(s)$ with $s \in C^*$ $complex\ hyperreal\ numbers\ set$ with $a^* > 0$ and with $s \neq 1$ .
However this new formula is very easily extendable at the whole complex plane with $s \neq 1$ .
Notable result is that with the definition of " non standard rearrangement " the commutative property of addition continues to hold even for simply convergent series (such as harmonic series with alternate signs).
Moreover the author, by means of the new formula of $\zeta^*(s)$ and the corresponding functional equation, gives a proof of the Riemann hypothesis.

## TEXT

The new formula (for the zeta function of Riemann) is the follower:

$$\zeta^*(s) = st\left(\sum_1^{\omega} n^{-s} - \frac{\omega^{-s+1}}{-s+1}\right) \qquad (I)$$

$s \in C^*$ $with\ form\ a + ib$ ( $a$ $and\ b\ hyperreal\ numbers$ ) $s \neq 1$ and $a^* > 0$ ( $^*>$ hyperreal order relation [7] ). (I) has a simple pole at $s = 1$ where : ω=infinite number positive (see definitions and preliminaries);*=asterisk indicates the extension of $\zeta(s)$ at non standard model of analysis (see definitions and preliminaries).By "$st$ "we indicate the standard part [6] [7] [10] of the expression in the right-hand side, that is, the real number closest to the hyperreal quantity in the right-hand side. By definition, the difference between the standard part of a hyperreal number and itself hyperreal number is an infinitesimal quantity. Mathematicians such as Euler and Leibniz "neglected" such infinitesimal quantities by setting them equal to zero in their calculations.
For the demonstration of Riemann's hypothesis we also use the functional equation of $\zeta(s)$:

$$\zeta(1-s) = 2^{1-s}\pi^{-s}\sin\left(\frac{1-s}{2}\pi\right)(s-1)!\,\zeta(s) \qquad (II) \qquad [2].$$

Applying to (I) the Euler-Maclaurin formula (E.M.F) we have:

$$\zeta^*(s)=\sum_1^R n^{-s}-\frac{(R+1)^{-s+1}}{-s+1}+\frac{1}{2}(R+1)^{-s}-\frac{1}{12}\left[-s(R+1)^{-s-1}\right]+$$

$$+\frac{1}{720}\left[-s(-s-1)(-s-2)R+1)^{-s-3}\right]+\ldots\ldots \quad \text{(III)}$$

R $\in N$ =1,2,3…

In this article by convention we will indicate this formula (E.M.F) without distinction with its extension at non standard model of analysis (see definitions and preliminaries), therefore also when this formula is applied in $\mathrm{R}^*$ (hyperreal numbers set) or in $\mathrm{C}^*$ complex hyperreal numbers set or with form $s^* = a + \mathrm{ib}$ (with $a$ and b hyperreal numbers).

It is easy to see that (I) with $a$ *>1 ( *> hyperreal order relation [7]) becomes:

$$\zeta^*(s)=\sum_1^\omega n^{-s}$$

that is the extension at non standard model of analysis of the classical definition:

$$\zeta(s)=\sum_1^\infty n^{-s} \text{ with } a>1 \text{ [1][2][3][9]}.$$

Therefore by principle of identity of analytic functions (and for the transfer principle,see definitions and preliminaries)

(I) is the analytic continuation of

$$\zeta^*(s)=\sum_1^\omega n^{-s} \text{ (with } a^*>1)$$

In fact it is not difficult to prove with the (E.M.F) that (I) converges with $a$ *>0.

We can formalize and generalize this in the following way.Let F($s$) and G($s$) be two analytic functions in a region $of\ C$ $R_1$ and $R_2$ respectively with $R_1 \cap R_2 \neq \emptyset$ (a small arc in common is sufficient)

$(\forall s \in (R_1 \wedge R_2)\ (F(s)=G(s)) \rightarrow (\forall s \in (R_1 \vee R_2)\ (F(s)=G(s))$.
estendit it at non standard model we have:

$(\forall s^* \in (R^*_1 \wedge R^*_2)\ (F^*(s) = G^*(s)) \rightarrow (\forall s^* \in (R^*_1 \vee R^*_2)\ (F^*(s) = G^*(s))$.
since the first formula written is true in $\mathbb{C}$ ( set of complex numbers), in fact it is represents the principle of identity of analytic functions which also guarantees that the zeta function admits a unique analytic continuation, the last formula is true too in $\mathbb{C}^*$ (complex hyperreal numbers set) according to the transfer principle or Leibniz' principle , in fact standard and non standard models are elementary equivalents ,it is not possible to find a formula ( of the first order language) that is true in one model that is not true in the other model ( see the transfer principle).
Moreover (III) permits to find all values of $\zeta^*(s)$ ( and therefore of $\zeta(s)$ )

for example:

For $\zeta^*(0)$ = ? using (III) and R=1

$\zeta^*(0) = 1-2+\frac{1}{2}(\frac{1}{2})^0 = -\frac{1}{2}$

For $\zeta^*(-1)$ = ? (R=1)

For $\zeta^*(-1) = 1-\frac{2^2}{2}+\frac{1}{2}\cdot 2 - \frac{1}{12}(1\cdot 2^0) = \frac{-1}{12}$

For $\zeta^*(-2)$ = ? (R=1)

$\zeta^*(-2) = 1-\frac{2^3}{3}+\frac{1}{2}2^2 - \frac{1}{12}(2\cdot 2) = 0$ (the first trivial zero of $\zeta(s)$)

For $\zeta^*(\frac{1}{2})$ = ? (R=3)

$\zeta^*(\frac{1}{2}) = 1+2^{-\frac{1}{2}} + 3^{-\frac{1}{2}} - \frac{4^{\frac{1}{2}}}{\frac{1}{2}} + \frac{1}{2}4^{-\frac{1}{2}} - \frac{1}{12}\left[-\frac{1}{2}\bullet 4^{-1,5}\right] +$
$\frac{1}{720}\left[-\frac{1}{2}\cdot -1,5\cdot -2,5\cdot 4^{-3,5}\right] + \ldots =$

$= -1,460354\ldots$

To deduce (III) it was taken into account that $\frac{\omega^{-s+1}}{-s+1}$ it is the primitive of the (non standard ) function $n^{-s}$calculates at the ω point.If you want to find the same formula (III),but obtained with methods of classical analysis,and in ways very different from those of this article see [3].Moreover for the error term to be attributed to (III),where the derivatives of odd order do not vanish,as in the

last example seen above, following Edwards [3] “it’s a general rule of thumb in applying the (E.M.F.) that as long as terms are decreasing rapidly in size,the bulk of the errors is in the term omitted".

Now we prove (I).

Stated harmonic series with alternate signs :

$$S = \sum_{1}^{\infty} \frac{(-1)^{n+1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \ldots = \ln 2 \qquad (1)$$

extending it at “non standard model” we have

$$st\left(\sum_{1}^{\omega} \frac{(-1)^{n+1}}{n}\right) = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \ldots = \ln 2 \qquad (1)^*$$

with ω = infinite number positive .

We call “non standard rearrangement” series of (1)*

a series with both the conditions following :

1)The rearranged series of (1)* must consist of all the same standard terms (finite numbers) contained in (1)* and no other terms.

2) The rearranged series of (1)* must consist of all the same terms non standard (infinitesimal numbers) contained in (1)* and no other terms.

Since real numbers are a subset of hyperreals ( see extension principle) the non standard rearrangement could be defined by a single condition ,but to Better highlight the difference between two rearrangements two were written.We give an example of standard rearrangement (using only (1) condition) and correspondent “non standard rearrangement” (using (1) and (2) conditions).

It knows that:

$$\left(1 - \frac{1}{2} - \frac{1}{4}\right) + \left(\frac{1}{3} - \frac{1}{6} - \frac{1}{8}\right) + \left(\frac{1}{5} - \frac{1}{10} - \frac{1}{12}\right) + \ldots =$$

$$= \sum_{n=1}^{\infty} \left(\frac{1}{2n-1} - \frac{1}{4n-2} - \frac{1}{4n}\right) = \underline{S} = \frac{1}{2} S = \frac{Ln2}{2} \qquad (2).$$

(2) is an example of “standard rearrangement "of (1).

According to Riemann-Dini ‘s theorem the result of (2) is different from result of (1),

on the contrary using the first and second condition we rewrite $(2)^*$ as :

$$(\sum_{1}^{\frac{\omega}{2}} \frac{1}{2n-1} - \sum_{1}^{\frac{\omega}{4}} \frac{1}{4n-2} - \sum_{1}^{\frac{\omega}{4}} \frac{1}{4n})$$

considering that: $\sum_{1}^{\frac{\omega}{4}} \frac{1}{4n-2} = \frac{1}{2} \sum_{1}^{\frac{\omega}{4}} \frac{1}{2n-1}$

we have:

$$\frac{1}{2}(\sum_{1}^{\frac{\omega}{4}} \frac{1}{2n-1} - \sum_{1}^{\frac{\omega}{4}} \frac{1}{2n}) + \sum_{\frac{\omega}{4}+1}^{\frac{\omega}{2}} \frac{1}{2n-1} \qquad (2)^* \quad \text{with :}$$

$$st(\sum_{1}^{\frac{\omega}{4}} \frac{1}{2n-1} - \frac{1}{2n})) = S^* = \ln 2$$

(it is the harmonic series with alternating signs calculated until to ω/4, by the Cauchy convergence test in non standard analysis [3] we have that the sum of harmonic series with alternating signs is invariant ,to less than infinitesimals ,from the particular infinite number that we use for to calculate it ,for example as in this case ω/4,or others infinite numbers as ω/2,or ω,or ω²,or….etc ) and using E.M.F.(Euler-Maclaurin formula):

$$\text{St}(\sum_{\frac{\omega}{4}+1}^{\frac{\omega}{2}} \frac{1}{2n-1}) = St\,(\int_{\frac{\omega}{4}}^{\frac{\omega}{2}} \frac{1}{2n-1} dn) = \frac{1}{2} Ln2$$ therefore we have:

$$\frac{1}{2}\ln 2 + \frac{1}{2}\ln 2 = S^* = \ln 2 \qquad (2)^*$$

This result confirms validity of commutative property of addition in opposition to Riemann-Dini’s theorem (or Riemann's rearrangement theorem) in fact:

$$st(\sum_{\frac{\omega}{2}+1}^{\omega} \frac{1}{2n-1} - \sum_{\frac{\omega}{4}+1}^{\omega} \frac{1}{4n-2} - \sum_{\frac{\omega}{4}+1}^{\omega} \frac{1}{4n}) = -\frac{1}{2} \ln 2 \qquad (3)^*$$

Since $(3)^*$ does not include in $(1)^*$ we must remove it (according to the second conditions before seen) from the simple extension at non standard model of (2) that is :

$st(\sum_{1}^{\omega}(\frac{1}{2n-1}-\frac{1}{4n-2}-\frac{1}{4n}))$ $\qquad\qquad$ $(2^{bis})^*$

therefore we have :

$(2^{bis})^* - (3)^* = (2)^* = S^* = \ln 2$

Now we generalize and formalize the two previous conditions which together constitute the new concept of "non standard rearrangement"of a series,for the first condition that characterize the classical standard rearrangement of a series we have:

give the series $\sum_k a_k$ with real or complex terms and one bijective function

$\pi : N \to N$ it's called rearranged series of $\sum_k a_k$ according $\pi$ , the series $\sum_k a_{\pi(k)}$ .

The second condition is the following:

given the series $\sum_k a_k{}^*$ with non standard and complex terms (infinitesimal complex and infinitesimal real numbers) and one bijective non standard function (see definition and preliminaries)

$\pi^* : N^* \to N^*$ ($N^*$ are also called infinite hypernatural numbers [6]).

It is called rearranged series according to $\pi^*$ the series:

$$\sum_k a_{\pi^*(k)}$$

The biunivocity of $\pi$ and $\pi^*$ ensure in particular that the number of terms of the rearranged series have the same number of terms as the originals series respectively in $N$ and in $N^*$.Note that Riemann - Dini theorem concerns simply convergent series in $\mathrm{R}$, while the concept of "non standard rearrangement" is

easily extendable at $\mathrm{C}$ and $\mathrm{C}^*$.It is shown in standard analysis (Ohm's rearrangement theorem) that in (1) taking $p$ positive and $q$ negative terms we have the sum :

$S$ = ln(2)+(1/2)ln($p$/$q$).In the previous standard example it was $p$ = 1 and $q$ = 2 instead with our definition of "non standard rearrangement" it is always $p^*$ = $q^*$ both infinite numbers in the previous non standard example it was

$p^*$ = ω/2 and $q^*$ = ω/4+ω/4 = ω/2 .

The validity of the commutative property in (1)* should not be surprising as this property is obviously valid for real numbers,according to the transfer principle it is also valid for hyperreal numbers in fact we have:

$(\forall x\ \forall y)\in \mathrm{R}\ (x+y=y+x)$
it is true in $\mathrm{R}$,extending it at non standard model we have :

$(\forall x^*\ \ \forall y^*) \in \mathrm{R}^*\ (x^*+y^*=y^*+x^*)$
($\mathrm{R}^*$= set of hyperreal numbers)
( Following common usage we omit the asterisk for the sum between hyperreal numbers).

Why is the Riemann -Dini theorem valid with " standard rearrangement" despite the commutativity of addition?The answer is simple in fact formalizing this theorem (by the first order predicate logic) we have: let $\sum_k a_k$ a simply convergent series

$(\forall x \in \mathrm{R} \cup \{-\infty,+\infty\}\ \exists \pi : N\leftrightarrow N)$:Mi ha mi mi p non

$$\lim\ N \to \infty \sum_{k=1}^{N} a_{\pi(k)} = x$$

Extending at non standard model we have:

$(\forall x^* \in \mathrm{R}^*\ \exists \pi^* : N^*\leftrightarrow N^*)$ :

$$\sum_{k=1}^{\omega} a_{\pi^*(k)} = x$$

(remembering that standard part $st$ is defined only for finite hyperreal numbers that is, only for complex hyperreal numbers with finite modulus) .It is precisely the limit with $N \to \infty$ $\sum_{k=1}^{N} a_{\pi(k)}$ or in non standard model $\sum_{k=1}^{\omega} a_{\pi^*(k)}$ that introduces an infinite quantity of infinitesimals whose sum is different from zero not existing in the original series (1)* these quantities are excluded only with the" non standard rearrangement" as seen in (3)*.

Now we study the following series with alternate signs:

$\eta(s) = 1 - \frac{1}{2^s} + \frac{1}{3^s} - \frac{1}{4^s} + \ldots \qquad (s \in C)$

it is convergent (with $a > 0$) [2] and we have

$\eta(s) = (1 + \frac{1}{2^s} + \frac{1}{3^s} + \ldots) - 2(\frac{1}{2^s} + \frac{1}{4^s} + \frac{1}{6^s} + \ldots.) =$

$= \zeta(s) - \frac{2\zeta(s)}{2^s} = (1 - \frac{2}{2^s})\,\zeta(s)$ (4)

but extending (4) at non standard model we have:

$\eta^*(s) = st[\sum_{1}^{\omega} n^{-s} - 2(\sum_{1}^{\omega} (2n)^{-s})]$ (4)*

but the last series on the right following the second condition of the " non standard rearrangement" seen before is not correct because all terms of:

$\sum_{\frac{\omega}{2}+1}^{\omega} (2n)^{-s}$ are not in $\zeta^*(s) = \sum_{1}^{\omega} n^{-s}$

therefore the correct formula is:

$\eta^*_{correct}(s) = St\ [\sum_{1}^{\omega} n^{-s} - \frac{2}{2^s}(\sum_{1}^{\omega} n^{-s} - \sum_{\frac{\omega}{2}+1}^{\omega} n^{-s})\ ]$ (5)*

in (5)* we have now $p^* = q^* = \omega$.Furthermore since (5)* is convergent with $a >^* 0$ we can apply the Cauchy convergence test which extended at non standard model [6] tells us as seen before that the value of (5)* is invariant for the particular infinite number that we use for p* e q* for example $2\omega$, $\omega/2$, $\omega$,... $\omega^2$etc

, the thing important for the " non standard rearrangement" of (5)* is that always is p* = q*)

therefore we have:

$\zeta^{*}(s)= \frac{\eta^{*}(s)}{1-\frac{2}{2^{s}}}$ which begins with our correction: .

$$\zeta^{*}(s) = \frac{\eta^{*}(s)_{correct}}{1-\frac{2}{2^{s}}} = \left(\sum_{1}^{\omega} n^{-s} + \frac{1}{2^{s-1}-1} \bullet \sum_{\frac{\omega}{2}+1}^{\omega} n^{-s}\right) \qquad (5^{*}bis)$$

using ( E.M.F.) we have with $s \neq 1$.

$$\frac{1}{2^{s-1}-1} \bullet \sum_{\frac{\omega}{2}+1}^{\omega} n^{-s} = \sum_{\frac{\omega}{2}}^{\omega} n^{-s} - \omega^{-s} = \left\| \frac{n^{-s+1}}{-s+1} \right\|_{\frac{\omega}{2}}^{\omega} \frac{\omega^{-s+1}}{-s+1} - \frac{\left(\frac{\omega}{2}\right)^{-s+1}}{-s+1}$$

$+ \frac{1}{2^{s-1}-1}$ [ ( $-\frac{1}{2}\ \omega^{-s}$ + $\frac{1}{2}$( $\frac{\omega^{-s}}{2}$ ) +.....] This quantity in square brackets ,with a real part $a$ > 0, is an infinitesimal number which can be denoted with I( $s$)] .At last we have (after simple calculations):

$$\zeta^{*}_{correct}(s)= \zeta^{*}(s) = St\left(\sum_{n=1}^{\omega} n^{-s} - \frac{\omega^{-s+1}}{-s+1}\right) \qquad (I)$$

with $s \in C$ * and $\neq 1$ (with $a$ *>0).

Now we demonstrate using (I) and (II) Riemann hypothesis (RH).However, let us first prove some facts concerning the zeros of $\zeta(s)$ (such facts can be extended to $\zeta^{*}(s)$ in a trivial way).

In (II) replacing $s$ with 2n+1 we have :

$\zeta(-2n)=2^{-2n}\pi^{-2n-1}\sin(-n\pi)(-2n)!\zeta(2n+1)$ (6).

The only factor that can be canceled at the second member of (6) is

$\sin(-n\pi) = 0 \quad \forall n \in N$

therefore (6) admits zeros with $s$ =−2n (−2, −4, −6,… $-2n$),they are called trivial zeros of $\zeta(s)$,and they do not concern the (RH) [2].

It is easy to prove using the Euler's product formula [1][2][3][8]

$\zeta(s) = \prod_{p=primes} \frac{p^s}{p^s-1}$ with $a > 1$ , $(s \in C)$

(p = primes = 2,3,5,7,….) that doesn't exist $s \in C$ for which $p^s$=0 (since the numerator never vanishes).

Therefore $\zeta(s)$ with $a > 1$ does not admit zeros,and using (II) we have that $\zeta(s)$ does not admit zeros with $a < 0$ (except for the trivial zeros).

Therefore all non trivial zeros of $\zeta(s)$ are in the range $0 \leq a \leq 1$.

In 1896 La Vallée' Poussin[8] and Jacques Hadamard [4] independently proved that $\zeta(s)$ has not zeros with $a$ = 1,and therefore using (II) it has not zeros even with $a$ = 0, so all non trivial zeros are in the range $0 < a < 1$ (this range is called "the critical strip" [2][3]) and we can use in this range the formula (I) and the corresponding equation below:

$\zeta^*(1-s) = st\,(\sum_1^{\omega} n^{s-1} - \frac{\omega^s}{s})$ with $s \in C^*$ and $s \neq 0$ (IV)

(IV) has a simple pole at $s = 0$ ,obtained from (I) replacing $s$ with $1-s$

in fact (IV) is valid with $a^* < 1$.Now let $s_k$ be the ( non trivial ) zeros of $\zeta^*(s)$ (with $k \in \mathrm{N}=1,2,3,\ldots$) by (II)* we obtain:

$\zeta^*(s_k) = \zeta^*(1 - s_k) = 0.$ $(V)$

Since in the critical strip the only factor of (II)* than can be canceled is precisely $\zeta^*(1 - s_k)$ [2].Obviously by (II)* we mean the extension of (II) at non- standard model i.e.

$\zeta^*(1\text{-}s) = [\, 2^{1-s}\, \pi^{-s}\, \sin^*(\frac{1-s}{2}\pi)(s-1)!^*\, \zeta^*(s)\,]$ $s \in \mathrm{C}^*$ (II)*

For (II)* as for all complex functions extended to the non- standard model,the obvious property applies $f^*(s) = f(s)$ $\forall s \in \mathrm{C}$ ( if $f(x)$ is real function $f^*(x) = f(x)$ $\forall x \in \mathrm{R}$). By means of $(V)$ we have that if $s_k = \frac{1}{2} + ib_k$ is a zero of $\zeta^*(s)$ then also $s_k = \frac{1}{2} - ib_k$ will be a zero of $\zeta^*(s)$ , but this is true also for all non trivial zeros by reflection principle [3] ( of Schwartz) , or the non trivial

zeros arrange themselves in pairs of conjugate complex ( hyper) real numbers [2]

Using (I) and (IV) into $(V)$ we obtain (with $s = s_k$):

$$(\sum_{1}^{\omega} n^{-s_k} - \frac{\omega^{-s_k+1}}{-s_k+1}) = (\sum_{1}^{\omega} n^{s_k-1} - \frac{\omega^{s_k}}{s_k}) = 0 \qquad \text{(VI)}.$$

( in reality Equation (VI) should be written as:

$$\sum_{1}^{\omega} n^{-s_k} - \frac{\omega^{-s_k+1}}{-s_k+1} = \mathrm{I}(s_k) \wedge$$

$$\sum_{1}^{\omega} n^{s_k-1} - \frac{\omega^{s_k}}{s_k} = \mathrm{I}(1-s_k)$$

where $\mathrm{I}(s_k)$ and $\mathrm{I}(1-s_k)$ are infinitesimal quantities derived from Eq. (5* bis); however, as will be demonstrated below, in reality $\mathrm{I}(s_k) = \mathrm{I}(1-s_k) = 0$ as the only possible solution).

Since the zeta function is a monodromous function [9] that is one value in the domain corresponds to one and only one value in the codomain,by the means of $(V)$ we have that the four elements of (VI) are also monodromous.It is easy to prove that (VI) is solved with

$$-s_k = s_k - 1 \quad \text{or} \quad -a_k \pm ib_k = a_k - 1 \pm ib_k$$

Separating the (hyper)real and the imaginary part we have:

$$-a_k = a_k - 1 \rightarrow a_k = \frac{1}{2}\,;\ \pm b_k i = \pm b_k i \quad (it\ is\ always\ walid\ )$$

Therefore (VI) is solved precisely by value admitted by the RH ($a_k = \frac{1}{2}$) for the non trivial zeros of $\zeta^*(s)$ (naturally also the real number $\frac{1}{2} \in \mathrm{R}^*$ it is a finite hyperreal).

But of course the (RH) absolutely demands that all non-trivial zeros must lie on the critical line $a_k = \frac{1}{2}$ in this regard it is good to recall a work of Hardy

[5] in which he demonstrates that ζ($s$) with $a_k = \frac{1}{2}$ admits infinitely many zeros.But infinitely many zeros do not mean all!

Alternatively (and equivalently) the (RH) can thus be stated “no zero of ζ*($s$) with $a_k$ *≠ $\frac{1}{2}$ can exist in the critical stripe” and this will be what we will prove.Note that $(V)$ is invariant under substitution : $s_k = 1 - s_k$ this substitution admits $s_k = \frac{1}{2} \pm b_k i$ as the only possible value for $(V)$ as we will now demonstrate ( that is we demonstrate the Riemann hypothesis).Now let $s_k = \frac{1}{2} + \varepsilon \pm b_k i$ with ε real number $0 \leq \varepsilon < \frac{1}{2}$ ( the case with ε positive infinitesimal we will see later)

If (VI) admits $s_k = \frac{1}{2} + \varepsilon \pm b_k i$ with $\varepsilon = 0$ as the only one solution

then (RH) is valid,if instead (VI) admits also only one solution with

ε≠0, $then\ (RH)\ is$ certainly false.Let therefore be fixed

$s_k = \frac{1}{2} + \varepsilon \pm b_k i$ with ( $0 \leq \varepsilon < \frac{1}{2}$) (VII).

(VI) in addition to trivial solution given equalizing (I) and (IV) to zero, admits another simple solution,it is easy to see that the other simple solution of (VI) is given by following chain of equalities (This solution is unique, provided we assume, for example, that $\sum_1^{\omega} n^{-s_k}$ = $\sum_1^{\omega} n^{s_k - 1}$ a statement that will be simply demonstrated in the sequel):

$$\sum_1^{\omega} n^{-s_k} = \frac{\omega^{-s_k+1}}{-s_k+1} = \sum_1^{\omega} n^{s_k-1} = \frac{\omega^{s_k}}{s_k} \qquad \text{(VIII).}$$

Since into (VIII ) no element can be zero, because we are in the critical strip,the (VIII) is the last possible solution of (VI) as it is easy verify as will be demonstrated below (as we will see shortly), for graphic simplicity we denote the four elements of (VI) in an orderly manner with A,B,C,D we can write (VI) as:

A−B=C−D=0 ⇒

A=B & C=D in particular from chain of equalities (VI) we have:

$A-B=C-D$ putting B and C to the other member :

$A-C=B-D$ that admits as ulterior solution (equalizing both members to zero)

$A=C$ & $B=D$ together the two solution can be writer as:

$A=B=C=D$ that is (VIII).

Strictly speaking, equation (VI) should have been written as $A-B=\mathrm{I}(s_k)$ and $C-D=\mathrm{I}(1-s_k)$ ,where both $\mathrm{I}(s_k)$ and $\mathrm{I}(1-s_k)$ are infinitesimal quantities derived from (5*bis) for $\zeta^*(s_k)$ and $\zeta^*(1-s_k)$ . Consequently, the aforementioned equality $A=B=C=D$ would be the only possible solution for ( VI) , for instance, $A=C$ i.e. $\sum_1^{\omega} n^{-s_k} = \sum_1^{\omega} n^{s_k-1}$ and, simultaneously, both $\mathrm{I}(s_k)$ and $\mathrm{I}(1-s_k)$ vanished to zero—the proof of which is rather straightforward and will be provided at the end of this paper; and for the time being, this will be taken for granted.Considering that by (VIII) in particular is :

$$\frac{\omega^{-s_k+1}}{-s_k+1} = \frac{\omega^{s_k}}{s_k} \qquad \text{(IX)}$$

dividing both the members of (IX) by the first member and inserting (VII) into (IX) we have : $\frac{\left(\frac{1}{2}-\varepsilon \pm ib_k\right)}{\left(\frac{1}{2}+\varepsilon \pm ib_k\right)} \omega^{2\varepsilon} = 1.$ (IX)

To the first member of (IX) the term $\omega^{2\varepsilon} = e^{2\varepsilon ln\omega}$ with $\varepsilon > 0$

is always infinite number ( this result for the extension principle is valid also for every positive hyperreal numbers excluding the case with $\varepsilon$ infinitesimal numbers that for definition is smaller than every positive real numbers even if $>^*0$ this case will be seen later) while the other term is finite term non -zero, therefore (IX) ( and therefore (VI) too) is not satisfied with $\varepsilon\ number\ real > 0$. For this the only admissible value for (IX) ( and therefore for (VI) too ) is $\varepsilon = 0$, in fact $\omega^{2\varepsilon}$ with $\varepsilon = 0$ is $e^{0ln\omega} = e^0 = 1$.

But this condition is necessary but not yet sufficient.In fact for (IX) to be satisfied it,must have for solution $\varepsilon = 0$ also the following finite term no-zero before seen:

$\frac{\left(\frac{1}{2}-\varepsilon \pm ib_k\right)}{\left(\frac{1}{2}+\varepsilon \pm ib_k\right)} = 1$ which admits ,effectively exactly the solution with $\varepsilon = 0$ (just substitute the value $\varepsilon = 0$ in the above equation).Since as just demonstrated there are no zeros of ζ*($s$) with $s_k = \frac{1}{2} + \varepsilon \pm ib_k$ with ε≠0 then for (II)* there can be no zeros of ζ($s$) with $s_k$ $= \frac{1}{2} - \varepsilon \pm i\, b_k$ $with\, \varepsilon \neq 0$ , $therefore\ now\ \forall s_k$ with $0 *< a_k *< 1$ (excluding the case with ε positive infinitesimal in (VII) and therefore in (IX)).The hypothesis remains that (IX) is also resolved with ε positive infinitesimal ,but this is not possible since the ratio of the two quantities in brackets in the first member of (IX) can never be a non- complex hyperreal number (i.e. without imaginary unit i) as instead is the second member of (IX) (or the number 1),except in the case before seen with $\varepsilon = 0$,or in the case with $b_k = 0$ ($but\ there\ is\ not$ non trivial zero for zeta function with $b = 0$ ), while $\omega^{2\varepsilon} = e^{2\varepsilon ln\omega}$ using Taylor series expansion for the extension principle we have : $e^{2\varepsilon ln\omega} = 1+2\varepsilon ln\omega + 2\varepsilon^2 ln^2\omega$ + ..... is not a complex number ( it is a pure hyperreal number or ∈ R*). For this reason finally the Riemann hypothesis is proven to be true, more precisely we have demonstrated that an extension at non standard model of ζ($s$) admits all its non trivial zeros only with $a_k = \frac{1}{2}$ this result can be transferred to the standard function ζ($s$) we in fact can write : be

$s_k * \in C*$ and $0 *< a_k *< 1$ $(\forall s_k * : \zeta^*(s_k)=0) \rightarrow a_k *= \frac{1}{2}$

or in standard model : be $s_k \in C$ and $0 < a_k < 1$

$(\forall s_k : \zeta(s_k)=0) \rightarrow a_k = \frac{1}{2}$. In .Since the first proposition is demonstrated true in C* the second proposition is true in C therefore Riemann hypothesis is demonstrated true also in C.Furthermore according to (VIII) the demonstration of the Riemann hypothesis is also equivalent to demonstration that the following equalities :

$$\sum_1^{\omega} n^{-s_k} = \sum_1^{\omega} n^{s_k - 1} \qquad (X)$$

is valid only with the value given by (VII) with $\varepsilon = 0$, or $s_k = \frac{1}{2} \pm b_k i$ $\forall s_k$ with $0 *< a_k *< 1$. In fact (X) is equivalently by means (VIII) to (IX) where we

already have demonstrated that the only admissible value is precisely that required for the validity of the Riemann hypothesis (*equation* (X) *will be proved directly later on by a simple application of Cauchy's convergence test)* .It is immediate to see by (VIII) that the two series of (X) are divergent since they are evaluated in the critical stripe . However (X) can be written equivalently if you prefer as :

$$\sum_{1}^{\omega}(n^{-s_k}-n^{s_k-1})=0. \qquad (X).$$

So this one series has a finite value. (X) finally in standard model becomes:

$$\sum_{1}^{\infty}(n^{-s_k}-n^{s_k-1})=0. \quad \text{with} \quad s_k \in C \text{ and } \quad 0<a_k<1$$ Now we see by relation $\zeta^{*}(s)=\frac{\eta^{*}(s)}{1-\frac{2}{2^{s}}}$ that the function $\eta^{*}(s)$ has exactly the same zeros as the function zeta. That is, if $s_k$ is a zero for one of them, it is necessarily also a zero for the other function.

$$\eta^{*}(s)=[\sum_{1}^{\omega}(2n-1)^{-s_k}-\sum_{1}^{\omega}(2n)^{-s_k}]=2^{-s_k}[\sum_{1}^{\omega}(n-\frac{1}{2})^{-s_k}-\sum_{1}^{\omega}n^{-s_k}]=0$$

(XI)

BY (XI) we have :

$$\sum_{1}^{\omega}n^{-s_k}=\sum_{1}^{\omega}(n-\frac{1}{2})^{-s_k} \qquad (XII)$$

BY (X) we have

$$\sum_{1}^{\omega}n^{-s_k}=\sum_{1}^{\omega}n^{s_k-1} \qquad (X)$$

Comparing (X) with (XII) we have therefore trivially :

$$\sum_{1}^{\omega}n^{-s_k}=\sum_{1}^{\omega}(n-\frac{1}{2})^{-s_k}=\sum_{1}^{\omega}n^{s_k-1} \qquad (XIII)$$

Therefore we have:

$$2^{s_k}\eta^{*}(s) \quad = \sum_{1}^{\omega} n^{-s_k} - \sum_{1}^{\omega} (n-\frac{1}{2})^{-s_k} = \sum_{1}^{\omega} n^{-s_k} - \sum_{1}^{\omega} n^{s_k-1} \quad = \frac{\omega^{-s_k+1}}{-s_k+1} - \frac{\omega^{s_k}}{s_k}$$

$= 0.$ (XlV)

(To deduce the last equality in (XlV), (VIII) was used) we have precisely at the last member of (XlV) a equality of (VIII) or $\frac{\omega^{-s_k+1}}{-s_k+1} = \frac{\omega^{s_k}}{s_k}$ which as we have demonstrated before is only possible with ε=0 therefore $[2^{s_k}\eta * (\frac{1}{2} + \varepsilon \pm b_k i)] = 0$ or $(since \;\; 2^{s_k} \; never \; cancels \; out)$ $[\eta * (\frac{1}{2} + \varepsilon \pm b_k i)] = 0$ is possible only with $\varepsilon = 0$ ,so all the zeros of $\eta^{*}(s)$ are on the line passing through $\frac{1}{2}$ and therefore all the zeros of $\zeta^{*}(s)$ are also on the line passing through $\frac{1}{2}$, that is, they are all on the critical line.Let it be now

$$\zeta^{*}(s) = \sum_{1}^{R} n^{-s} - \frac{(R+1)^{-s+1}}{-s+1} + \frac{1}{2}(R+1)^{-s} - \frac{1}{12}\left[-s(R+1)^{-s-1}\right] +$$

$$+\frac{1}{720}\left[-s(-s-1)(-s-2)R+1)^{-s-3}\right] + \ldots\ldots \qquad (III)$$

R ∈ $N^{*}$ =1,2,3,....

(III) It holds for every integer - finite or infinite- between 1 and ω. We can choose N* = $\frac{\omega}{2}$ ( $Any \; infinite \; integer \; number \; H \; would \; be \; fine$) and $b_k \in$ R(hence a finite Number) therefore we have:

$$\zeta^{*}(s_k) = \sum_{1}^{\frac{\omega}{2}} n^{-s_k} - \frac{(\frac{\omega}{2}+1)^{-s_k+1}}{-s_k+1} +$$

$$\frac{1}{2}(\frac{\omega}{2}+1)^{-s_k} + \frac{1}{2}(\omega+1)^{-s_k} - \frac{1}{12}\left[-s_k(\frac{\omega}{2}+1)^{-s_k-1})\right] +$$

$$\frac{1}{720}\left[-s_k(-s_k-1)(-s_k-2)(\frac{\omega}{2}+1)^{-s_k-3}\right] + \ldots\ldots = 0. \qquad (XV)$$

$$\zeta^*(1-s_k)=\sum_1^{\frac{\omega}{2}} n^{s_k-1}-\frac{(\frac{\omega}{2}+1)^{s_k}}{s_k}+\frac{1}{2}(\frac{\omega}{2}+1)^{s_k-1}+$$

$$+\frac{1}{2}(\omega+1)^{s_k-1}-\frac{1}{12}\left[(s_k-1)(\frac{\omega}{2}+1)^{s_k-2}\right]+$$

$$\frac{1}{720}\left[(s_k-1)(s_k-2)(s_k-3)(\frac{\omega}{2}+1)^{s_k-4}\right]+\ldots.=0. \qquad \text{(XVl)}$$

Since it is $by\ (V)\ \zeta^*(s_k)-\zeta^*(1-s_k)=0.$ (XVll) or (XV)−(XVl)=0.

Or

$$\sum_1^{\frac{\omega}{2}} n^{-s_k}-\frac{(\frac{\omega}{2}+1)^{-s_k+1}}{-s_k+1}+$$

$$\frac{1}{2}(\frac{\omega}{2}+1)^{-s_k}+\frac{1}{2}(\omega+1)^{-s_k}-\frac{1}{12}\left[-s_k(\frac{\omega}{2}+1)^{-s_k-1}\right]$$

$$+\frac{1}{720}\left[(s_k-1)(s_k-2)(s_k-3)(\frac{\omega}{2}+1)^{s_k-4}\right]$$

$$-\sum_1^{\frac{\omega}{2}} n^{s_k-1}+\frac{(\frac{\omega}{2}+1)^{s_k}}{s_k}-\frac{1}{2}(\frac{\omega}{2}+1)^{s_k-1}-\frac{1}{2}(\omega+1)^{s_k-1}+$$

$$+\frac{1}{12}\left[(s_k-1)(\frac{\omega}{2}+1)^{s_k-2}\right]+$$

$$-\frac{1}{720}\left[(s_k-1)(s_k-2)(s_k-3)(\frac{\omega}{2}+1)^{s_k-4}\right]+\ldots\ldots=0 \qquad \text{(XVll)}$$

It is easy seen that with $s_k=\frac{1}{2}\pm b_k i$ the equation (XVll) it is perfectly resolved,let's see if with $s_k=\frac{1}{2}+\delta\pm b_k i\ \ with\ 0\le\delta<\frac{1}{2}\ (\delta\wedge b_k)\in R$ there is at least one solution with $\delta\neq0$ in which case the Riemann hypothesis is false. If instead the solution $\delta=0$ is unique then the Riemann hypothesis is true, we observe that each term of (XV) and (XVI) can only be an infinitesimal number of a certain order (or in the case of the first two on the left in (XV) and (XVl) infinite numbers of a certain order), therefore for any finite number (different from zero) if it is multiplied by an infinitesimal term of (XV) or (XV), the corresponding order of the infinitesimal does not change (just as, if the quantity

is infinitely large, the order of infinity does not change when multiplied by any finite, non-zero number furthermore, the infinitesimals of the derivatives in equations (XV) and (XVI) increase by two orders at each step and have alternating signs)this implies—given that δ is a real number—that all the individual terms of (XV) and (XVI) possess an order of magnitude whose value cannot be nullified by other individual terms having mutually incompatible orders of magnitude, or a finite or infinite sum of the other terms,for example in (XV) the term:

$$\frac{1}{2}(\omega + 1)^{-\frac{1}{2}-\delta\pm b_k i}$$

it is an infinitesimal of order $(\frac{1}{2}+ \delta\ )$ $While\ corresponding\ \ term\ of\ \ (XVI)$

$\frac{1}{2}(\omega + 1)^{-\frac{1}{2}+\delta\pm b_k i}$ $it\ is\ an\ infinitesimal\ number\ of\ order\ (\frac{1}{2} -\delta)$

It is immediately apparent that they are equal, and therefore their difference is zero only with $\delta=0$. If instead we consider in (XV) the term

$$\frac{1}{12}\left[ - s_k(\frac{\omega}{2} + 1\ )^{-1,5-\delta\pm b_k i}\ \right]$$

It is an infinitesimal number of order $(1,5+ \delta)$, While in(XV) the corresponding term is $\frac{1}{12}\left[(s_k - 1)(\frac{\omega}{2} + 1\ )^{-1,5+\delta\pm b_k i}\ \right]$

which is infinitesimal term of order $( 1,5 \ - \delta)$ the two have the same order of infinitesimals only when $\delta=0$ that is in this case 1,5 since (XVll) all corresponding terms (those sharing the same order of infinity or infinitesimality) have coefficients that coincide only when $\delta=0$ (where $\delta\ is$ a positive real number less than one-half or possibly zero)and naturally have opposite signs ,as we shall soon see, the same applies to infinite numbers of (XVll). For example, let us consider the two infinite numbers given by the difference (XV) − (XVI) = 0.

$$-\frac{(\frac{\omega}{2}+1)^{-s_k+1}}{-s_k+1} + \frac{(\frac{\omega}{2}+1)^{s_k}}{s_k} \quad \text{that is} \quad -\frac{(\frac{\omega}{2}+1)^{0,5-\delta\pm b_k i}}{0,5-\delta\pm b_k i} + \frac{(\frac{\omega}{2}+1)^{0,5+\delta\pm b_k i}}{0,5+\delta\pm b_k i}$$

They cancel each other out only when $\delta=0$ , in fact their orders of infinity are respectively $(\frac{1}{2} - \delta) and\ (\frac{1}{2} + \delta)$ which vanish only when $\delta=0$ (it should be noted that if all the other terms in equation ( XVII) did not also vanish with $\delta=0$,

it would imply that no zeros exist for the zeta function in the interval between zero and one-half ! "In other words, were (XVII) not to vanish at $\delta=0$ it would imply that the zeta function possesses no zeros on the critical line as well namely for $a = \frac{1}{2}$). The same holds for the divergent series $\sum_{1}^{\frac{\omega}{2}} n^{-s_k}$ $- \sum_{1}^{\frac{\omega}{2}} n^{s_k-1}$

which according to (I) and ( IV) and (V) have the compatible order of infinity seen just above $(\frac{1}{2} - \delta)\ and\ (\frac{1}{2} + \delta)$ respectively ( the orders of infinity of this series are equal only with $\delta = 0$) . The procedure of eliminating pairs whose elements are equal and opposite to each other when $\delta=0$ extends indefinitely to all terms that lie beyond the truncation in (XVll) in fact a generic term of (XV)relative to the derivatives is (with $B_{2n}$ $Bernoulli's\ numbers$) :

$$\frac{1}{2n!} B_{2n} \quad -s_k(-s_k-1)(-s_k-2)(-s_k-3).....(-s_k-2n+2)$$
$$(\frac{\omega}{2}+1)^{-s_k-2n+1}$$

Whereas for (XVl) the general term ,starting from $n = 1$,that is the first derivative is:

$$\frac{1}{2n!} B_{2n} \quad (s_k-1)(s_k-2)(s_k-3).....(s_k-2n+1)(\frac{\omega}{2}+1)^{s_k-2n}$$

in fact is necessary but not sufficient condition for two infinitesimals of opposite signs to cancel each other out is that they must both have the same order of infinitesimal, i.e., for $s_k = \frac{1}{2} + \delta \pm b_k i$ always with being $\delta$ a real number either 0 or positive between 0 and $\frac{1}{2}$ $and\ b_k$ a real number therefore arbitrarily large but finite, for la (XV)we have an order of infinitesimal of $(-s_k - 2n + 1$ )that is ( $\frac{1}{2} + \delta + 2n$) while for (XVI) ( $s_k - 2n$) that is ( $\frac{1}{2} - \delta + 2n$) and ( $\frac{1}{2} + \delta + 2n$)= ($\frac{1}{2} - \delta + 2n$) only if $\delta=0$.

For example,comparing the first derivatives of ( XV) and (XVI) we have for the coefficients $-s_k = s_k - 1$ *and simultaneously the relative*

exponents must satisfy $-s_k - 1 = s_k - 2$ ,both conditions satisfied only with $a_k = \frac{1}{2}$ in fact in a similar way, it can be proven that the generic coefficients also:

$$-\frac{1}{2n!} B_{2n} \qquad s_k\left(-s_k - 1\right)\left(-s_k - 2\right)\left(-s_k - 3\right)\ldots\left(-s_k - 2n + 2\right)$$

and

$$\frac{1}{2n!} B_{2n} \qquad \left(s_k - 1\right)\left(s_k - 2\right)\left(s_k - 3\right)\ldots\left(s_k - 2n + 1\right)$$

admit only $\delta=0$ as the unique solution to cancel the pair of infinitesimals. This means that the Riemann hypothesis is valid.". Now we directly prove (X), which is merely a special case of (VIII). Previously, we implicitly assumed that if (VIII) held, then the Riemann hypothesis necessarily followed; however, (VIII) was always valid if and only if A=C $I(s_k)=I(1 - s_k) = 0$ . Conditions that we now prove to be necessary (moreover, just above we have already demonstrated that in fact $I(s_k)=I(1 - s_k) = 0$ . however, we will provide another simple proof below)

Let us consider (5*bis) without the standard part or other infinitesimal approximations,always with $s_k$zeros of $\zeta^*(s)$ *with* $0 \le \delta < \frac{1}{2}$ $(\delta \wedge b_k) \in$ R there is at

$$\zeta^*(s_k)=\left(\sum_1^{\omega} n^{-s_k} + \frac{1}{2^{s_k-1}-1} \bullet \sum_{\frac{\omega}{2}+1}^{\omega} n^{-s_k}\right) = 0. \qquad (5^* \text{ bis})$$

$\sum_{\frac{\omega}{2}+1}^{\omega} n^{-s_k} = \sum_{\frac{\omega}{2}}^{\omega} n^{-s_k} - \left(\frac{\omega}{2}\right)^{-s_k}$ Thus $(5^* \; bis)\; becomes\; \zeta^*(s_k)=\sum_1^{\omega} n^{-s_k}$

$$-\frac{\omega^{-s_k+1}}{-s_k+1} + \frac{1}{2^{s_k-1}-1}\{\left[-\frac{1}{2}\left(\frac{\omega}{2}\right)^{-s_k} + \frac{1}{2}\omega^{-s_k} +\right.$$

$$\left.-\frac{1}{12}\left(1 - 2^{s_k+1}\right) s_k\left[\omega^{-s_k-1} - \left(\frac{\omega}{2}\right)^{-s_k-1}\right]+\ldots\ldots\right\}=0 \qquad (XVlll)$$

Therefore:

$$\zeta^{*}(1-s_k)=\sum_{1}^{\omega} n^{s_k-1}-\frac{\omega^{s_k}}{s_k}+\frac{1}{2^{-s_k}-1}\{[-\frac{1}{2}(\frac{\omega}{2})^{s_k-1}+$$

$$+\frac{1}{2}\omega^{s_k-1}+\frac{1}{12}(1-2^{-s_k})(s_k-1)\,\omega^{s_k-2}+....]\}=0. \qquad \text{(IXX)}$$

Putting $s_k=\frac{1}{2}+\delta\pm b_k i \quad with \quad 0\le\delta<\frac{1}{2}$ and $b_k\in R$ . Multiplying both sides of (XVIII) by $(\frac{\frac{1}{2}+\delta\pm b_k i}{\frac{1}{2}-\delta\pm b_k i})\,\omega^{2\delta}$ equation (XVIII) becomes:

$$(\frac{\frac{1}{2}+\delta\pm b_k i}{\frac{1}{2}-\delta\pm b_k i})\,\omega^{2\delta}\bullet\sum_{1}^{\omega} n^{-\frac{1}{2}-\delta\pm b_k i}-\frac{(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i})\,\omega^{2\delta}\bullet\omega^{\frac{1}{2}-\delta\pm b_k i}}{\frac{1}{2}-\delta\pm b_k i}+(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i})\,\omega^{2\delta}\bullet$$

$$\frac{1}{2^{-\frac{1}{2}-\delta\pm b_k}-1}\{[-\frac{1}{2}(\frac{\omega}{2})^{-\frac{1}{2}-\delta\pm b_k i}+\frac{1}{2}\omega^{-\frac{1}{2}-\delta\pm b_k i}+$$

$$-\frac{1}{12}(1-2^{\frac{3}{2}+\delta\pm b_k i})(\frac{1}{2}+\delta\pm b_k)\,\omega^{-\frac{3}{2}-\delta\pm b_k i}]+.....\}=0$$

$It\ is\ worth\ noting\ that:\ \frac{(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i})\,\omega^{2\delta}\bullet\omega^{\frac{1}{2}-\delta\pm b_k i}}{\frac{1}{2}-\delta\pm b_k i}=\frac{\omega^{\frac{1}{2}+\delta\pm b_k i}}{\frac{1}{2}+\delta\pm b_k i}$ which is the second term of the left - hand side of $\zeta^{*}(1-s_k)$ calculated with $s_k=\frac{1}{2}+\delta\pm b_k i$ .Calling $K(\delta)=(\frac{\frac{1}{2}+\delta\pm b_k i}{\frac{1}{2}-\delta\pm b_k i})\,\omega^{2\delta}$ and calculating $K(\delta)$(XVIII)−(IXX)=0 that is $K(\delta)\,\zeta^{*}(s_k)-\zeta^{*}(1-s_k)=0-0=0$ (and moving all the infinitesimals it contains to the second member ,denoting them as $I(s_k)$ and $I(1-s_k)$ respectively ,*noting that in* $(XVIII)$ $\frac{(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i})\,\omega^{2\delta}\bullet\omega^{\frac{1}{2}-\delta\pm b_k i}}{\frac{1}{2}-\delta\pm b_k i}=\frac{\omega^{\frac{1}{2}+\delta\pm b_k i}}{\frac{1}{2}+\delta\pm b_k i}$ which is the second term of the left- band side of (IXX) evaluated at $s_k=\frac{1}{2}+\delta\pm b_k i$ we have :

$$K(\delta)(\text{XVIII})-(\text{IXX}) = K(\delta)\sum_{1}^{\omega} n^{-\frac{1}{2}-\delta\pm b_k i} - \frac{\omega^{\frac{1}{2}+\delta\pm b_k i}}{\frac{1}{2}+\delta\pm b_k i} + \frac{K(\delta)I(\ s_k)}{2^{-\frac{1}{2}+\delta\pm b_k}-1}\ ) -[$$

$$\sum_{1}^{\omega} n^{-\frac{1}{2}+\delta\pm b_k i} - \frac{\omega^{\frac{1}{2}+\delta\pm b_k i}}{\frac{1}{2}+\delta\pm b_k i} + \frac{I(1-s_k)\ ]}{2^{-\frac{1}{2}-\delta\pm i\, b_k}-1}] = K(\delta)\sum_{1}^{\omega} n^{-\frac{1}{2}-\delta\pm b_k i} -$$

$$\sum_{1}^{\omega} n^{-\frac{1}{2}+\delta\pm b_k i} = \frac{I(1-s_k)}{2^{-\frac{1}{2}-\delta\pm i\, b_k}-1} - \frac{K(\delta)\, I(\ s_k)}{2^{-\frac{1}{2}+\delta\pm b_k}-1}$$

Where $K(\delta)\ I(\ s_k)$is given by the following expression:

$$K(\delta)\ I(\ s_k) = \{(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i})\ \frac{\omega^{2\,\delta}}{2^{-\frac{1}{2}-\delta\pm b_k}-1} \bullet [\ -\frac{1}{2}(\frac{\omega}{2})^{-\frac{1}{2}-\delta\pm b_k i}$$

$$-\frac{1}{12}(\frac{1}{2}+\ \delta\pm b_k i)(\ 1\ -\ 2^{\frac{3}{2}+\delta\pm b_k i})\ \omega^{-\frac{3}{2}-\delta\pm b_k i} +\text{.....}]$$

Therefore $K(\delta)\ I(\ s_k) = \{(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i}) \bullet \frac{1}{2^{-\frac{1}{2}-\delta\pm b_k}-1}\}[$

$$-\frac{1}{2}\,2^{\frac{1}{2}+\delta\pm b_k i}\ \omega^{-\frac{1}{2}+\delta\pm b_k i} - \frac{1}{12}(1\ -\ 2^{\frac{3}{2}+\delta\pm b_k i})(\frac{1}{2}+\ \delta\pm b_k i)\ \omega^{-\frac{3}{2}+\delta\pm b_k i}$$

$$+\text{.....}]\}$$

Since the expression in the curly brackets can only yield finite complex numbers,then $K\ (\delta)\bullet I(\ s_k)$can only yield infinitesimal values for any $0\leq \delta < \frac{1}{2}$ and $b_k \in$ R therefore,it can be written as

$$K(\delta)\bullet\sum_{1}^{\omega} n^{-\frac{1}{2}-\delta\pm b_k i} - \sum_{1}^{\omega} n^{-\frac{1}{2}+\delta\pm b_k i} \approx 0. \qquad (\text{XX})$$

That is, the two expressions are almost identical,meaning that their differences is infinitesimal,given that $K(\delta)\ = (\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i})\ \omega^{2\,\delta}$ where the expression in parentheses is always a finite quantity $\forall\delta$ previously bounded,while $\omega^{2\,\delta}$ is always an infinite number $\forall\delta$ except for the single value $\delta=0$ for which $\omega^{0}=1$.Let us see under what conditions ( if any)equation (XX) is correct. Since both (5* bis) and its application to the functional equation (II)* are independent of the specific infinite hypernatural number chosen (in our case, the number ω), the standard part of the result in (XX) remains unchanged whether it is

calculated using one infinite hypernatural number or another (even if infinitely larger or smaller). In other words, the difference between (XX) evaluated at two distinct infinite hypernatural numbers can only be infinitesimal (see Cauchy's convergence test for example)

$$\sum_{1}^{\omega}(K(\delta)\bullet n^{-\frac{1}{2}-\delta\pm b_k i} - n^{-\frac{1}{2}+\delta\pm b_k i})\approx 0. \qquad \text{(XX)}$$

Since (XX) does not depend on the specific infinite number, the standard part of the sum of the series is preserved (i.e., it is zero). Applying Cauchy' s convergence test to ( XX) we have $\sum_{1}^{H}(K(\delta)\bullet n^{-\frac{1}{2}-\delta\pm b_k i}\, n^{-\frac{1}{2}+\delta\pm b_k i})\approx 0$ with H an infinite number ,(furthermore the relation always hold

$$\sum_{1}^{\omega}F(n) = \sum_{1}^{H-1}F(n) + \sum_{H}^{\omega}F(n)$$

Being $\sum_{1}^{\omega}F(n)\approx 0$ and $\sum_{1}^{H-1}F(n)\approx 0$ therefore also $\sum_{H}^{\omega}F(n)\approx 0$ that is:

$$\sum_{H}^{\omega}(K(\delta)\bullet n^{-\frac{1}{2}-\delta\pm b_k i} - n^{-\frac{1}{2}+\delta\pm b_k i})\approx 0\,. \qquad \text{(XXI)}$$

Let H be an infinite number to simplify the calculations, let us assume it is infinitely smaller than $\omega$ that is $\frac{H}{\omega}\approx 0$.We therefore have calculating (XXI)in detail:

$$\sum_{H}^{\omega}(K(\delta)\, n^{-\frac{1}{2}-\delta\pm b_k i} - n^{-\frac{1}{2}+\delta\pm b_k i}) = (\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i})\bullet \omega^{2\delta}[\frac{\omega^{\frac{1}{2}-\delta\pm ib_k}}{\frac{1}{2}-\delta\pm b_k i}$$

$$-\frac{H^{\frac{1}{2}-\delta\pm ib_k}}{\frac{1}{2}-\delta\pm b_k i} + \frac{1}{2}\omega^{-\frac{1}{2}-\delta\pm b_k i} + \frac{1}{2}H^{-\frac{1}{2}-\delta\pm b_k i} +$$

$$\frac{1}{12}[(\frac{1}{2} - \delta\pm b_k i)\ (\omega^{-\frac{3}{2}-\delta\pm b_k i} - H^{-\frac{3}{2}-\delta\pm b_k i})+.....] =[\frac{\omega^{\frac{1}{2}+\delta\pm ib_k}}{\frac{1}{2}+\delta\pm b_k i} -$$

$$\frac{H^{\frac{1}{2}-\delta\pm ib_k}\bullet\omega^{2\delta}}{\frac{1}{2}+\delta\pm b_k i}+$$

$$+\frac{1}{2}\left(\frac{1}{2}-\delta\pm b_k i\right)\left(\frac{\omega^{-\frac{1}{2}+\delta\pm ib_k}}{\frac{1}{2}+\delta\pm b_k i}\right)+\frac{1}{2}H^{-\frac{1}{2}-\delta\pm b_k i}\bullet\left(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i}\right)\bullet\omega^{2\delta}+$$

$$-\frac{1}{12}\left(\frac{1}{2}-\delta\pm b_k i\right)\left(\omega^{-\frac{3}{2}+\delta\pm b_k i}-\left(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i}\right)\bullet\omega^{2\delta}\bullet\left(H^{-\frac{3}{2}-\delta\pm b_k i}\right)+\ldots\ldots\ldots\right]$$

$$-\left[\frac{\omega^{\frac{1}{2}+\delta\pm ib_k}}{\frac{1}{2}+\delta\pm b_k i}-\frac{H^{\frac{1}{2}+\delta\pm ib_k}}{\frac{1}{2}+\delta\pm b_k i}+\frac{1}{2}\left(\frac{\omega^{-\frac{1}{2}+\delta\pm ib_k}}{\frac{1}{2}+\delta\pm b_k i}\right)+\frac{1}{2}H^{-\frac{1}{2}+\delta\pm b_k i}+\right.$$

$$\left.-\frac{1}{12}\left(\frac{1}{2}+\delta\pm b_k i\right)\left(\omega^{-\frac{3}{2}+\delta\pm b_k i}-H^{-\frac{3}{2}+\delta\pm b_k i}+\ldots\ldots\ldots\right]\approx 0.\right.$$ That is, explicitly writing only the non-infinitesimal terms

$$-\frac{H^{\frac{1}{2}-\delta\pm ib_k}\bullet\omega^{2\delta}}{\frac{1}{2}+\delta\pm b_k i}+\frac{H^{\frac{1}{2}+\delta\pm ib_k}}{\frac{1}{2}+\delta\pm b_k i}$$

$$+\frac{1}{2}H^{-\frac{1}{2}-\delta\pm b_k i}\bullet\left(\frac{\frac{1}{2}-\delta\pm b_k i}{\frac{1}{2}+\delta\pm b_k i}\right)\bullet\omega^{2\delta}+\ldots\ldots\ )\approx 0. \qquad \text{(XXI)}$$

We observe that the first term in (XXI) with $\delta > 0$ is always infinitely larger than all the others; therefore, for such values of δ, equation (XXI) cannot have a solution. The only remaining possibility is that $\delta = 0$, a value which is indeed compatible with (XXI) in fact from (XXI) only the second term $\frac{H^{\frac{1}{2}+\delta\pm ib_k}}{\frac{1}{2}+\delta\pm b_k i}$ is of the same order of infinity as the first term if and only if $\delta = 0$

$-\frac{H^{\frac{1}{2}-\delta\pm ib_k}\bullet\omega^{2\delta}}{\frac{1}{2}+\delta\pm b_k i}+\frac{H^{\frac{1}{2}+\delta\pm ib_k}}{\frac{1}{2}+\delta\pm b_k i}=0.$ Since we have $\omega^0 = e^{0ln\omega}=1$ it follows that K(0)=1 and

$-\frac{H^{\frac{1}{2}\pm ib_k}}{\frac{1}{2}\pm b_k i}+\frac{H^{\frac{1}{2}\pm ib_k}}{\frac{1}{2}\pm b_k i}=0$ . By inserting the only possible value, δ=0, into the other terms, the equation (XXI) is effectively equal to zero. This is still an acceptable value for equation (XXI) as well, because there are other infinitesimal terms to be considered. However, it is easy to see that with δ=0 since we have $\omega^0 = e^{0ln\omega}=1$ it follows that K(0)=1 and with $s_k = \frac{1}{2} \pm b_k i$ (δ=0) we obtain:

$$I(s_k)=I(1-s_k)=0$$

Therefore (XXI) becomes : $$(\sum_{H}^{\omega}(n^{-\frac{1}{2}-\delta\pm b_k i}-n^{-\frac{1}{2}+\delta\pm b_k i})=0.$$

While equation (XX) with $K(0)=1$ becomes:

$$\sum_{1}^{\omega} n^{-\frac{1}{2}\pm b_k i}=\sum_{1}^{\omega} n^{-\frac{1}{2}\pm b_k i}$$

That is, (XX) is exactly equal to (X) with $s_k=\frac{1}{2}\pm b_k i$ :

## REFERENCES


[1]E.Bombieri,"Problems of millennium:the Riemann hypothesis",2000".

[2]J.Derbyshire,"Prime obsession.Bernhard Riemann and greatest unsolved problem in mathematics' ',2003.

[3]H.M.Edwards,"Riemann's zeta function",1974

[4]J.S.Hadamard,"Sur la distribution des zéros de la fonction ζ($s$) et ses conséquences arithmétique",1896

[5]G.H.Hardy,"Sur le zeros de la fonction ζ($s$),1914

[6]G.H.Keisler ,"Elementary calculus:an infinitesimal approach",1976,2000 (second edition)

[7] G.H.Keisler," Foundations of infinitesimal calculus" 2007, online edition 2020 .

[8]C.H.J. La Valleè Poussin,"Sur la fonction de Riemann et le nombre de nombres premiers inferierieurs à une limite donnée",1896

[9]B.Riemann,"Ueber die Anzahl der Primzahlen unter gegebenen Gr$\ddot{o}$sse",1859

[10] A.Robinson,"Non standard analysis",1966