

# A Three Field Theory of Gravity

Bernd Schmutzler

Bernd.Schmutzler@gmx.de

*09126 Chemnitz, Germany*

May 5, 2025

## Abstract

We propose a tensor-vector-scalar field theory of gravity by defining a corresponding Lagrange function. Deriving the field equations we observe, that the coupling to the tensor field gives mass to the vector field. After considerations of stability and convergence we develop the systems of differential equations for two cases. In the static case of a central mass we start with the Schwarzschild metric as a first approximation and get a behavior corresponding to the observation of galaxy rotation for sufficiently large distances from the center. Using this result a second approximation gives a qualitatively stronger gravity for very large distances, which could possibly describe the cohesion of galaxy clusters. When the distance tends to infinity, the gravity returns to a Newton-like decay due to the mass term of the vector field. In cosmology we find an expansion of a homogeneous, isotropic, flat universe at a lower rate than expected, but with a time contraction towards the past. Together these effects cause an observed redshift of light with far distant sources, which suggests an accelerated expansion of the universe.

## 1 Introduction

There are different attempts to describe gravity without the need of dark matter by relativistic theories based on the Einstein tensor field and extended by vector and scalar fields, for example the Brans-Dicke theory [3], TeVeS by Bekenstein [1] or newer versions [22] of Modified Newtonian dynamics and MOG by Moffat [15]. In most cases a scalar field represents a generalization of the gravitational constant. In our theory, the scalar field should fill the gap between the observed gravity and that calculated from visible matter by

the Newton/Einstein theory. The scalar field is coupled to the vector field by the scalar potential, while the vector field is coupled to the tensor field. Only the tensor field has a direct source, which is the usual stress-energy tensor.

In the next section we define the Lagrange function consisting of the three parts representing tensor, vector and scalar field, while the fourth part corresponds to the stress-energy tensor. The following section starts with a decomposition of the coupling between tensor and vector field. It turns out that the term quadratic in the vector field represents a mass term, where the mass is given by  $\frac{3}{2}$  of the coupling constant. Next, we discuss the relation of the resulted system of equations to general relativity for small coupling constants. In the case of a flat metric we investigate the character of the equations for the vector field and determine the number of degrees of freedom for the remaining system using the Hamilton-Dirac analysis.

The following section is devoted to the spherically symmetric case of a central mass  $M$ . First, it turns out that we had to use Cartesian instead of spherical coordinates because of their intrinsic warping. We derive the time dependent equations, but reduce the consideration to the static case. Starting with the Schwarzschild metric we describe the behavior of the vector and the scalar field in powers and logarithms of the distance  $r$  to the central mass. In a first approximation the scalar field generates a gravity proportional to  $M^{\frac{1}{2}}r^{-1}$  multiplied by a logarithmic factor with a negative exponent. This corresponds to observations of galaxy rotations, which in the outer regions can be approximated by the square root of Newtonian gravity. Using this result for a second approximation of the metric, we get a higher rate of gravity proportional to  $M^{\frac{1}{2}}r^{-\frac{1}{2}}$  again multiplied by a logarithmic term for larger distances. Possibly, this can describe galaxy clusters, where the gravity given by the first approximation is too weak. Finally, for distances tending to infinity, the mass term dominates the equation for the vector field. Then, the scalar field generates a gravity proportional to  $r^{-2}$ , which corresponds to Newtonian gravity.

In the section about cosmology we apply the theory to a homogeneous, isotropic, flat universe, dominated by mass neglecting radiation. In contrast to the Robertson-Walker metric we need a variable factor for the time component. Using again the Ansatz of a power series with logarithmic factors, here depending on  $t$ , we get only a logarithmic expansion and a Hubble function proportional to  $t^{-1}$  up to a logarithmic term. On the other hand, there is a time contraction with a factor  $t^{-2}$  and a logarithmic term with negative exponent. For observed light from the early universe this gives a redshift with a factor  $t$  multiplied by a logarithmic term with positive exponent. Interpreted in the Robertson-Walker metric this suggests an accelerated expansion of the universe.

## 2 Lagrange formulation

In a Minkowski space with coordinates  $x^\alpha|_{\alpha=0}^3$ , where for simplicity we identify  $x^0$  with time  $t$  even if it should be  $\frac{1}{c}x^0$ , we will describe gravity by the metric tensor  $g = g_{\alpha,\beta}|_{\alpha,\beta=0}^3$  and two additional fields, a vector field  $\mathcal{U} = u_\alpha|_{\alpha=0}^3$  and a scalar field  $v$ . For the metric we will use the signature  $(-, +, +, +)$ , i.e. in absence of matter the metric should be  $g = \text{diag}(-1, +1, +1, +1)$ .

The Christoffel symbols  $\Gamma_{\alpha,\beta}^\gamma$ , the Ricci tensor  $\mathcal{R}_{\alpha\beta}$  and the Ricci scalar  $R$  will be considered as differential operators

$$\Gamma_{\alpha\beta}^\gamma(\partial) = \frac{1}{2} g^{\gamma\sigma} (\partial_\alpha g_{\sigma\beta} + \partial_\beta g_{\alpha\sigma} - \partial_\sigma g_{\alpha\beta}) \quad (1)$$

$$\mathcal{R}_{\alpha\beta}(\partial) = \partial_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \partial_\beta \Gamma_{\alpha\gamma}^\gamma(\partial) + \Gamma_{\alpha\beta}^\gamma(\partial) \Gamma_{\gamma\sigma}^\sigma(\partial) - \Gamma_{\alpha\sigma}^\gamma(\partial) \Gamma_{\gamma\beta}^\sigma(\partial) \quad (2)$$

with  $\partial = \partial_\alpha|_{\alpha=0}^3$ . One can show, that  $\mathcal{R}_{\alpha\beta}$  is symmetric in  $\alpha, \beta$ , what is only nontrivial for the second term of (2). To preserve this symmetry for more general differential operators  $D = D_\alpha|_{\alpha=0}^3$  we define

$$\mathcal{R}_{\alpha\beta}(D) := \frac{1}{2} [\mathcal{R}_{\alpha\beta}(\partial)|_{\partial \rightarrow D} + \mathcal{R}_{\beta\alpha}(\partial)|_{\partial \rightarrow D}] \quad (3)$$

$$R(D) := g^{\alpha\beta} \mathcal{R}_{\alpha\beta}(D) \quad (4)$$

The tensor part  $\mathcal{L}_T$  of the Lagrangian can now be defined by

$$\mathcal{L}_T = \frac{1}{2} R(\partial + \mu \mathcal{U}) \quad (5)$$

where  $\partial + \mu \mathcal{U} = \partial_\alpha + \mu u_\alpha|_{\alpha=0}^3$  and  $\mu$  is the coupling constant between the tensor and the vector field. The vector part  $\mathcal{L}_V$  and the scalar part  $\mathcal{L}_S$  of the Lagrangian are

$$\mathcal{L}_V = -\frac{1}{4} (\nabla_\alpha u_\beta - \nabla_\beta u_\alpha) (\nabla^\alpha u^\beta - \nabla^\beta u^\alpha) - \frac{1}{2} \nabla_\alpha u^\alpha \nabla_\beta u^\beta \quad (6)$$

$$\mathcal{L}_S = -\frac{1}{2} \nabla_\alpha v \nabla^\alpha v - V(\mathcal{U}) v^2, \quad V(\mathcal{U}) = \frac{1}{2} (m^2 + \nu^2 \nabla_\alpha u^\alpha \nabla_\beta u^\beta) \quad (7)$$

where  $\nu$  describes the strength of the coupling between the scalar and the vector field and the covariant derivative of a vector or a scalar field is defined by

$$\nabla_\alpha u_\beta = \partial_\alpha u_\beta - \Gamma_{\alpha\beta}^\gamma(\partial) u_\gamma, \quad \nabla_\alpha u^\beta = \partial_\alpha u^\beta + \Gamma_{\alpha\gamma}^\beta(\partial) u^\gamma, \quad \nabla_\alpha v = \partial_\alpha v \quad (8)$$

Note that  $\nabla_\alpha u^\alpha \nabla_\beta u^\beta = \left( \sum_\gamma \nabla_\gamma u^\gamma \right)^2 \geq 0$ .

Together with the matter part  $\mathcal{L}_M$ , which is defined later in the usual way, depending only on the metric  $g$  we can write the complete Lagrangian as

$$\mathcal{L} = \mathcal{L}_T + \mathcal{L}_V + \mathcal{L}_S + \mathcal{L}_M \quad (9)$$

and the action

$$\mathcal{S} = \int \mathcal{L} \epsilon d^4x, \quad \epsilon = \sqrt{-\det g} \quad (10)$$

### 3 Field Equations

First, we shall decompose the expression  $R(\partial + \mu\mathcal{U})$  in the tensor part of the Lagrangian. For

$$\Gamma_{\alpha\beta}^\gamma(\partial + \mu\mathcal{U}) = \Gamma_{\alpha\beta}^\gamma(\partial) + \mu \Upsilon_{\alpha\beta}^\gamma \quad (11)$$

with

$$\begin{aligned} \Upsilon_{\alpha\beta}^\gamma &= \frac{1}{2} g^{\gamma\sigma} (u_\alpha g_{\sigma\beta} + u_\beta g_{\alpha\sigma} - u_\sigma g_{\alpha\beta}) \\ &= \frac{1}{2} (\delta_\beta^\gamma u_\alpha + \delta_\alpha^\gamma u_\beta - g_{\alpha\beta} u^\gamma) \end{aligned} \quad (12)$$

we have

$$\begin{aligned} \mathcal{R}_{\alpha\beta}(\partial + \mu\mathcal{U}) &= \partial_\gamma [\Gamma_{\alpha\beta}^\gamma(\partial) + \mu \Upsilon_{\alpha\beta}^\gamma] - \frac{1}{2} \partial_\beta [\Gamma_{\alpha\gamma}^\gamma(\partial) + \mu \Upsilon_{\alpha\gamma}^\gamma] \\ &\quad - \frac{1}{2} \partial_\alpha [\Gamma_{\beta\gamma}^\gamma(\partial) + \mu \Upsilon_{\beta\gamma}^\gamma] + \mu u_\gamma [\Gamma_{\alpha\beta}^\gamma(\partial) + \mu \Upsilon_{\alpha\beta}^\gamma] \\ &\quad - \frac{1}{2} \mu u_\beta [\Gamma_{\alpha\gamma}^\gamma(\partial) + \mu \Upsilon_{\alpha\gamma}^\gamma] - \frac{1}{2} \mu u_\alpha [\Gamma_{\beta\gamma}^\gamma(\partial) + \mu \Upsilon_{\beta\gamma}^\gamma] \\ &\quad + [\Gamma_{\alpha\beta}^\gamma(\partial) + \mu \Upsilon_{\alpha\beta}^\gamma] [\Gamma_{\gamma\sigma}^\sigma(\partial) + \mu \Upsilon_{\gamma\sigma}^\sigma] \\ &\quad - [\Gamma_{\alpha\sigma}^\gamma(\partial) + \mu \Upsilon_{\alpha\sigma}^\gamma] [\Gamma_{\gamma\beta}^\sigma(\partial) + \mu \Upsilon_{\gamma\beta}^\sigma] \\ &= \mathcal{R}_{\alpha\beta}(\partial) + \mu [\partial_\gamma \Upsilon_{\alpha\beta}^\gamma - \frac{1}{2} \partial_\beta \Upsilon_{\alpha\gamma}^\gamma - \frac{1}{2} \partial_\alpha \Upsilon_{\beta\gamma}^\gamma \\ &\quad + \Gamma_{\alpha\beta}^\gamma(\partial) \Upsilon_{\gamma\sigma}^\sigma + \Gamma_{\gamma\sigma}^\sigma(\partial) \Upsilon_{\alpha\beta}^\gamma - \Gamma_{\alpha\sigma}^\gamma(\partial) \Upsilon_{\gamma\beta}^\sigma - \Gamma_{\gamma\beta}^\sigma(\partial) \Upsilon_{\alpha\sigma}^\gamma] \\ &\quad + \mu [u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} u_\beta \Gamma_{\alpha\gamma}^\gamma(\partial) - \frac{1}{2} u_\alpha \Gamma_{\beta\gamma}^\gamma(\partial)] \\ &\quad + \mu^2 [u_\gamma \Upsilon_{\alpha\beta}^\gamma - \frac{1}{2} u_\beta \Upsilon_{\alpha\gamma}^\gamma - \frac{1}{2} u_\alpha \Upsilon_{\beta\gamma}^\gamma + \Upsilon_{\alpha\beta}^\gamma \Upsilon_{\gamma\sigma}^\sigma - \Upsilon_{\alpha\sigma}^\gamma \Upsilon_{\gamma\beta}^\sigma] \end{aligned}$$

For the last term an easy calculation using (12) shows

$$\begin{aligned} \mathcal{R}_{\alpha\beta}(\mathcal{U}) &= u_\gamma \Upsilon_{\alpha\beta}^\gamma - \frac{1}{2} u_\beta \Upsilon_{\alpha\gamma}^\gamma - \frac{1}{2} u_\alpha \Upsilon_{\beta\gamma}^\gamma + \Upsilon_{\alpha\beta}^\gamma \Upsilon_{\gamma\sigma}^\sigma - \Upsilon_{\alpha\sigma}^\gamma \Upsilon_{\gamma\beta}^\sigma \\ &= -\frac{1}{2} u_\alpha u_\beta - g_{\alpha\beta} u_\gamma u^\gamma \end{aligned}$$

With the relations

$$\begin{aligned} \nabla_\gamma \Upsilon_{\alpha\beta}^\gamma - \nabla_\beta \Upsilon_{\alpha\gamma}^\gamma &= \partial_\gamma \Upsilon_{\alpha\beta}^\gamma - \partial_\beta \Upsilon_{\alpha\gamma}^\gamma \\ &\quad + \Gamma_{\alpha\beta}^\gamma(\partial) \Upsilon_{\gamma\sigma}^\sigma + \Gamma_{\gamma\sigma}^\sigma(\partial) \Upsilon_{\alpha\beta}^\gamma - \Gamma_{\alpha\sigma}^\gamma(\partial) \Upsilon_{\gamma\beta}^\sigma - \Gamma_{\gamma\beta}^\sigma(\partial) \Upsilon_{\alpha\sigma}^\gamma \\ \nabla_\gamma \Upsilon_{\alpha\beta}^\gamma - \nabla_\alpha \Upsilon_{\beta\gamma}^\gamma &= \partial_\gamma \Upsilon_{\alpha\beta}^\gamma - \partial_\alpha \Upsilon_{\beta\gamma}^\gamma \\ &\quad + \Gamma_{\alpha\beta}^\gamma(\partial) \Upsilon_{\gamma\sigma}^\sigma + \Gamma_{\gamma\sigma}^\sigma(\partial) \Upsilon_{\alpha\beta}^\gamma - \Gamma_{\beta\sigma}^\gamma(\partial) \Upsilon_{\gamma\alpha}^\sigma - \Gamma_{\gamma\alpha}^\sigma(\partial) \Upsilon_{\beta\sigma}^\gamma \end{aligned}$$

we obtain

$$\begin{aligned} \mathcal{L}_T &= \frac{1}{2} g^{\alpha\beta} \mathcal{R}_{\alpha\beta}(\partial + \mu\mathcal{U}) \\ &= \frac{1}{2} R(\partial) + \frac{1}{2} \mu g^{\alpha\beta} [\nabla_\gamma \Upsilon_{\alpha\beta}^\gamma - \frac{1}{2} \nabla_\beta \Upsilon_{\alpha\gamma}^\gamma - \frac{1}{2} \nabla_\alpha \Upsilon_{\beta\gamma}^\gamma] \\ &\quad + \frac{1}{2} \mu g^{\alpha\beta} [u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} u_\beta \Gamma_{\alpha\gamma}^\gamma(\partial) - \frac{1}{2} u_\alpha \Gamma_{\beta\gamma}^\gamma(\partial)] - \frac{9}{4} \mu^2 u_\gamma u^\gamma \end{aligned}$$

By renaming some indices we can now write

$$\mathcal{L}_T = \frac{1}{2} \mathcal{L}_T^{(0)} + \frac{1}{2} \mu \mathcal{L}_T^{(1)} + \frac{1}{4} \mu^2 \mathcal{L}_T^{(2)} + \frac{1}{2} \mu \nabla_\gamma \mathcal{K}_T^\gamma \quad (13)$$

with

$$\mathcal{L}_T^{(0)} = R(\partial) \quad (14)$$

$$\mathcal{L}_T^{(1)} = g^{\alpha\beta} \left[ u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} u_\beta \Gamma_{\alpha\gamma}^\gamma(\partial) - \frac{1}{2} u_\alpha \Gamma_{\beta\gamma}^\gamma(\partial) \right] \quad (15)$$

$$= u_\gamma \left[ g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} g^{\alpha\gamma} \Gamma_{\alpha\beta}^\beta(\partial) - \frac{1}{2} g^{\beta\gamma} \Gamma_{\alpha\beta}^\alpha(\partial) \right] \quad (16)$$

$$\mathcal{L}_T^{(2)} = 9 u_\gamma u^\gamma \quad (17)$$

$$\mathcal{K}_T^\gamma = g^{\alpha\beta} \Upsilon_{\alpha\beta}^\gamma - \frac{1}{2} g^{\alpha\gamma} \Upsilon_{\alpha\beta}^\beta - \frac{1}{2} g^{\beta\gamma} \Upsilon_{\alpha\beta}^\alpha = -3 u^\gamma \quad (18)$$

where  $\mathcal{K}_T^\gamma$  was simply calculated using (12).

Having defined the vector field as massless, we now see a mass term

$$\frac{1}{4} \mu^2 \mathcal{L}_T^{(2)} = \left( \frac{3}{2} \mu \right)^2 u_\gamma u^\gamma$$

generated by the coupling to the tensor field with a mass proportional to the coupling constant.

For the field equations we have to describe the variation of the action  $\mathcal{S}$  by the variation of  $\mathcal{L} \epsilon$  in the form

$$\delta[\mathcal{L} \epsilon] = \left[ \mathcal{M}_{\alpha\beta}^{(g)} \delta g^{\alpha\beta} + \mathcal{M}_\gamma^{(u)} \delta u^\gamma + \mathcal{M}^{(v)} \delta v + \nabla_\gamma \mathcal{K}^\gamma \right] \epsilon \quad (19)$$

Then, the field equations are

$$\mathcal{M}_{\alpha\beta}^{(g)} = 0 \text{ for } \alpha, \beta = 0, \dots, 3, \quad \mathcal{M}_\gamma^{(u)} = 0 \text{ for } \gamma = 0, \dots, 3, \quad \mathcal{M}^{(v)} = 0 \quad (20)$$

First, we have

$$\delta[\mathcal{L} \epsilon] = \left[ \delta \mathcal{L} - \frac{1}{2} \mathcal{L} g_{\alpha\beta} \delta g^{\alpha\beta} \right] \epsilon \quad (21)$$

since

$$\delta \epsilon = -\frac{1}{2} \epsilon^{-1} \delta(\det g) \quad \text{and} \quad \delta(\det g) = \det g g^{\alpha\beta} \delta g_{\alpha\beta} = \epsilon^2 g_{\alpha\beta} \delta g^{\alpha\beta}$$

One can show [4] (appendix A) that

$$\mathcal{A}^\gamma = g^{\alpha\beta} \delta \Gamma_{\alpha\beta}^\gamma(\partial) - g^{\alpha\gamma} \delta \Gamma_{\alpha\beta}^\beta(\partial) = g_{\alpha\beta} \nabla^\gamma \delta g^{\alpha\beta} - \nabla_\beta \delta g^{\beta\gamma} \quad (22)$$

Using this relation for the zero part of  $\mathcal{L}_T$  we get

$$\delta \mathcal{L}_T^{(0)} = \mathcal{R}_{\alpha\beta}(\partial) \delta g^{\alpha\beta} + \nabla_\gamma \left[ g_{\alpha\beta} \nabla^\gamma \delta g^{\alpha\beta} - \nabla_\beta \delta g^{\beta\gamma} \right] \quad (23)$$

The variation of the next part gives

$$\begin{aligned}
\delta\mathcal{L}_T^{(1)} &= \delta u_\gamma \left[ g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} g^{\alpha\gamma} \Gamma_{\alpha\beta}^\beta(\partial) - \frac{1}{2} g^{\beta\gamma} \Gamma_{\alpha\beta}^\alpha(\partial) \right] \\
&+ u_\gamma \left[ \delta g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} \delta g^{\alpha\gamma} \Gamma_{\alpha\beta}^\beta(\partial) - \frac{1}{2} \delta g^{\beta\gamma} \Gamma_{\alpha\beta}^\alpha(\partial) \right] \\
&+ u_\gamma \left[ g^{\alpha\beta} \delta\Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} g^{\alpha\gamma} \delta\Gamma_{\alpha\beta}^\beta(\partial) - \frac{1}{2} g^{\beta\gamma} \delta\Gamma_{\alpha\beta}^\alpha(\partial) \right] \quad (24)
\end{aligned}$$

We can write

$$\begin{aligned}
\delta u_\gamma &= \delta(g_{\gamma\rho} u^\rho) = g_{\gamma\rho} \delta u^\rho - u^\rho g_{\gamma\sigma} g_{\tau\rho} \delta g^{\sigma\tau} = g_{\gamma\rho} \delta u^\rho - u_\tau g_{\gamma\sigma} \delta g^{\sigma\tau} \\
&= g_{\gamma\rho} \delta u^\rho - \frac{1}{2} u_\tau g_{\gamma\sigma} \delta g^{\sigma\tau} - \frac{1}{2} u_\sigma g_{\gamma\tau} \delta g^{\sigma\tau}
\end{aligned}$$

to preserve symmetry of indices in the  $\delta g$  terms. With  $g_{\gamma\rho} g^{\alpha\gamma} = \delta_\rho^\alpha$  and applying (22) again we obtain

$$\begin{aligned}
\delta\mathcal{L}_T^{(1)} &= \left[ g_{\gamma\rho} g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} \Gamma_{\rho\beta}^\beta(\partial) - \frac{1}{2} \Gamma_{\alpha\rho}^\alpha(\partial) \right] \delta u^\rho \\
&- \frac{1}{2} \left[ g_{\gamma\sigma} g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} \Gamma_{\sigma\beta}^\beta(\partial) - \frac{1}{2} \Gamma_{\alpha\sigma}^\alpha(\partial) \right] u_\tau \delta g^{\sigma\tau} \\
&- \frac{1}{2} \left[ g_{\gamma\tau} g^{\alpha\beta} \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} \Gamma_{\tau\beta}^\beta(\partial) - \frac{1}{2} \Gamma_{\alpha\tau}^\alpha(\partial) \right] u_\sigma \delta g^{\sigma\tau} \\
&+ \left[ u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} u_\beta \Gamma_{\alpha\gamma}^\gamma(\partial) - \frac{1}{2} u_\alpha \Gamma_{\gamma\beta}^\gamma(\partial) \right] \delta g^{\alpha\beta} \\
&+ \left[ g_{\alpha\beta} \nabla^\gamma \delta g^{\alpha\beta} - \frac{1}{2} \nabla_\beta \delta g^{\beta\gamma} - \frac{1}{2} \nabla_\alpha \delta g^{\alpha\gamma} \right] u_\gamma \quad (25)
\end{aligned}$$

The last line can be transformed by

$$\begin{aligned}
u_\gamma g_{\alpha\beta} \nabla^\gamma \delta g^{\alpha\beta} &= \nabla_\rho \left[ u^\rho g_{\alpha\beta} \delta g^{\alpha\beta} \right] - \nabla_\rho u^\rho g_{\alpha\beta} \delta g^{\alpha\beta} \\
u_\gamma \left[ \frac{1}{2} \nabla_\beta \delta g^{\beta\gamma} + \frac{1}{2} \nabla_\alpha \delta g^{\alpha\gamma} \right] &= \frac{1}{2} \nabla_\beta \left[ u_\gamma \delta g^{\beta\gamma} \right] + \frac{1}{2} \nabla_\alpha \left[ u_\gamma \delta g^{\alpha\gamma} \right] \\
&- \frac{1}{2} \nabla_\beta u_\gamma \delta g^{\beta\gamma} - \frac{1}{2} \nabla_\alpha u_\gamma \delta g^{\alpha\gamma} \quad (26)
\end{aligned}$$

having used  $\nabla^\gamma = g^{\rho\gamma} \nabla_\rho$  and  $g^{\rho\gamma} u_\gamma = u^\rho$ . After changing indices we get

$$\begin{aligned}
\delta\mathcal{L}_T^{(1)} &= \left[ u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} (u_\beta g_{\alpha\gamma} + u_\alpha g_{\beta\gamma}) g^{\sigma\tau} \Gamma_{\sigma\tau}^\gamma(\partial) \right] \\
&+ \frac{1}{2} (\nabla_\alpha u_\beta + \nabla_\beta u_\alpha) - \nabla_\gamma u^\gamma g_{\alpha\beta} \delta g^{\alpha\beta} \\
&+ \left[ g_{\gamma\rho} g^{\sigma\tau} \Gamma_{\sigma\tau}^\rho(\partial) - \Gamma_{\gamma\rho}^\rho(\partial) \right] \delta u^\gamma \\
&+ \nabla_\gamma \left[ u^\gamma g_{\sigma\tau} \delta g^{\sigma\tau} - u_\rho \delta g^{\gamma\rho} \right] \quad (27)
\end{aligned}$$

For the remaining parts we obtain

$$\begin{aligned}
\delta\mathcal{L}_T^{(2)} &= 9 \delta \left[ g_{\gamma\rho} u^\rho u^\gamma \right] \\
&= 9 g_{\gamma\rho} \left[ \delta u^\rho u^\gamma + u^\rho \delta u^\gamma \right] - 9 u^\gamma u^\rho g_{\gamma\sigma} g_{\rho\tau} \delta g^{\sigma\tau} \\
&= 9 \left[ 2 u_\gamma \delta u^\gamma - u_\alpha u_\beta \delta g^{\alpha\beta} \right] \quad (28)
\end{aligned}$$

and

$$\delta\mathcal{K}_T^\gamma = -3\delta u^\gamma \quad (29)$$

From (23), (27), (28) and (29) we get

$$\begin{aligned} \delta\mathcal{L}_T &= \frac{1}{2} \left\{ \mathcal{R}_{\alpha\beta}(\partial) + \mu [u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2}(u_\beta g_{\alpha\gamma} + u_\alpha g_{\beta\gamma}) g^{\sigma\tau} \Gamma_{\sigma\tau}^\gamma(\partial) \right. \\ &\quad + \frac{1}{2} (\nabla_\alpha u_\beta + \nabla_\beta u_\alpha) - \nabla_\gamma u^\gamma g_{\alpha\beta} \left. \right\} \delta g^{\alpha\beta} \\ &\quad + \frac{1}{2} \mu [g_{\gamma\rho} g^{\sigma\tau} \Gamma_{\sigma\tau}^\rho(\partial) - \Gamma_{\gamma\rho}^\rho(\partial) + 9\mu u_\gamma] \delta u^\gamma \\ &\quad + \frac{1}{2} \nabla_\gamma [g_{\alpha\beta} \nabla^\gamma \delta g^{\alpha\beta} - \nabla_\beta \delta g^{\beta\gamma} \\ &\quad + \mu (u^\gamma g_{\sigma\tau} \delta g^{\sigma\tau} - u_\rho \delta g^{\gamma\rho} - 3\delta u^\gamma)] \end{aligned} \quad (30)$$

We write the vector part of the Lagrangian (6) in the form

$$\mathcal{L}_V = -\frac{1}{2} \mathcal{L}_V^{(0)} - \mathcal{L}_V^{(1)} \quad (31)$$

with

$$\mathcal{L}_V^{(0)} = \frac{1}{2} (\nabla_\alpha u_\beta - \nabla_\beta u_\alpha) (\nabla^\alpha u^\beta - \nabla^\beta u^\alpha), \quad \mathcal{L}_V^{(1)} = \frac{1}{2} \nabla_\alpha u^\alpha \nabla_\beta u^\beta \quad (32)$$

The first part can be simplified by changing indices in two terms

$$\begin{aligned} \mathcal{L}_V^{(0)} &= \frac{1}{2} [\nabla_\alpha u_\beta \nabla^\alpha u^\beta + \nabla_\beta u_\alpha \nabla^\beta u^\alpha - \nabla_\alpha u_\beta \nabla^\beta u^\alpha - \nabla_\beta u_\alpha \nabla^\alpha u^\beta] \\ &= \nabla_\alpha u_\beta (\nabla^\alpha u^\beta - \nabla^\beta u^\alpha) = \nabla^\alpha u^\beta (\nabla_\alpha u_\beta - \nabla_\beta u_\alpha) \end{aligned} \quad (33)$$

With  $u_\beta = g_{\beta\gamma} u^\gamma$  the variation gives

$$\begin{aligned} \delta\mathcal{L}_V^{(0)} &= \delta g_{\beta\gamma} \nabla_\alpha u^\gamma (\nabla^\alpha u^\beta - \nabla^\beta u^\alpha) + g_{\beta\gamma} \nabla_\alpha \delta u^\gamma (\nabla^\alpha u^\beta - \nabla^\beta u^\alpha) \\ &\quad + \nabla_\alpha u_\beta (\nabla^\alpha \delta u^\beta - \nabla^\beta \delta u^\alpha) \\ &= -\delta g^{\rho\sigma} g_{\beta\rho} g_{\sigma\gamma} \nabla_\alpha u^\gamma (\nabla^\alpha u^\beta - \nabla^\beta u^\alpha) \\ &\quad + \nabla_\alpha [\delta u^\gamma (\nabla^\alpha u_\gamma - \nabla_\gamma u^\alpha)] - \delta u^\gamma \nabla_\alpha (\nabla^\alpha u_\gamma - \nabla_\gamma u^\alpha) \\ &\quad + \nabla_\alpha u_\beta \nabla_\gamma (g^{\gamma\alpha} \delta u^\beta - g^{\gamma\beta} \delta u^\alpha) \\ &= -\nabla_\alpha u_\sigma (\nabla^\alpha u_\rho - \nabla_\rho u^\alpha) \delta g^{\rho\sigma} \\ &\quad + \nabla_\alpha [(\nabla^\alpha u_\gamma - \nabla_\gamma u^\alpha) \delta u^\gamma] - \nabla_\alpha (\nabla^\alpha u_\gamma - \nabla_\gamma u^\alpha) \delta u^\gamma \\ &\quad + \nabla_\gamma [\nabla^\gamma u_\beta \delta u^\beta - \nabla_\alpha u^\gamma \delta u^\alpha] \\ &\quad - [\nabla_\gamma \nabla^\gamma u_\beta \delta u^\beta - \nabla_\gamma \nabla_\alpha u^\gamma \delta u^\alpha] \end{aligned}$$

and after changing indices

$$\begin{aligned} \delta\mathcal{L}_V^{(0)} &= -\nabla^\gamma u_\beta (\nabla_\gamma u_\alpha - \nabla_\alpha u_\gamma) \delta g^{\alpha\beta} - 2\nabla^\rho (\nabla_\rho u_\gamma - \nabla_\gamma u_\rho) \delta u^\gamma \\ &\quad + 2\nabla_\gamma [(\nabla^\gamma u_\rho - \nabla_\rho u^\gamma) \delta u^\rho] \end{aligned} \quad (34)$$

For the second part we get

$$\begin{aligned}\delta\mathcal{L}_V^{(1)} &= \nabla_\rho u^\rho \nabla_\gamma \delta u^\gamma \\ &= \nabla_\gamma [\nabla^\rho u_\rho \delta u^\gamma] - \nabla_\gamma [\nabla^\rho u_\rho] \delta u^\gamma\end{aligned}\quad (35)$$

and together

$$\begin{aligned}\delta\mathcal{L}_V &= -\frac{1}{2}\delta\mathcal{L}_V^{(0)} - \delta\mathcal{L}_V^{(1)} \\ &= \frac{1}{4} [\nabla^\gamma u_\beta (\nabla_\gamma u_\alpha - \nabla_\alpha u_\gamma) + \nabla^\gamma u_\alpha (\nabla_\gamma u_\beta - \nabla_\beta u_\gamma)] \delta g^{\alpha\beta} \\ &\quad + \nabla^\rho \nabla_\rho u_\gamma \delta u^\gamma - \nabla_\gamma [(\nabla^\gamma u_\rho - \nabla_\rho u^\gamma) \delta u^\rho + \nabla^\rho u_\rho \delta u^\gamma]\end{aligned}\quad (36)$$

Here we have used  $\delta g^{\alpha\beta} = \delta g^{\beta\alpha}$  and changed the indices  $\alpha, \beta$  in 1/2 of its coefficient to symmetrize the expression.

Splitting the scalar part of the Lagrangian (7) in the form

$$\mathcal{L}_S = -\frac{1}{2}\mathcal{L}_S^{(0)} - \mathcal{L}_S^{(1)}\quad (37)$$

with

$$\mathcal{L}_S^{(0)} = \nabla_\alpha v \nabla^\alpha v, \quad \mathcal{L}_S^{(1)} = V(\mathcal{U}) v^2,\quad (38)$$

the variation gives

$$\begin{aligned}\delta\mathcal{L}_S^{(0)} &= \delta [g^{\alpha\beta} \nabla_\alpha v \nabla_\beta v] \\ &= \delta g^{\alpha\beta} \nabla_\alpha v \nabla_\beta v + g^{\alpha\beta} \nabla_\alpha \delta v \nabla_\beta v + g^{\alpha\beta} \nabla_\alpha v \nabla_\beta \delta v \\ &= \nabla_\alpha v \nabla_\beta v \delta g^{\alpha\beta} + 2 \nabla^\gamma v \nabla_\gamma \delta v \\ &= \nabla_\alpha v \nabla_\beta v \delta g^{\alpha\beta} + 2 \nabla_\gamma [\nabla^\gamma v \delta v] - 2 \nabla_\gamma \nabla^\gamma v \delta v\end{aligned}\quad (39)$$

and similar to  $\delta\mathcal{L}_V^{(1)}$

$$\begin{aligned}\delta\mathcal{L}_S^{(1)} &= \frac{1}{2} \delta [m^2 v^2 + \nu^2 \nabla_\gamma u^\gamma \nabla_\rho u^\rho v^2] \\ &= m^2 v \delta v + \nu^2 \nabla^\gamma u_\gamma \nabla^\rho u_\rho v \delta v + \nu^2 \nabla_\gamma [\nabla^\rho u_\rho v^2 \delta u^\gamma] \\ &\quad - \nu^2 \nabla_\gamma [\nabla^\rho u_\rho v^2] \delta u^\gamma\end{aligned}\quad (40)$$

Together we have

$$\begin{aligned}\delta\mathcal{L}_S &= -\frac{1}{2}\delta\mathcal{L}_S^{(0)} - \delta\mathcal{L}_S^{(1)} \\ &= [\nabla_\gamma \nabla^\gamma v - m^2 v - \nu^2 \nabla^\gamma u_\gamma \nabla^\rho u_\rho v] \delta v \\ &\quad + \nu^2 \nabla_\gamma [\nabla^\rho u_\rho v^2] \delta u^\gamma - \frac{1}{2} \nabla_\alpha v \nabla_\beta v \delta g^{\alpha\beta} \\ &\quad - \nabla_\gamma [\nabla^\gamma v \delta v + \nu^2 \nabla^\rho u_\rho v^2 \delta u^\gamma]\end{aligned}\quad (41)$$

Finally, the variation of the matter part should give the stress energy tensor  $\mathcal{T}$

$$\delta [\mathcal{L}_M \epsilon] = -\frac{1}{2} \kappa \mathcal{T}_{\alpha\beta} \epsilon \delta g^{\alpha\beta}\quad (42)$$

where  $\kappa = 8\pi \frac{G}{c^4}$  with the gravitational constant  $G$  and the speed of light in vacuum  $c$ .

So we obtain the functions in (19) by (21), (30), (36), (41) and (42)

$$\begin{aligned} \mathcal{M}_{\alpha\beta}^{(g)} &= \frac{1}{2} \mathcal{R}_{\alpha\beta}(\partial) - \frac{1}{2} \mathcal{L} g_{\alpha\beta} - \frac{1}{2} \kappa \mathcal{T}_{\alpha\beta} \\ &\quad + \frac{1}{2} \mu \left[ u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} (u_\beta g_{\alpha\gamma} + u_\alpha g_{\beta\gamma}) g^{\sigma\tau} \Gamma_{\sigma\tau}^\gamma(\partial) \right] \\ &\quad + \frac{1}{4} \mu (\nabla_\alpha u_\beta + \nabla_\beta u_\alpha) - \frac{1}{2} \mu \nabla_\gamma u^\gamma g_{\alpha\beta} - \frac{9}{4} \mu^2 u_\alpha u_\beta \\ &\quad + \frac{1}{4} [\nabla^\gamma u_\beta (\nabla_\gamma u_\alpha - \nabla_\alpha u_\gamma) + \nabla^\gamma u_\alpha (\nabla_\gamma u_\beta - \nabla_\beta u_\gamma)] \\ &\quad - \frac{1}{2} \nabla_\alpha v \nabla_\beta v \end{aligned} \quad (43)$$

$$\begin{aligned} \mathcal{M}_\gamma^{(u)} &= \nabla_\rho \nabla^\rho u_\gamma + \frac{9}{2} \mu^2 u_\gamma + \nu^2 \nabla_\gamma [\nabla^\rho u_\rho v^2] \\ &\quad + \frac{1}{2} \mu [g_{\gamma\rho} g^{\sigma\tau} \Gamma_{\sigma\tau}^\rho(\partial) - \Gamma_{\gamma\rho}^\rho(\partial)] \end{aligned} \quad (44)$$

$$\mathcal{M}^{(v)} = \nabla_\gamma \nabla^\gamma v - m^2 v - \nu^2 \nabla^\gamma u_\gamma \nabla^\rho u_\rho v \quad (45)$$

$$\begin{aligned} \mathcal{K}^\gamma &= \frac{1}{2} [g_{\alpha\beta} \nabla^\gamma \delta g^{\alpha\beta} - \nabla_\beta \delta g^{\beta\gamma} + \mu (u^\gamma g_{\sigma\tau} \delta g^{\sigma\tau} - u_\rho \delta g^{\gamma\rho})] \\ &\quad - (\nabla^\gamma u_\rho - \nabla_\rho u^\gamma) \delta u^\rho - (\nu^2 v^2 + 1) \nabla^\rho u_\rho \delta u^\gamma - \frac{3}{2} \mu \delta u^\gamma \\ &\quad - \nabla^\gamma v \delta v \end{aligned} \quad (46)$$

Inserting the parts of  $\mathcal{L}$  from (7), (13) - (18), (33) we get the field equations defined by (20)

$$\begin{aligned} &\mathcal{R}_{\alpha\beta}(\partial) - \frac{1}{2} R(\partial) g_{\alpha\beta} - \frac{1}{2} \mu g^{\sigma\tau} [u_\gamma \Gamma_{\sigma\tau}^\gamma(\partial) - u_\sigma \Gamma_{\tau\gamma}^\gamma(\partial)] g_{\alpha\beta} \\ &\quad + \mu [u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) - \frac{1}{2} (u_\beta g_{\alpha\gamma} + u_\alpha g_{\beta\gamma}) g^{\sigma\tau} \Gamma_{\sigma\tau}^\gamma(\partial)] \\ &\quad + \mu \left[ \frac{1}{2} (\nabla_\alpha u_\beta + \nabla_\beta u_\alpha) - \nabla_\gamma u^\gamma g_{\alpha\beta} \right] - \frac{9}{4} \mu^2 (2 u_\alpha u_\beta + u_\gamma u^\gamma g_{\alpha\beta}) \\ &\quad + \frac{1}{2} [\nabla^\gamma u_\beta (\nabla_\gamma u_\alpha - \nabla_\alpha u_\gamma) + \nabla^\gamma u_\alpha (\nabla_\gamma u_\beta - \nabla_\beta u_\gamma)] \\ &\quad + \frac{1}{2} \nabla^\gamma u^\rho (\nabla_\gamma u_\rho - \nabla_\rho u_\gamma) g_{\alpha\beta} - \nabla_\alpha v \nabla_\beta v + \frac{1}{2} \nabla_\gamma v \nabla^\gamma v g_{\alpha\beta} \\ &\quad + \frac{1}{2} m^2 v^2 g_{\alpha\beta} + \frac{1}{2} (\nu^2 v^2 + 1) \nabla^\gamma u_\gamma \nabla^\rho u_\rho g_{\alpha\beta} \\ &= \kappa \mathcal{T}_{\alpha\beta} \quad \text{for } \alpha, \beta = 0, \dots, 3, \end{aligned} \quad (47)$$

$$\begin{aligned} &\nabla_\rho \nabla^\rho u_\gamma + \frac{9}{2} \mu^2 u_\gamma + \nu^2 \nabla_\gamma [\nabla^\rho u_\rho v^2] \\ &= \frac{1}{2} \mu [\Gamma_{\gamma\rho}^\rho(\partial) - g_{\gamma\rho} g^{\sigma\tau} \Gamma_{\sigma\tau}^\rho(\partial)] \quad \text{for } \gamma = 0, \dots, 3, \end{aligned} \quad (48)$$

$$\nabla_\gamma \nabla^\gamma v - m^2 v - \nu^2 \nabla^\gamma u_\gamma \nabla^\rho u_\rho v = 0 \quad (49)$$

## 4 Relation to General Relativity, the case of flat metric

We assume now, that  $\mu$  and  $m$  are very small. For a fixed metric  $g$  we introduce an additional parameter  $\lambda$  with  $0 \leq \lambda \leq 1$  and denote by  $u_\alpha(x, \lambda)$ ,

$v(x, \lambda)$  the solutions of equations (48), (49) for  $\mu = \lambda \mu_0$ ,  $m = \lambda m_0$  with some fixed  $\mu_0, m_0$ . Omitting the part proportional to  $\mu^2$  in (48) the linearity of the equation yields the proportionality to  $\mu$  for  $u_\alpha$ . Putting this property into equation (49) we can also describe the dependence on  $\lambda$  of  $v$ :

$$u_\alpha(x, \lambda) \sim \lambda u_\alpha(x, 1), \quad v(x, \lambda) \sim v(y, 1), \quad \nabla_\beta v(x, \lambda) \sim \lambda \nabla_\beta v(y, 1) \quad (50)$$

for  $y = \lambda x$ . With these results we can write (47) in the form

$$\mathcal{R}_{\alpha\beta}(\partial) - \frac{1}{2} R(\partial) g_{\alpha\beta} - \kappa \mathcal{T}_{\alpha\beta} \sim \lambda^2 \tilde{\mathcal{F}}_{\alpha\beta}^{(g)} \quad (51)$$

where  $\tilde{\mathcal{F}}_{\alpha\beta}^{(g)}$  depends polynomially on  $\lambda$ ,  $u_\gamma(x, 1)$ ,  $v(y, 1)$  and its first derivatives. So we can expect that the corresponding metric  $g_{\alpha\beta}(x, \lambda)$  tends to the solution of General Relativity, if  $\lambda$  resp.  $\mu$  and  $m$  tend to 0.

Now, we want to investigate the system of equations for the vector and the scalar field in the case of a flat metric  $g = \eta = \text{diag}(-1, +1, +1, +1)$ . The equations (48), (49) then take the form

$$\eta_\rho \partial_\rho^2 u_\gamma + \frac{9}{2} \mu^2 u_\gamma + \nu^2 \eta_\rho \partial_\gamma [\partial_\rho u_\rho v^2] = 0 \quad \text{for } \gamma = 0, \dots, 3, \quad (52)$$

$$\eta_\gamma \partial_\gamma^2 v - m^2 v - \nu^2 \eta_\gamma \eta_\rho \partial_\gamma u_\gamma \partial_\rho u_\rho v = 0 \quad (53)$$

where for simplicity we wrote  $\eta_\gamma := \eta_{\gamma\gamma}$ .

The vector field  $\mathcal{U}$  does not satisfy the condition of a nondynamical zero component imposed in [6], i.e. the equations for  $\mathcal{U}$  should not include a term  $\partial_0 u_0$ . So we will have a closer look to this system. The equations (52) can be written in the form

$$\mathcal{P}(x, \partial) \mathcal{U} = p_{\alpha\beta}^{(2)}(x) \partial_\alpha \partial_\beta \mathcal{U} + p_\gamma^{(1)}(x) \partial_\gamma \mathcal{U} + p^{(0)}(x) \mathcal{U} = 0 \quad (54)$$

with  $4 \times 4$  matrices  $p_{\alpha\beta}^{(2)}$ ,  $p_\gamma^{(1)}$ ,  $p^{(0)}$ . The character of this system is determined by its main part

$$\mathcal{P}^{(2)}(x, \partial) = p_{\alpha\beta}^{(2)}(x) \partial_\alpha \partial_\beta \quad (55)$$

and the corresponding main symbol  $\mathcal{P}^{(2)}(x, \xi)$  with  $\xi = \xi_\alpha|_{\alpha=0}^3$ . The system is called hyperbolic, if for all  $x$  the equation

$$\det \mathcal{P}^{(2)}(x, \lambda, \hat{\xi}) = 0 \quad (56)$$

with  $\hat{\xi} = \xi_i|_{i=1}^3$  has only real solutions  $\lambda$  and strictly hyperbolic, if additionally for  $\hat{\xi} \neq 0$  all these solutions are different.

For (52) the main part is

$$\mathcal{P}^{(2)}(x, \partial) \mathcal{U} = [\eta_\rho \partial_\rho^2 u_\gamma + \nu^2 v^2 \eta_\rho \partial_\gamma \partial_\rho u_\rho]_{\gamma=0}^3 \quad (57)$$

and the corresponding main symbol

$$\mathcal{P}^{(2)}(x, \xi) = \left( \left| \hat{\xi} \right|^2 - \xi_0^2 \right) \mathcal{I} + \nu^2 v^2 \begin{bmatrix} -\xi_0^2 & \xi_0 \xi_1 & \xi_0 \xi_2 & \xi_0 \xi_3 \\ -\xi_0 \xi_1 & \xi_1^2 & \xi_1 \xi_2 & \xi_1 \xi_3 \\ -\xi_0 \xi_2 & \xi_1 \xi_2 & \xi_2^2 & \xi_2 \xi_3 \\ -\xi_0 \xi_3 & \xi_1 \xi_3 & \xi_2 \xi_3 & \xi_3^2 \end{bmatrix} \quad (58)$$

with the identical matrix  $\mathcal{I}$ . The equation

$$\det \mathcal{P}^{(2)}(x, \lambda, \hat{\xi}) = (\nu^2 v^2 + 1) \left( \left| \hat{\xi} \right|^2 - \lambda^2 \right)^4 = 0 \quad (59)$$

has only real, but multiple solutions  $\lambda = \left| \hat{\xi} \right|$  and  $\lambda = -\left| \hat{\xi} \right|$ . Hence, the system (52) is hyperbolic, but not strictly hyperbolic. For such systems there are a priori estimates showing that the solution does not grow exponential in case of bounded right hand side and initial conditions, for example [7].

Next, we want to determine the number of degrees of freedom for the system (52), (53) using the Hamilton-Dirac analysis [16]. For this purpose we have to derive the Hamiltonian from the Lagrangian, which in the flat case  $g = \eta$  is reduced to

$$\begin{aligned} \bar{\mathcal{L}} = & -\frac{1}{2} \eta_{\alpha\beta} \eta_{\gamma} [\partial_{\alpha} u_{\beta} (\partial_{\alpha} u_{\beta} - \partial_{\beta} u_{\alpha}) + \partial_{\alpha} u_{\alpha} \partial_{\beta} u_{\beta}] + \left(\frac{3}{2}\mu\right)^2 \eta_{\gamma} u_{\gamma}^2 \\ & -\frac{1}{2} [\eta_{\alpha} (\partial_{\alpha} v)^2 + (m^2 + \nu^2 \eta_{\alpha\beta} \eta_{\gamma} \partial_{\alpha} u_{\alpha} \partial_{\beta} u_{\beta}) v^2] \end{aligned} \quad (60)$$

Here, we used equations (6), (7), (9), (13) – (17) and (33). In this expression we have to separate time derivatives  $\partial_0 u_0 =: \dot{u}_0$ ,  $\partial_0 u_i =: \dot{u}_i$ ,  $i = 1, 2, 3$ ,  $\partial_0 v =: \dot{v}$  and spatial derivatives  $\partial_j u_0$ ,  $\partial_j u_i$ ,  $\partial_j v$ ,  $i, j = 1, 2, 3$  and get

$$\begin{aligned} \bar{\mathcal{L}} = & \frac{1}{2} \sum_i (\partial_i u_0 - \dot{u}_i)^2 - \frac{1}{2} \sum_{i,j} \partial_i u_j (\partial_i u_j - \partial_j u_i) \\ & -\frac{1}{2} (\dot{u}_0 - \sum_i \partial_i u_i)^2 + \left(\frac{3}{2}\mu\right)^2 (u_0^2 - \sum_i u_i^2) \\ & +\frac{1}{2} \dot{v}^2 - \frac{1}{2} \sum_i (\partial_i v)^2 - \frac{1}{2} [m^2 + \nu^2 (\dot{u}_0 - \sum_i \partial_i u_i)^2] v^2 \end{aligned} \quad (61)$$

Determining the conjugate momenta

$$\pi^0 := \frac{\partial \bar{\mathcal{L}}}{\partial \dot{u}_0} = - (1 + \nu^2 v^2) (\dot{u}_0 - \sum_i \partial_i u_i) \quad (62)$$

$$\pi^i := \frac{\partial \bar{\mathcal{L}}}{\partial \dot{u}_i} = \dot{u}_i - \partial_i u_0 \quad (63)$$

$$\pi^4 := \frac{\partial \bar{\mathcal{L}}}{\partial \dot{v}} = \dot{v} \quad (64)$$

we obtain the base Hamiltonian defined by

$$\begin{aligned}
\mathcal{H}_0 &:= \pi^0 \dot{u}_0 + \sum_i \pi^i \dot{u}_i + \pi^4 \dot{v} - \bar{\mathcal{L}} \\
&= -\frac{1}{2} \sum_i (\partial_i u_0 - \dot{u}_i) (\partial_i u_0 + \dot{u}_i) + \frac{1}{2} \sum_{i,j} \partial_i u_j (\partial_i u_j - \partial_j u_i) \\
&\quad + \left(\frac{3}{2} \mu\right)^2 (u_0^2 - \sum_i u_i^2) + \frac{1}{2} \dot{v}^2 + \frac{1}{2} \sum_i (\partial_i v)^2 + \frac{1}{2} m^2 v^2 \\
&\quad - \frac{1}{2} (1 + \nu^2 v^2) (\dot{u}_0 - \sum_i \partial_i u_i) (\dot{u}_0 + \sum_i \partial_i u_i)
\end{aligned} \tag{65}$$

The fact, that the matrix

$$\left. \frac{\partial^2 \bar{\mathcal{L}}}{\partial \dot{u}_i \partial \dot{u}_j} \right|_{i,j=0}^4 = \text{diag} \left( - (1 + \nu^2 v^2), 1, 1, 1, 1 \right) \tag{66}$$

with  $\dot{u}_4 := \dot{v}$  is invertible implies, that there are no constraints, the base Hamiltonian is already the final Hamiltonian and can be expressed by spatial derivatives of the original fields and the conjugate momenta

$$\begin{aligned}
\mathcal{H} = \mathcal{H}_0 &= -\frac{1}{2} \sum_i \pi^i (2\partial_i u_0 + \pi^i) + \frac{1}{2} \sum_{i,j} \partial_i u_j (\partial_i u_j - \partial_j u_i) \\
&\quad + \left(\frac{3}{2} \mu\right)^2 (u_0^2 - \sum_i u_i^2) + \frac{1}{2} (\pi^4)^2 + \frac{1}{2} \sum_i (\partial_i v)^2 + \frac{1}{2} m^2 v^2 \\
&\quad + \frac{1}{2} \pi^0 \left( 2 \sum_i \partial_i u_i - \frac{1}{1 + \nu^2 v^2} \pi^0 \right)
\end{aligned} \tag{67}$$

Without any constraints the number of degrees of freedom is equal to the number of field components. This property, respectively the invertibility of matrix (66), should be stable under small perturbations and hence preserved for spacetime with small warping.

## 5 The spherically symmetric case

Usually, the spherically symmetric case is described by coordinates  $(t, r, \phi, \theta)$  which are defined by the relations

$$\tilde{x}_1 = r \cos \theta \cos \phi, \quad \tilde{x}_2 = r \cos \theta \sin \phi, \quad \tilde{x}_3 = r \sin \theta \tag{68}$$

to cartesian coordinates, and a metric tensor

$$\tilde{g} = \text{diag} (g_0, g_1, r^2, r^2 \cos^2 \theta) \tag{69}$$

with functions  $g_0 = g_0(r, t)$ ,  $g_1 = g_1(r, t)$ , But in our system of equations this is not suitable, since the spherical coordinates have an intrinsic warping. The problem can be seen in equation (48). Here, the right hand side should

vanish in the flat case, but it does not in spherical coordinates. Therefore, we transform the metric (69) to Cartesian coordinates using the formula

$$g_{\alpha\beta} = \frac{\partial \tilde{x}^\mu}{\partial x^\alpha} \frac{\partial \tilde{x}^\nu}{\partial x^\beta} \tilde{g}_{\mu\nu} \quad (70)$$

and obtain

$$g_{00} = g_0, \quad g_{0i} = g_{i0} = 0, \quad g^{00} = g_0^{-1}, \quad g^{0i} = g^{i0} = 0, \quad (71)$$

$$g_{ij} = \delta_{ij} + (g_1 - 1) \frac{x_i x_j}{r^2}, \quad g^{ij} = \delta_{ij} + (g_1^{-1} - 1) \frac{x_i x_j}{r^2} \quad (72)$$

or

$$g = \begin{bmatrix} g_0 & 0 & 0 & 0 \\ 0 & (g_1 - 1) \frac{x_1^2}{r^2} + 1 & (g_1 - 1) \frac{x_1 x_2}{r^2} & (g_1 - 1) \frac{x_1 x_3}{r^2} \\ 0 & (g_1 - 1) \frac{x_1 x_2}{r^2} & (g_1 - 1) \frac{x_2^2}{r^2} + 1 & (g_1 - 1) \frac{x_2 x_3}{r^2} \\ 0 & (g_1 - 1) \frac{x_1 x_3}{r^2} & (g_1 - 1) \frac{x_2 x_3}{r^2} & (g_1 - 1) \frac{x_3^2}{r^2} + 1 \end{bmatrix} \quad (73)$$

and a similar representation for  $g^{-1}$  with  $g_0$  and  $g_1$  replaced by its inverse  $g_0^{-1}$  and  $g_1^{-1}$ .

In the following we will write as usual a prime for a differentiation with respect to  $r$  and a dot for a differentiation with respect to  $t$ , i.e. for  $a = a(r, t)$

$$\frac{\partial a}{\partial r} =: a', \quad \frac{\partial a}{\partial t} =: \dot{a}, \quad \frac{\partial^2 a}{\partial r^2} =: a'', \quad \frac{\partial^2 a}{\partial t^2} =: \ddot{a}, \quad \frac{\partial^2 a}{\partial r \partial t} =: \dot{a}'.$$

For the Christoffel symbols (1) we get

$$\Gamma_{00}^0 = \frac{1}{2} \frac{\dot{g}_0}{g_0}, \quad \Gamma_{i0}^0 = \frac{1}{2} \frac{g_0'}{g_0} \frac{x_i}{r}, \quad \Gamma_{00}^k = -\frac{1}{2} \frac{g_0'}{g_1} \frac{x_k}{r}, \quad \Gamma_{i0}^k = \frac{1}{2} \frac{\dot{g}_1}{g_1} \frac{x_i x_k}{r^2}, \quad (74)$$

$$\Gamma_{ij}^0 = -\frac{1}{2} \frac{\dot{g}_1}{g_0} \frac{x_i x_j}{r^2}, \quad \Gamma_{ij}^k = \left[ \left( \frac{r}{2} \frac{g_1'}{g_1} + \frac{1}{g_1} - 1 \right) \frac{x_i x_j}{r^2} + \delta_{ij} \left( 1 - \frac{1}{g_1} \right) \right] \frac{x_k}{r^2} \quad (75)$$

Next, we can evaluate the Ricci tensor  $\mathcal{R}_{\alpha\beta}$  (2) and the Ricci scalar  $R$  (4). For a simple representation we first define an additional scalar  $\tilde{R}$  by

$$\tilde{R} := \frac{1}{g_1} \left[ -\frac{g_0''}{g_0} + \frac{1}{2} \frac{(g_0')^2}{g_0^2} + \frac{1}{2} \frac{g_0' g_1'}{g_0 g_1} \right] + \frac{1}{g_0} \left[ -\frac{\ddot{g}_1}{g_1} + \frac{1}{2} \frac{(\dot{g}_1)^2}{g_1^2} + \frac{1}{2} \frac{\dot{g}_0 \dot{g}_1}{g_0 g_1} \right] \quad (76)$$

Then, we obtain

$$\mathcal{R}_{00} = \frac{1}{2} g_0 \tilde{R} - \frac{1}{r} \frac{g'_0}{g_1}, \quad \mathcal{R}_{0i} = \mathcal{R}_{i0} = \frac{\dot{g}_1}{g_1} \frac{x_i}{r^2}, \quad (77)$$

$$\begin{aligned} \mathcal{R}_{ij} = & \left[ \frac{1}{2} g_1 \tilde{R} + \frac{1}{r} \frac{g'_1}{g_1} \right] \frac{x_i x_j}{r^2} \\ & + \frac{1}{2r g_1} \left[ \frac{g'_1}{g_1} - \frac{g'_0}{g_0} + \frac{2}{r} (g_1 - 1) \right] \left( \delta_{ij} - \frac{x_i x_j}{r^2} \right) \end{aligned} \quad (78)$$

and

$$R = \tilde{R} + \frac{2}{r g_1} \left[ \frac{g'_1}{g_1} - \frac{g'_0}{g_0} + \frac{1}{r} (g_1 - 1) \right] \quad (79)$$

The terms  $\mathcal{R}_{\alpha\beta} - \frac{1}{2} R g_{\alpha\beta}$  with non-vanishing  $g_{\alpha\beta}$  are

$$\mathcal{R}_{00} - \frac{1}{2} R g_{00} = \frac{1}{r} \frac{g_0}{g_1} \left[ \frac{g'_1}{g_1} + \frac{1}{r} (g_1 - 1) \right] \quad (80)$$

$$\begin{aligned} \mathcal{R}_{ij} - \frac{1}{2} R g_{ij} = & \frac{1}{r} \left[ \frac{g'_0}{g_0} - \frac{1}{r} (g_1 - 1) \right] \frac{x_i x_j}{r^2} \\ & - \frac{1}{2} \left[ \tilde{R} + \frac{1}{r g_1} \left( \frac{g'_1}{g_1} - \frac{g'_0}{g_0} \right) \right] \left( \delta_{ij} - \frac{x_i x_j}{r^2} \right) \end{aligned} \quad (81)$$

In the spherically symmetric case the vector and scalar fields can be represented in the form

$$u_0 = u_o(r, t), \quad u_i = u_e(r, t) \frac{x_i}{r}, \quad v = v(r, t) \quad (82)$$

with scalar functions  $u_o$ ,  $u_e$  and  $v$ . Now, we may calculate some terms of the equations (47) - (49). Such containing no or only linear vector field components are

$$\mathcal{N}^{(1)}(\mathcal{U}) := g^{\sigma\tau} [u_\gamma \Gamma_{\sigma\tau}^\gamma(\partial) - u_\sigma \Gamma_{\tau\gamma}^\gamma(\partial)] \quad (83)$$

$$\mathcal{N}_{\alpha\beta}^{(2)}(\mathcal{U}) := u_\gamma \Gamma_{\alpha\beta}^\gamma(\partial) \quad (84)$$

$$\mathcal{N}_\alpha^{(3)} := g_{\alpha\gamma} g^{\sigma\tau} \Gamma_{\sigma\tau}^\gamma(\partial) \quad (85)$$

$$\mathcal{N}_\gamma^{(4)} := \Gamma_{\gamma\rho}^\rho(\partial) - g_{\gamma\rho} g^{\sigma\tau} \Gamma_{\sigma\tau}^\rho(\partial) \quad (86)$$

In this case we get

$$\mathcal{N}^{(1)}(\mathcal{U}) = -\frac{1}{g_0 g_1} (\dot{g}_1 u_o + g'_0 u_e) + \frac{2}{r} \left( 1 - \frac{1}{g_1} \right) u_e \quad (87)$$

$$\mathcal{N}_{00}^{(2)}(\mathcal{U}) = \frac{1}{2} \left[ \frac{\dot{g}_0}{g_0} u_o - \frac{g'_0}{g_0} u_e \right] \quad (88)$$

$$\mathcal{N}_{0i}^{(2)}(\mathcal{U}) = \mathcal{N}_{i0}^{(2)}(\mathcal{U}) = \frac{1}{2} \left[ \frac{g'_0}{g_0} u_o + \frac{\dot{g}_1}{g_1} u_e \right] \frac{x_i}{r} \quad (89)$$

$$\mathcal{N}_{ij}^{(2)}(\mathcal{U}) = \frac{1}{2} \left[ -\frac{\dot{g}_1}{g_0} u_o + \frac{g'_1}{g_1} u_e \right] \frac{x_i x_j}{r^2} \quad (90)$$

$$+ \frac{1}{r} \left( 1 - \frac{1}{g_1} \right) \left( \delta_{ij} - \frac{x_i x_j}{r^2} \right) u_e \quad (91)$$

$$\mathcal{N}_0^{(3)} = \frac{1}{2} \left( \frac{\dot{g}_0}{g_0} - \frac{\dot{g}_1}{g_1} \right) \quad (92)$$

$$\mathcal{N}_i^{(3)} = \left[ \frac{1}{2} \left( -\frac{g'_0}{g_0} + \frac{g'_1}{g_1} \right) + \frac{2}{r} (g_1 - 1) \right] \frac{x_i}{r} \quad (93)$$

$$\mathcal{N}_0^{(4)} = \frac{\dot{g}_1}{g_1}, \quad \mathcal{N}_i^{(4)} = \left[ \frac{g'_0}{g_0} - \frac{2}{r} (g_1 - 1) \right] \frac{x_i}{r} \quad (94)$$

The quadratic term is

$$\mathcal{N}^{(5)}(\mathcal{U}) := u_\gamma u^\gamma = \frac{1}{g_0} u_o^2 + \frac{1}{g_1} u_e^2 \quad (95)$$

For the terms with derivatives of first order

$$\mathcal{N}^{(6)}(\mathcal{U}, \partial\mathcal{U}) := \nabla_\gamma u^\gamma \quad (96)$$

$$\mathcal{N}_{\alpha\beta}^{(7)}(\mathcal{U}, \partial\mathcal{U}) := \nabla^\gamma u_\beta (\nabla_\gamma u_\alpha - \nabla_\alpha u_\gamma) \quad (97)$$

$$\mathcal{N}^{(8)}(\mathcal{U}, \partial\mathcal{U}) := \nabla^\gamma u^\rho (\nabla_\gamma u_\rho - \nabla_\rho u_\gamma) \quad (98)$$

$$\mathcal{N}^{(9)}(v, \partial v) := \nabla_\gamma v \nabla^\gamma v \quad (99)$$

we obtain

$$\mathcal{N}^{(6)}(\mathcal{U}, \partial\mathcal{U}) = \frac{1}{g_0} \dot{u}_o + \frac{1}{g_1} u'_e + \frac{1}{2} \frac{1}{g_0} \left( -\frac{\dot{g}_0}{g_0} + \frac{\dot{g}_1}{g_1} \right) u_o + \frac{1}{2} \frac{1}{g_1} \left( \frac{g'_0}{g_0} - \frac{g'_1}{g_1} \right) u_e \quad (100)$$

$$\mathcal{N}_{00}^{(7)}(\mathcal{U}, \partial\mathcal{U}) = \frac{1}{g_1} \left[ u'_o - \frac{1}{2} \left( \frac{g'_0}{g_0} u_o + \frac{\dot{g}_1}{g_1} u_e \right) \right] (u'_o - \dot{u}_e) \quad (101)$$

$$\mathcal{N}_{i0}^{(7)}(\mathcal{U}, \partial\mathcal{U}) = \frac{1}{g_0} \left[ -\dot{u}_o + \frac{1}{2} \left( \frac{\dot{g}_0}{g_0} u_o - \frac{g'_0}{g_1} u_e \right) \right] (u'_o - \dot{u}_e) \frac{x_i}{r} \quad (102)$$

$$\mathcal{N}_{0j}^{(7)}(\mathcal{U}, \partial\mathcal{U}) = \frac{1}{g_1} \left[ u'_e + \frac{1}{2} \left( \frac{\dot{g}_1}{g_0} u_o - \frac{g'_1}{g_1} u_e \right) \right] (u'_o - \dot{u}_e) \frac{x_j}{r} \quad (103)$$

$$\mathcal{N}_{ij}^{(7)}(\mathcal{U}, \partial\mathcal{U}) = \frac{1}{g_0} \left[ -\dot{u}_e + \frac{1}{2} \left( \frac{g'_0}{g_0} u_o + \frac{\dot{g}_1}{g_1} u_e \right) \right] (u'_o - \dot{u}_e) \frac{x_i x_j}{r^2} \quad (104)$$

$$\mathcal{N}^{(8)}(\mathcal{U}, \partial\mathcal{U}) = \frac{1}{g_0 g_1} (u'_o - \dot{u}_e)^2, \quad \mathcal{N}^{(9)}(v, \partial v) = \frac{1}{g_0} (\dot{v})^2 + \frac{1}{g_1} (v')^2 \quad (105)$$

Terms with second order derivatives of the vector or the scalar field are

$$\mathcal{N}_\gamma^{(10)}(\mathcal{U}, \partial\mathcal{U}, \partial^2\mathcal{U}) := \nabla_\rho \nabla^\rho u_\gamma, \quad \mathcal{N}^{(11)}(v, \partial v, \partial^2 v) := \nabla_\gamma \nabla^\gamma v \quad (106)$$

$$\begin{aligned} \mathcal{N}_0^{(10)}(\mathcal{U}, \partial\mathcal{U}, \partial^2\mathcal{U}) &= \frac{1}{g_0} \ddot{u}_o + \frac{1}{g_1} u''_o + \frac{1}{g_0} \left( -\frac{3}{2} \frac{\dot{g}_0}{g_0} + \frac{1}{2} \frac{\dot{g}_1}{g_1} \right) \dot{u}_o + \\ &+ \frac{1}{g_1} \left( -\frac{1}{2} \frac{g'_0}{g_0} - \frac{1}{2} \frac{g'_1}{g_1} + \frac{2}{r} \right) u'_o + \\ &+ \frac{1}{g_1} \left( \frac{g'_0}{g_0} \dot{u}_e - \frac{\dot{g}_1}{g_1} u'_e \right) + \mathcal{S}_0^{(o)} u_o + \mathcal{S}_0^{(e)} u_e \end{aligned} \quad (107)$$

$$\begin{aligned} \mathcal{N}_i^{(10)}(\mathcal{U}, \partial\mathcal{U}, \partial^2\mathcal{U}) &= \left[ \frac{1}{g_0} \ddot{u}_e + \frac{1}{g_1} u''_e + \frac{1}{g_0} \left( -\frac{1}{2} \frac{\dot{g}_0}{g_0} - \frac{1}{2} \frac{\dot{g}_1}{g_1} \right) \dot{u}_e + \right. \\ &+ \frac{1}{g_1} \left( \frac{1}{2} \frac{g'_0}{g_0} - \frac{3}{2} \frac{g'_1}{g_1} + \frac{2}{r} \right) u'_e + \\ &\left. + \frac{1}{g_0} \left( -\frac{g'_0}{g_0} \dot{u}_o + \frac{\dot{g}_1}{g_1} u'_o \right) + \mathcal{S}_1^{(o)} u_o + \mathcal{S}_1^{(e)} u_e \right] \frac{x_i}{r} \end{aligned} \quad (108)$$

where

$$\begin{aligned} \mathcal{S}_0^{(o)} &= \frac{1}{g_0} \left[ -\frac{1}{2} \frac{\ddot{g}_0}{g_0} + \left( \frac{\dot{g}_0}{g_0} \right)^2 - \frac{1}{4} \frac{\dot{g}_0 \dot{g}_1}{g_0 g_1} - \frac{1}{4} \left( \frac{\dot{g}_1}{g_1} \right)^2 \right] \\ &+ \frac{1}{g_1} \left[ -\frac{1}{2} \frac{g''_0}{g_0} + \frac{1}{4} \left( \frac{g'_0}{g_0} \right)^2 + \frac{1}{4} \frac{g'_0 g'_1}{g_0 g_1} - \frac{1}{r} \frac{g'_0}{g_0} \right] \end{aligned} \quad (109)$$

$$\mathcal{S}_0^{(e)} = \frac{1}{g_1} \left[ \frac{1}{2} \frac{\dot{g}'_0}{g_0} - \frac{1}{2} \frac{\dot{g}'_1}{g_1} - \frac{1}{2} \frac{\dot{g}_0 g'_0}{g_0^2} - \frac{1}{2} \frac{g'_0 \dot{g}_1}{g_0 g_1} + \frac{\dot{g}_1 g'_1}{g_1^2} - \frac{1}{r} \frac{\dot{g}_1}{g_1} \right] \quad (110)$$

$$\mathcal{S}_1^{(o)} = \frac{1}{g_0} \left[ -\frac{1}{2} \frac{\dot{g}'_0}{g_0} + \frac{1}{2} \frac{\dot{g}'_1}{g_1} + \frac{\dot{g}_0 g'_0}{g_0^2} - \frac{1}{2} \frac{g'_0 \dot{g}_1}{g_0 g_1} - \frac{1}{2} \frac{\dot{g}_1 g'_1}{g_1^2} + \frac{1}{r} \frac{\dot{g}_1}{g_1} \right] \quad (111)$$

$$\begin{aligned} \mathcal{S}_1^{(e)} &= \frac{1}{g_1} \left[ -\frac{1}{2} \frac{g''_1}{g_1} - \frac{1}{4} \left( \frac{g'_0}{g_0} \right)^2 - \frac{1}{4} \frac{g'_0 g'_1}{g_0 g_1} + \left( \frac{g'_1}{g_1} \right)^2 - \frac{1}{r} \frac{g'_1}{g_1} - \frac{2}{r^2} \right] \\ &+ \frac{1}{g_0} \left[ -\frac{1}{2} \frac{\ddot{g}_1}{g_1} + \frac{1}{4} \left( \frac{\dot{g}_1}{g_1} \right)^2 + \frac{1}{4} \frac{\dot{g}_0 \dot{g}_1}{g_0 g_1} \right] \end{aligned} \quad (112)$$

$$\mathcal{N}^{(11)}(v, \partial v, \partial^2 v) = \frac{1}{g_0} \ddot{v} + \frac{1}{g_1} v'' + \frac{2}{r} v' \quad (113)$$

With a reduced Lagrangian, defined by

$$\begin{aligned}
\tilde{\mathcal{L}} &= \frac{1}{2} \mathcal{N}^{(8)}(\mathcal{U}, \partial\mathcal{U}) - \mu \mathcal{N}^{(6)}(\mathcal{U}, \partial\mathcal{U}) - \frac{1}{2} \mu \mathcal{N}^{(1)}(\mathcal{U}) - \frac{9}{4} \mu^2 \mathcal{N}^{(5)}(\mathcal{U}) \\
&\quad + \frac{1}{2} (\nu^2 v^2 + 1) [\mathcal{N}^{(6)}(\mathcal{U}, \partial\mathcal{U})]^2 + \frac{1}{2} \mathcal{N}^{(9)}(v, \partial v) + \frac{1}{2} m^2 v^2 \quad (114) \\
&= \frac{1}{2} \frac{1}{g_0 g_1} (u'_o - \dot{u}_e)^2 + \frac{1}{2} \left[ \frac{1}{g_0} (\dot{v})^2 + \frac{1}{g_1} (v')^2 \right] + \frac{1}{2} (\nu^2 v^2 + 1) * \\
&\quad * \left[ \frac{1}{g_0} \dot{u}_o + \frac{1}{g_1} u'_e + \frac{1}{2} \frac{1}{g_0} \left( -\frac{\dot{g}_0}{g_0} + \frac{\dot{g}_1}{g_1} \right) u_o + \frac{1}{2} \frac{1}{g_1} \left( \frac{g'_0}{g_0} - \frac{g'_1}{g_1} \right) u_e \right]^2 \\
&\quad + \mu \left[ -\frac{1}{g_0} \dot{u}_o - \frac{1}{g_1} u'_e + \frac{1}{2} \frac{\dot{g}_0}{g_0^2} u_o + \frac{1}{2} \frac{g'_1}{g_1^2} u_e - \frac{1}{r} \left( 1 - \frac{1}{g_1} \right) u_e \right] \\
&\quad - \frac{9}{4} \mu^2 \left( \frac{1}{g_0} u_o^2 + \frac{1}{g_1} u_e^2 \right) + \frac{1}{2} m^2 v^2 \quad (115)
\end{aligned}$$

we can write equation (47) in the form

$$\begin{aligned}
\mathcal{R}_{\alpha\beta}(\partial) - \frac{1}{2} R(\partial) g_{\alpha\beta} + \tilde{\mathcal{L}} g_{\alpha\beta} + \frac{1}{2} \left[ \mathcal{N}_{\alpha\beta}^{(7)}(\mathcal{U}, \partial\mathcal{U}) + \mathcal{N}_{\beta\alpha}^{(7)}(\mathcal{U}, \partial\mathcal{U}) \right] \\
- \nabla_\alpha v \nabla_\beta v + \mu \left[ \mathcal{N}_{\alpha\beta}^{(2)}(\mathcal{U}) - \frac{1}{2} \left( u_\beta \mathcal{N}_\alpha^{(3)} + u_\alpha \mathcal{N}_\beta^{(3)} \right) \right] \\
+ \frac{1}{2} \mu (\nabla_\alpha u_\beta + \nabla_\beta u_\alpha) - \frac{9}{2} \mu^2 u_\alpha u_\beta = \kappa \mathcal{T}_{\alpha\beta} \quad (116)
\end{aligned}$$

For indices  $(\alpha, \beta) = (0, 0)$  we get

$$\begin{aligned}
\frac{1}{r} \frac{g_0}{g_1} \left[ \frac{g'_1}{g_1} + \frac{1}{r} (g_1 - 1) \right] + \tilde{\mathcal{L}} g_0 + \frac{1}{g_1} \left[ u'_o - \frac{1}{2} \left( \frac{g'_0}{g_0} u_o + \frac{\dot{g}_1}{g_1} u_e \right) \right] * \\
* (u'_o - \dot{u}_e) + \mu \left[ \dot{u}_o + \frac{1}{2} \left( -\frac{\dot{g}_0}{g_0} + \frac{\dot{g}_1}{g_1} \right) u_o + \frac{1}{2} \left( -\frac{g'_0}{g_0} + \frac{g'_1}{g_1} \right) u_e \right] \\
- \frac{9}{2} \mu^2 u_o^2 - \dot{v}^2 = \kappa \mathcal{T}_{00}, \quad (117)
\end{aligned}$$

for  $(\alpha, \beta) = (i, 0)$  or  $(\alpha, \beta) = (0, i)$

$$\begin{aligned}
\left\{ \frac{1}{r} \frac{\dot{g}_1}{g_1} + \frac{1}{2} \left[ -\frac{1}{g_0} \dot{u}_o + \frac{1}{g_1} u'_e + \frac{1}{2} \frac{1}{g_0} \left( \frac{\dot{g}_0}{g_0} + \frac{\dot{g}_1}{g_1} \right) u_o - \right. \right. \\
- \frac{1}{2} \frac{1}{g_1} \left( \frac{g'_0}{g_0} + \frac{g'_1}{g_1} \right) u_e \left. \right] (u'_o - \dot{u}_e) + \frac{1}{2} \mu \left[ u'_o + \dot{u}_e + \right. \\
+ \left( \frac{1}{2} \frac{g'_0}{g_0} - \frac{1}{2} \frac{g'_1}{g_1} - \frac{2}{r} (g_1 - 1) \right) u_o + \left. \left( -\frac{1}{2} \frac{\dot{g}_0}{g_0} + \frac{\dot{g}_1}{g_1} \right) u_e \right] \\
\left. - \frac{9}{2} \mu^2 u_o u_e - \dot{v} v' \right\} \frac{x_i}{r} = \kappa \mathcal{T}_{i0}, \quad (118)
\end{aligned}$$

and for  $(\alpha, \beta) = (i, j)$

$$\begin{aligned}
& \frac{1}{r} \left[ \frac{g'_0}{g_0} - \frac{1}{r} (g_1 - 1) \right] \frac{x_i x_j}{r^2} - \frac{1}{2} \left[ \tilde{R} + \frac{1}{r g_1} \left( \frac{g'_1}{g_1} - \frac{g'_0}{g_0} \right) \right] \left( \delta_{ij} - \frac{x_i x_j}{r^2} \right) \\
& + \tilde{\mathcal{L}} \left[ \delta_{ij} + (g_1 - 1) \frac{x_i x_j}{r^2} \right] + \frac{1}{g_0} \left[ -\dot{u}_e + \frac{1}{2} \left( \frac{g'_0}{g_0} u_o + \frac{\dot{g}_1}{g_1} u_e \right) \right] * \\
& * (u'_o - \dot{u}_e) \frac{x_i x_j}{r^2} + \mu \left[ u'_e + \left( \frac{1}{2} \frac{g'_0}{g_0} - \frac{1}{2} \frac{g'_1}{g_1} - \frac{2}{r} (g_1 - 1) \right) u_e \right] \frac{x_i x_j}{r^2} \\
& + \mu \frac{2}{r} u_e \left( \delta_{ij} - \frac{x_i x_j}{r^2} \right) - \frac{9}{2} \mu^2 u_e^2 \frac{x_i x_j}{r^2} - (v')^2 \frac{x_i x_j}{r^2} = \kappa \mathcal{T}_{ij} \quad (119)
\end{aligned}$$

For equation (48) in the case  $\gamma = 0$  we obtain

$$\begin{aligned}
& \frac{1}{g_0} \ddot{u}_o + \frac{1}{g_1} u''_o + \frac{1}{g_0} \left( -\frac{3}{2} \frac{\dot{g}_0}{g_0} + \frac{1}{2} \frac{\dot{g}_1}{g_1} \right) \dot{u}_o + \frac{1}{g_1} \left( -\frac{1}{2} \frac{g'_0}{g_0} - \frac{1}{2} \frac{g'_1}{g_1} + \frac{2}{r} \right) u'_o \\
& + \frac{1}{g_1} \left( \frac{g'_0}{g_0} \dot{u}_e - \frac{\dot{g}_1}{g_1} u'_e \right) + \mathcal{S}_0^{(o)} u_o + \mathcal{S}_0^{(e)} u_e + \frac{9}{2} \mu^2 u_o + \nu^2 * \\
& * \frac{\partial}{\partial t} \left\{ \left[ \frac{1}{g_0} \dot{u}_o + \frac{1}{g_1} u'_e + \frac{1}{2} \frac{1}{g_0} \left( -\frac{\dot{g}_0}{g_0} + \frac{\dot{g}_1}{g_1} \right) u_o + \frac{1}{2} \frac{1}{g_1} \left( \frac{g'_0}{g_0} - \frac{g'_1}{g_1} \right) u_e \right] v^2 \right\} \\
& = \frac{1}{2} \mu \frac{\dot{g}_1}{g_1} \quad (120)
\end{aligned}$$

and for  $\gamma = i$

$$\begin{aligned}
& \frac{1}{g_0} \ddot{u}_e + \frac{1}{g_1} u''_e - \frac{1}{g_0} \left( \frac{1}{2} \frac{\dot{g}_0}{g_0} + \frac{1}{2} \frac{\dot{g}_1}{g_1} \right) \dot{u}_e + \frac{1}{g_1} \left( \frac{1}{2} \frac{g'_0}{g_0} - \frac{3}{2} \frac{g'_1}{g_1} + \frac{2}{r} \right) u'_e \\
& + \frac{1}{g_0} \left( -\frac{g'_0}{g_0} \dot{u}_o + \frac{\dot{g}_1}{g_1} u'_o \right) + \mathcal{S}_1^{(o)} u_o + \mathcal{S}_1^{(e)} u_e + \frac{9}{2} \mu^2 u_e + \nu^2 * \\
& * \frac{\partial}{\partial r} \left\{ \left[ \frac{1}{g_0} \dot{u}_o + \frac{1}{g_1} u'_e + \frac{1}{2} \frac{1}{g_0} \left( -\frac{\dot{g}_0}{g_0} + \frac{\dot{g}_1}{g_1} \right) u_o + \frac{1}{2} \frac{1}{g_1} \left( \frac{g'_0}{g_0} - \frac{g'_1}{g_1} \right) u_e \right] v^2 \right\} \\
& = \frac{1}{2} \mu \left[ \frac{g'_0}{g_0} - \frac{2}{r} (g_1 - 1) \right] \quad (121)
\end{aligned}$$

where we omitted the factor  $\frac{x_i}{r}$  on both sides. Finally, equation (49) gives

$$\begin{aligned}
& \frac{1}{g_0} \ddot{v} + \frac{1}{g_1} v'' + \frac{2}{r} v' - m^2 v \\
& - \nu^2 \left[ \frac{1}{g_0} \dot{u}_o + \frac{1}{g_1} u'_e + \frac{1}{2} \frac{1}{g_0} \left( -\frac{\dot{g}_0}{g_0} + \frac{\dot{g}_1}{g_1} \right) u_o + \frac{1}{2} \frac{1}{g_1} \left( \frac{g'_0}{g_0} - \frac{g'_1}{g_1} \right) u_e \right]^2 v \\
& = 0 \quad (122)
\end{aligned}$$

Now, we consider the stationary case  $g_0 = g_0(r)$ ,  $g_1 = g_1(r)$ ,  $u_o = u_o(r)$ ,  $u_e = u_e(r)$  and  $v = v(r)$ . From (51) we obtain that for  $\lambda = 0$  resp.  $\mu = 0$  and  $m = 0$  the equations (117) - (119) pass over to general relativity, giving the well known Schwarzschild solution for a central mass  $M$ . So, as a first approximation for  $g_0, g_1$  we use the Schwarzschild metric

$$g_{0(1)} = - \left(1 - \frac{r_S}{r}\right), \quad g_{1(1)} = \left(1 - \frac{r_S}{r}\right)^{-1}, \quad \text{where } r_S = 2 \frac{MG}{c^2} \quad (123)$$

and derive the behavior of corresponding functions  $u_{o(1)}, u_{e(1)}, v_{(1)}$  from equations (120) - (122), which in the stationary case after multiplication with  $g_1$  are reduced to

$$\begin{aligned} & u_o'' + \left(-\frac{1}{2} \frac{g_0'}{g_0} - \frac{1}{2} \frac{g_1'}{g_1} + \frac{2}{r}\right) u_o' \\ & + \left[-\frac{1}{2} \frac{g_0''}{g_0} + \frac{1}{4} \left(\frac{g_0'}{g_0}\right)^2 + \frac{1}{4} \frac{g_0' g_1'}{g_0 g_1} - \frac{1}{r} \frac{g_0'}{g_0}\right] u_o + \frac{9}{2} \mu^2 g_1 u_o = 0 \end{aligned} \quad (124)$$

$$\begin{aligned} & u_e'' + \left(\frac{1}{2} \frac{g_0'}{g_0} - \frac{3}{2} \frac{g_1'}{g_1} + \frac{2}{r}\right) u_e' \\ & + \left[-\frac{1}{2} \frac{g_1''}{g_1} - \frac{1}{4} \left(\frac{g_0'}{g_0}\right)^2 - \frac{1}{4} \frac{g_0' g_1'}{g_0 g_1} + \left(\frac{g_1'}{g_1}\right)^2 - \frac{1}{r} \frac{g_1'}{g_1} - \frac{2}{r^2}\right] u_e \\ & + \frac{9}{2} \mu^2 g_1 u_e + \nu^2 g_1 \frac{\partial}{\partial r} \left\{ \frac{1}{g_1} \left[ u_e' + \frac{1}{2} \left(\frac{g_0'}{g_0} - \frac{g_1'}{g_1}\right) u_e \right] v^2 \right\} \\ & = \frac{1}{2} \mu g_1 \left[ \frac{g_0'}{g_0} - \frac{2}{r} (g_1 - 1) \right] \end{aligned} \quad (125)$$

$$v'' + \frac{2}{r} g_1 v' - m^2 g_1 v - \nu^2 \frac{1}{g_1} \left[ u_e' + \frac{1}{2} \left(\frac{g_0'}{g_0} - \frac{g_1'}{g_1}\right) u_e \right]^2 v = 0 \quad (126)$$

We see that in the system (124) - (126) the function  $u_o$  is decoupled from  $u_e$  and  $v$ . So, from the vanishing right hand side of (124) we can derive  $u_o = 0$ .

To describe functions as power series with logarithmic factors by terms of highest order we will use the following relation. For two functions  $f^{(0)}, f^{(1)}$  defined by converging series

$$\begin{aligned} f^{(i)}(r) &= \sum_{k,l \geq 0} f_{kl}^{(i)} r^{p_i - k} (\ln r)^{q_i - l} \quad i = 1, 2 \quad \text{we write} \\ f_1 &\sim f_2 \quad \text{if } p_1 = p_2, q_1 = q_2, f_{00}^{(1)} = f_{00}^{(2)} \end{aligned} \quad (127)$$

Some simple properties are

$$f_1 + f_2 \sim \begin{cases} f_{00}^{(1)} r^{p_1} (\ln r)^{q_1} & \text{for } p_1 = p_2 \text{ and } q_1 > q_2 \\ & \text{or } p_1 > p_2 \\ \left( f_{00}^{(1)} + f_{00}^{(2)} \right) r^{p_1} (\ln r)^{q_1} & \text{for } p_1 = p_2 \text{ and } q_1 = q_2 \end{cases} \quad (128)$$

$$f_1 f_2 \sim f_{00}^{(1)} f_{00}^{(2)} r^{p_1+p_2} (\ln r)^{q_1+q_2}, \quad (129)$$

and

$$\frac{d}{dr} f_i \sim \begin{cases} f_{00}^{(i)} p_i r^{p_i-1} (\ln r)^{q_i} & \text{for } p_i \neq 0 \text{ or } p_i = 0, q_i = 0 \\ f_{00}^{(i)} q_i r^{-1} (\ln r)^{q_i-1} & \text{for } p_i = 0 \text{ and } q_i \neq 0 \end{cases} \quad (130)$$

First, we observe

$$g_{0(1)} \sim -1, \quad g_{1(1)} \sim 1, \quad \frac{g'_{0(1)}}{g_{0(1)}} \sim \frac{g'_{1(1)}}{g_{1(1)}} \sim \frac{r_S}{r^2}, \quad \frac{g''_{0(1)}}{g_{0(1)}} \sim \frac{g''_{1(1)}}{g_{1(1)}} \sim -2 \frac{r_S}{r^3} \quad (131)$$

We will now assume, that  $r_S \ll r$  and  $r \ll \frac{2}{3} \frac{1}{\mu}$ , i.e.  $\frac{9}{2} \mu^2 \ll \frac{2}{r^2}$ . Furthermore, we set  $m = 0$ . Then, from (125) and (126) we get

$$u''_{e(1)} + \frac{2}{r} u'_{e(1)} - \frac{2}{r^2} u_{e(1)} + \nu^2 \frac{d}{dr} [u'_{e(1)} v_{(1)}^2] \sim -\frac{1}{2} \mu \frac{r_S}{r^2} \quad (132)$$

$$v''_{(1)} + \frac{2}{r} v'_{(1)} - \nu^2 [u'_{e(1)}]^2 v_{(1)} \sim 0 \quad (133)$$

Defining for some  $r_0 > r_S$  the variables

$$\tilde{r} := \frac{r}{r_0}, \quad \tilde{s} := \ln \left( \frac{r}{r_0} \right) \quad (134)$$

we use the Ansatz

$$u_{e(1)}(r) \sim \tilde{u}_{e(1)} \tilde{r}^{p_{1(1)}} \tilde{s}^{q_{1(1)}}, \quad v_{(1)}(r) \sim \tilde{v}_{(1)} \tilde{r}^{p_{2(1)}} \tilde{s}^{q_{2(1)}}, \quad (135)$$

To ensure that the coupling term in equation (133) has the same order of  $r$  resp.  $\tilde{r}$  as the remaining terms,  $u'_{e(1)}$  should have order  $-1$ , hence  $p_{1(1)} = 0$ . In equation (132) the coupling term should have the same order as the right hand side, i.e.  $u'_{e(1)} v_{(1)}^2$  must be of order  $-1$ , so that the derivative has order  $-2$ . Then follows  $p_{2(1)} = 0$ . Inserting now (135) with  $p_{1(1)} = p_{2(1)} = 0$  into (132) and (133) we get

$$\begin{aligned} & -\frac{q_{1(1)}}{r^2} \tilde{u}_{e(1)} \tilde{s}^{q_{1(1)}-1} + \frac{2q_{1(1)}}{r^2} \tilde{u}_{e(1)} \tilde{s}^{q_{1(1)}-1} - \frac{2}{r^2} \tilde{u}_{e(1)} \tilde{s}^{q_{1(1)}} \\ & + \nu^2 q_{1(1)} \tilde{u}_{e(1)} \tilde{v}_{(1)}^2 \frac{d}{dr} \left[ \frac{1}{r} \tilde{s}^{q_{1(1)}+2q_{2(1)}-1} \right] \sim \\ & -\frac{\tilde{u}_{e(1)}}{r^2} [2\tilde{s}^{q_{1(1)}} + \nu^2 q_{1(1)} \tilde{v}_{(1)}^2 \tilde{s}^{q_{1(1)}+2q_{2(1)}-1}] \sim -\frac{1}{2} \mu \frac{r_S}{r^2} \end{aligned} \quad (136)$$

$$\begin{aligned}
& -\frac{q_{2(1)}}{r^2} \tilde{v}_{(1)} \tilde{s}^{q_{2(1)}-1} + \frac{2q_{2(1)}}{r^2} \tilde{v}_{(1)} \tilde{s}^{q_{2(1)}-1} \\
& \quad - \nu^2 \frac{q_{1(1)}^2}{r^2} \tilde{u}_{e(1)}^2 \tilde{v}_{(1)} \tilde{s}^{2q_{1(1)}+q_{2(1)}-2} \sim \\
& \frac{\tilde{v}_{(1)}}{r^2} [q_{2(1)} \tilde{s}^{q_{2(1)}-1} - \nu^2 q_{1(1)}^2 \tilde{u}_{e(1)}^2 \tilde{s}^{2q_{1(1)}+q_{2(1)}-2}] \sim 0 \quad (137)
\end{aligned}$$

Equation (137) implies  $q_{2(1)} - 1 = 2q_{1(1)} + q_{2(1)} - 2$ , i.e.  $q_{1(1)} = \frac{1}{2}$ . Then, (136) gives  $q_{1(1)} + 2q_{2(1)} - 1 = 0$ , i.e.  $q_{2(1)} = \frac{1}{4}$ . The coefficients are obtained from

$$\nu^2 q_{1(1)} \tilde{u}_{e(1)} \tilde{v}_{(1)}^2 = \frac{1}{2} \mu r_S \quad \text{and} \quad q_{2(1)} = \nu^2 q_{1(1)}^2 \tilde{u}_{e(1)}^2$$

We get

$$\tilde{u}_{e(1)} = \frac{1}{\nu}, \quad \tilde{v}_{(1)} = \left( \frac{\mu}{\nu} r_S \right)^{\frac{1}{2}} \quad (138)$$

and

$$u_{e(1)}(r) \sim \frac{1}{\nu} \left[ \ln \left( \frac{r}{r_0} \right) \right]^{\frac{1}{2}}, \quad v_{(1)}(r) \sim \left( \frac{\mu}{\nu} r_S \right)^{\frac{1}{2}} \left[ \ln \left( \frac{r}{r_0} \right) \right]^{\frac{1}{4}} \quad (139)$$

The resulting gravity  $-\text{grad } v_{(1)}$ , generated by the scalar field, has the amount

$$\mathcal{G}_{S(1)} \sim \frac{d}{dr} v_{(1)} \sim \frac{1}{4} \left( 2 \frac{\mu}{\nu} G \right)^{\frac{1}{2}} \frac{1}{c} M^{\frac{1}{2}} \frac{1}{r} \left[ \ln \left( \frac{r}{r_0} \right) \right]^{-\frac{3}{4}} \quad (140)$$

This corresponds to the description of observed gravity  $\mathcal{G}_O$  by Newtonian gravity  $\mathcal{G}_N$  for the outer regions of galaxies in [12] and [17] by

$$\mathcal{G}_O \approx (\mathcal{G}_L \mathcal{G}_N)^{\frac{1}{2}} = (\mathcal{G}_L G)^{\frac{1}{2}} M^{\frac{1}{2}} \frac{1}{r} \quad (141)$$

up to the weak logarithmic term, when  $\mathcal{G}_N < \mathcal{G}_L \approx 10^{-10} \text{m s}^{-2}$ .

Now, we can use a Robertson series [5] to get a second approximation of the metric:

$$\begin{aligned}
g_{0(2)} &= -1 + \frac{2}{c^2} v_{(1)} = -1 + \hat{v}_{(1)} \tilde{s}^{\frac{1}{4}}, \\
g_{1(2)} &= 1 + \frac{2}{c^2} v_{(1)} = 1 + \hat{v}_{(1)} \tilde{s}^{\frac{1}{4}},
\end{aligned} \quad \text{where} \quad \hat{v}_{(1)} = \frac{2}{c^2} \tilde{v}_{(1)} \quad (142)$$

assuming

$$\hat{v}_{(1)} \tilde{s}^{\frac{1}{4}} \ll 1 \quad \text{or} \quad r \ll r_0 e^{\frac{1}{\hat{v}_{(1)}}} \quad \text{with} \quad \frac{1}{\hat{v}_{(1)}} = \frac{4c^2}{\left( \frac{\mu}{\nu} MG \right)^2} \quad (143)$$

For the derivatives of the metric elements we get

$$g'_{0(2)} \sim g'_{1(2)} \sim \frac{1}{4r} \hat{v}_{(1)} \tilde{s}^{-\frac{3}{4}}, \quad g''_{0(2)} \sim g''_{1(2)} \sim -\frac{1}{4r^2} \hat{v}_{(1)} \tilde{s}^{-\frac{3}{4}}, \quad (144)$$

while according to (143) we set

$$g_{0(2)} \approx -1, \quad g_{1(2)} \approx 1 \quad (145)$$

Again we assume  $r_S \ll r$ ,  $r \ll \frac{2}{3}\frac{1}{\mu}$  and  $m = 0$ . Then, for corresponding functions  $u_{o(2)}$ ,  $u_{e(2)}$ ,  $v_{(2)}$  equations (120) - (122) give  $u_{o(2)} = 0$  and

$$u''_{e(2)} + \frac{2}{r} u'_{e(2)} - \frac{2}{r^2} u_{e(2)} + \nu^2 \frac{d}{dr} [u'_{e(2)} v_{(2)}^2] \sim -\mu \hat{v}_{(1)}^2 \frac{1}{r} \tilde{s}^{\frac{1}{2}}, \quad (146)$$

$$v''_{(2)} + \frac{2}{r} v'_{(2)} - \nu^2 [u'_{e(2)}]^2 v_{(2)} \sim 0 \quad (147)$$

For the Ansatz

$$u_{e(2)}(r) \sim \tilde{u}_{e(2)} \tilde{r}^{p_{1(2)}} \tilde{s}^{q_{1(2)}}, \quad v_{(2)}(r) \sim \tilde{v}_{(2)} \tilde{r}^{p_{2(2)}} \tilde{s}^{q_{2(2)}}, \quad (148)$$

we get again  $p_{1(2)} = 0$  from equation (147) and for the coupling term in (146) in comparison with the right hand side  $p_{1(2)} + 2p_{2(2)} - 2 = -1$ , i.e.  $p_{2(2)} = \frac{1}{2}$ . With these first results we insert (148) into (146) and (147) and get

$$\begin{aligned} & \frac{\tilde{u}_{e(2)}}{r^2} [q_{1(2)} \tilde{s}^{q_{1(2)}-1} + 2q_{1(2)} \tilde{s}^{q_{1(2)}-1} - 2 \tilde{s}^{q_{1(2)}}] \\ & + \frac{\nu^2}{r_0} q_{1(2)} \tilde{u}_{e(2)} \tilde{v}_{(2)}^2 \frac{d}{dr} [\tilde{s}^{q_{1(2)}+2q_{2(2)}-1}] \sim \\ & \frac{\nu^2}{r_0} q_{1(2)} (q_{1(2)} + 2q_{2(2)} - 1) \tilde{u}_{e(2)} \tilde{v}_{(2)}^2 \frac{1}{r} \tilde{s}^{q_{1(2)}+2q_{2(2)}-2} \sim -\mu \hat{v}_{(1)}^2 \frac{1}{r} \tilde{s}^{\frac{1}{2}} \end{aligned} \quad (149)$$

$$\begin{aligned} & -\frac{1}{4} \tilde{v}_{(2)} r^{-\frac{3}{2}} \tilde{s}^{q_{2(2)}} + \tilde{v}_{(2)} r^{-\frac{3}{2}} \tilde{s}^{q_{2(2)}} \\ & -\nu^2 q_{1(2)}^2 \tilde{u}_{e(2)}^2 \tilde{v}_{(2)} r^{-\frac{3}{2}} \tilde{s}^{2q_{1(2)}+q_{2(2)}-2} \sim \\ & \tilde{v}_{(2)} r^{-\frac{3}{2}} \left[ \frac{3}{4} \tilde{s}^{q_{2(2)}} - \nu^2 q_{1(2)}^2 \tilde{u}_{e(2)}^2 \tilde{s}^{2q_{1(2)}+q_{2(2)}-2} \right] \sim 0 \end{aligned} \quad (150)$$

Now, equation (150) implies  $q_{2(2)} = 2q_{1(2)} + q_{2(2)} - 2$ , i.e.  $q_{1(2)} = 1$ . Then, (149) gives  $q_{1(2)} + 2q_{2(2)} - 2 = \frac{1}{2}$ , i.e.  $q_{2(2)} = \frac{3}{4}$ . The coefficients are obtained from

$$\frac{\nu^2}{r_0} q_{1(2)} (q_{1(2)} + 2q_{2(2)} - 1) \tilde{u}_{e(2)} \tilde{v}_{(2)}^2 = -\mu \hat{v}_{(1)}^2 \quad \text{and} \quad \frac{3}{4} = \nu^2 q_{1(2)}^2 \tilde{u}_{e(2)}^2$$

We get

$$\tilde{u}_{e(2)} = -\frac{3^{\frac{1}{2}}}{2} \frac{1}{\nu}, \quad \tilde{v}_{(2)} = \frac{2}{3^{\frac{3}{4}}} \left(\frac{\mu}{\nu}\right)^{\frac{1}{2}} r_0^{\frac{1}{2}} \hat{v}_{(1)} = \frac{4}{3^{\frac{3}{4}}} \frac{\mu}{\nu} \frac{1}{c^2} r_0^{\frac{1}{2}} r_S^{\frac{1}{2}} \quad (151)$$

and

$$u_{e(2)}(r) \sim -\frac{3^{\frac{1}{2}}}{2} \frac{1}{\nu} \ln\left(\frac{r}{r_0}\right), \quad v_{(2)}(r) \sim \frac{4}{3^{\frac{3}{4}}} \frac{\mu}{\nu} \frac{1}{c^2} r_S^{\frac{1}{2}} r^{\frac{1}{2}} \left[\ln\left(\frac{r}{r_0}\right)\right]^{\frac{3}{4}} \quad (152)$$

The resulting gravity, generated by the scalar field, has the amount

$$\mathcal{G}_{S(2)} \sim \frac{d}{dr} v_{(2)} \sim \frac{2}{3^{\frac{3}{4}}} \frac{\mu}{\nu} (2G)^{\frac{1}{2}} \frac{1}{c^3} M^{\frac{1}{2}} \frac{1}{r^{\frac{1}{2}}} \left[\ln\left(\frac{r}{r_0}\right)\right]^{\frac{3}{4}} \quad (153)$$

and is stronger than  $\mathcal{G}_{S(1)}$  in the sense of a greater exponent in  $r^{-\frac{1}{2}}$  than that in  $r^{-1}$ , but with a lower coefficient. Therefore, the higher gravity would act on longer distances and could describe the cohesion of galaxy clusters, which can not be explained sufficiently by gravity with the strength of  $\mathcal{G}_{S(1)}$ .

Finally, we will consider the case of very large  $r \gg \frac{2}{3} \frac{1}{\mu}$ , so that we have to take into account the term  $\frac{9}{2} \mu^2 g_1 u_e$  in equation (125). For the searched functions, in this case denoted by  $u_{o(3)}$ ,  $u_{e(3)}$ ,  $v_{(3)}$  we get  $u_{o(3)} = 0$  and

$$u_{e(3)} \sim \frac{1}{9} \frac{1}{\mu} \left[ \frac{g'_0}{g_0} - \frac{2}{r} (g_1 - 1) \right] \quad (154)$$

Then, for the Ansatz

$$u_{e(3)}(r) \sim \tilde{u}_{e(3)} \tilde{r}^{p_1(3)} \tilde{s}^{q_1(3)}, \quad v_{(3)}(r) \sim \tilde{v}_{(3)} \tilde{r}^{p_2(3)} \tilde{s}^{q_2(3)}, \quad (155)$$

we can assume  $p_1(3) \leq -1$  and  $g_0 \approx 1$  for large  $r$ . For  $m = 0$  equation (126) implies that

$$v_{(3)}'' + \frac{2}{r} v_{(3)}' = \tilde{v}_{(3)} \frac{\tilde{r}^{p_2(3)}}{r^2} \left[ p_2(3) (p_2(3) + 1) \tilde{s}^{q_2(3)} + (2p_2(3) + 1) q_2(3) \tilde{s}^{q_2(3)-1} + q_2(3) (q_2(3) - 1) \tilde{s}^{q_2(3)-2} \right] \quad (156)$$

must have an order  $p_2(3) + 2p_1(3) - 2 \leq p_2(3) - 4$  with respect to  $r$  resp.  $\tilde{r}$ . This is satisfied only for  $q_2(3) = 0$  and  $p_2(3) = 0$  or  $p_2(3) = -1$ . For  $p_2(3) = 0$  the function  $v_{(3)}$  would be constant and the gradient gives no contribution to gravity. For  $p_2(3) = -1$  we get

$$v_{(3)}(r) \sim \tilde{v}_{(3)} \tilde{r}^{-1} \quad \text{and} \quad \mathcal{G}_{S(3)} \sim \frac{d}{dr} v_{(3)} \sim -\tilde{v}_{(3)} \frac{r_0}{r^2} \quad (157)$$

i.e. for very large distances the gravity generated by the scalar field has the order of Newtonian gravity.

## 6 Cosmology

Next, we shall describe the dynamics of a homogeneous, isotropic, flat universe. The corresponding metric should be diagonal with identical components for the three spatial dimensions, depending only on time, i.e.

$$g = \text{diag}(g_0, g_1, g_1, g_1) \quad \text{with} \quad g_0 = g_0(t), \quad g_1 = g_1(t), \quad (158)$$

The only non-vanishing Christoffel symbols (1) are

$$\Gamma_{00}^0 = \frac{1}{2} \frac{\dot{g}_0}{g_0}, \quad \Gamma_{i0}^i = \frac{1}{2} \frac{\dot{g}_1}{g_1}, \quad \Gamma_{ii}^0 = -\frac{1}{2} \frac{\dot{g}_1}{g_0} \quad (159)$$

Using again the scalar  $\tilde{R}$ , defined in (76)

$$\tilde{R} := \frac{1}{g_0} \left[ -\frac{\ddot{g}_1}{g_1} + \frac{1}{2} \frac{(\dot{g}_1)^2}{g_1^2} + \frac{1}{2} \frac{\dot{g}_0 \dot{g}_1}{g_0 g_1} \right] \quad (160)$$

for the Ricci tensor and the Ricci scalar we get

$$\begin{aligned} \mathcal{R}_{00} &= \frac{3}{2} g_0 \tilde{R}, \quad \mathcal{R}_{ii} = \frac{3}{2} g_1 \tilde{R}, \quad R = 6 \tilde{R}, \\ \mathcal{R}_{00} - \frac{1}{2} R g_{00} &= -\frac{3}{2} g_0 \tilde{R}, \quad \mathcal{R}_{ii} - \frac{1}{2} R g_{ii} = -\frac{3}{2} g_1 \tilde{R} \end{aligned} \quad (161)$$

Because of the isotropy the spatial components  $u_i$  of the vector field vanish. Hence, the additional fields are  $u_0 = u_0(t)$  and  $v = v(t)$ .

We may now calculate the terms  $\mathcal{N}^{(1)}$  -  $\mathcal{N}^{(11)}$ , defined in (83) - (86), (95) - (99) and (106).

$$\mathcal{N}^{(1)}(\mathcal{U}) = \frac{1}{g_0} \left( \frac{1}{2} \frac{\dot{g}_0}{g_0} - 3 \frac{\dot{g}_1}{g_1} \right) u_o \quad (162)$$

$$\mathcal{N}_{00}^{(2)}(\mathcal{U}) = \frac{1}{2} \frac{\dot{g}_0}{g_0} u_o, \quad \mathcal{N}_{ii}^{(2)}(\mathcal{U}) = -\frac{1}{2} \frac{\dot{g}_1}{g_0} u_o, \quad (163)$$

$$\mathcal{N}_{i0}^{(2)}(\mathcal{U}) = \mathcal{N}_{oi}^{(2)}(\mathcal{U}) = \mathcal{N}_{ij}^{(2)}(\mathcal{U}) = 0 \quad (i \neq j) \quad (164)$$

$$\mathcal{N}_0^{(3)} = \frac{1}{2} \frac{\dot{g}_0}{g_0} - \frac{3}{2} \frac{\dot{g}_1}{g_1}, \quad \mathcal{N}_i^{(3)} = 0 \quad (165)$$

$$\mathcal{N}_0^{(4)} = 3 \frac{\dot{g}_1}{g_1}, \quad \mathcal{N}_i^{(4)} = 0 \quad (166)$$

$$\mathcal{N}^{(5)}(\mathcal{U}) = \frac{1}{g_0} u_o^2 \quad (167)$$

$$\mathcal{N}^{(6)}(\mathcal{U}, \partial\mathcal{U}) = \frac{1}{g_0} \left[ \dot{u}_o + \left( -\frac{1}{2} \frac{\dot{g}_0}{g_0} + \frac{3}{2} \frac{\dot{g}_1}{g_1} \right) u_o \right] \quad (168)$$

$$\mathcal{N}_{\alpha\beta}^{(7)}(\mathcal{U}, \partial\mathcal{U}) = \mathcal{N}^{(8)}(\mathcal{U}, \partial\mathcal{U}) = 0 \quad (169)$$

$$\mathcal{N}^{(9)}(v, \partial v) = \frac{1}{g_0} \dot{v}^2 \quad (170)$$

$$\mathcal{N}_0^{(10)}(\mathcal{U}, \partial\mathcal{U}, \partial^2\mathcal{U}) = \frac{1}{g_0} \left\{ \ddot{u}_o + \left[ -\frac{3}{2} \frac{\dot{g}_0}{g_0} + \frac{3}{2} \frac{\dot{g}_1}{g_1} \right] \dot{u}_o + \right. \quad (171)$$

$$\left. + \left[ -\frac{1}{2} \frac{\ddot{g}_0}{g_0} + \left( \frac{\dot{g}_0}{g_0} \right)^2 - \frac{3}{4} \frac{\dot{g}_0 \dot{g}_1}{g_0 g_1} - \frac{3}{4} \left( \frac{\dot{g}_1}{g_1} \right)^2 \right] \right\} \quad (172)$$

$$\mathcal{N}_i^{(10)}(\mathcal{U}, \partial\mathcal{U}, \partial^2\mathcal{U}) = 0 \quad (173)$$

$$\mathcal{N}^{(11)}(v, \partial v, \partial^2 v) = \frac{1}{g_0} \ddot{v} \quad (174)$$

For the reduced Lagrangian, defined in (114) we get

$$\begin{aligned} \tilde{\mathcal{L}} = & \frac{1}{g_0} \left\{ \frac{1}{2} (\nu^2 v^2 + 1) \frac{1}{g_0} \left[ \dot{u}_o + \left( -\frac{1}{2} \frac{\dot{g}_0}{g_0} + \frac{3}{2} \frac{\dot{g}_1}{g_1} \right) u_o \right]^2 \right. \\ & \left. - \mu \left( \dot{u}_o - \frac{1}{4} \frac{\dot{g}_0}{g_0} u_o \right) - \frac{9}{4} \mu^2 u_o^2 + \frac{1}{2} \dot{v}^2 + \frac{1}{2} m^2 g_0 v^2 \right\} \quad (175) \end{aligned}$$

Then, equation (116) gives

$$\mu \left[ \dot{u}_o + \left( -\frac{1}{2} \frac{\dot{g}_0}{g_0} + \frac{3}{2} \frac{\dot{g}_1}{g_1} \right) u_o \right] - \frac{9}{2} \mu^2 u_o^2 + \frac{1}{2} \dot{v}^2 - \frac{3}{2} g_0 \tilde{R} + g_0 \tilde{\mathcal{L}} = \kappa \mathcal{T}_{00} \quad (176)$$

$$- \frac{3}{2} g_1 \tilde{R} + g_1 \tilde{\mathcal{L}} = \kappa \mathcal{T}_{ii} \quad (177)$$

for  $(\alpha, \beta) = (0, 0)$  and  $(\alpha, \beta) = (i, i)$ , while equation (48) is relevant only in the case  $\gamma = 0$

$$\begin{aligned} & \frac{1}{g_0} \ddot{u}_o + \frac{1}{g_0} \left( -\frac{3}{2} \frac{\dot{g}_0}{g_0} + \frac{3}{2} \frac{\dot{g}_1}{g_1} \right) \dot{u}_o \\ & + \frac{1}{g_0} \left[ -\frac{1}{2} \frac{\ddot{g}_0}{g_0} + \left( \frac{\dot{g}_0}{g_0} \right)^2 - \frac{3}{4} \frac{\dot{g}_0 \dot{g}_1}{g_0 g_1} - \frac{3}{4} \left( \frac{\dot{g}_1}{g_1} \right)^2 \right] u_o + \frac{9}{2} \mu^2 u_o \\ & + \nu^2 \frac{d}{dt} \left\{ \frac{1}{g_0} \left[ \dot{u}_o + \left( -\frac{1}{2} \frac{\dot{g}_0}{g_0} + \frac{3}{2} \frac{\dot{g}_1}{g_1} \right) u_o \right] v^2 \right\} = \frac{3}{2} \mu \frac{\dot{g}_1}{g_1} \quad (178) \end{aligned}$$

Equation (49) gives

$$\frac{1}{g_0} \ddot{v} - m^2 v - \nu^2 \frac{1}{g_0^2} \left[ \dot{u}_o + \left( -\frac{1}{2} \frac{\dot{g}_0}{g_0} + \frac{3}{2} \frac{\dot{g}_1}{g_1} \right) u_o \right]^2 v = 0 \quad (179)$$

We consider the universe as matter dominated, i.e.

$$\mathcal{T}_{\alpha\beta} = \rho \phi_\alpha \phi_\beta \quad \text{with} \quad \phi_\gamma \phi^\gamma = g^{\alpha\beta} \phi_\alpha \phi_\beta = c^2, \quad (180)$$

where  $\rho$  is the matter density and  $\phi_\gamma$  the four-velocity of the matter. In the isotropic case we have  $\phi_i = 0$  for  $i > 0$  and hence  $g_0^{-1} \phi_0^2 = c^2$ . So we get

$$\mathcal{T}_{\alpha\beta} = \text{diag} (c^2 g_0 \rho, 0, 0, 0) \quad (181)$$

with  $\rho = \rho(t)$ . Neglecting radiation the preservation of mass gives

$$\rho(t) [\chi_1(t)]^3 = \rho(t_0) [\chi_1(t_0)]^3 =: M_0 \quad (182)$$

for a starting point  $t_0$ , where  $\chi_1(t)$  describes the expansion of the universe and can be defined by  $\chi_1^2 = g_1$ . Now, the system (176), (177) is equivalent to

$$\mu \left[ \dot{u}_o + \left( -\frac{1}{2} \frac{\dot{g}_0}{g_0} + \frac{3}{2} \frac{\dot{g}_1}{g_1} \right) u_o \right] - \frac{9}{2} \mu^2 u_o^2 + \frac{1}{2} \dot{v}^2 = \kappa c^2 M_0 g_0 g_1^{-\frac{3}{2}} \quad (183)$$

$$g_0 \tilde{\mathcal{L}} = \frac{3}{2} g_0 \tilde{R} \quad (184)$$

To find the behavior of the unknown functions for large times we use again an Ansatz similar to (135) with the variables

$$\tilde{t} := \frac{t}{t_0}, \quad \tilde{w} := \ln \left( \frac{t}{t_0} \right) \quad (185)$$

$$\begin{aligned} g_0(t) &\sim \bar{g}_0 \tilde{t}^{p_0} \tilde{w}^{q_0}, & g_1(t) &\sim \bar{g}_1 \tilde{t}^{p_1} \tilde{w}^{q_1}, \\ u_o(t) &\sim \bar{u}_o \tilde{t}^{p_2} \tilde{w}^{q_2}, & v(t) &\sim \bar{v} \tilde{t}^{p_3} \tilde{w}^{q_3}. \end{aligned} \quad (186)$$

First, we want to evaluate the exponents  $p_i$  so, that equations (178), (179) and (183), (184) can be fulfilled. Since  $\tilde{R}$  is of order  $-2$ , from (175) and (184) we obtain  $p_2 = -1$ . Then, equation (178) multiplied by  $g_0$ , gives  $p_2 - 2 = p_2 + 2p_3 - 2 = p_0 - 1$ , hence  $p_0 = -2$  and  $p_3 = 0$ . Finally, from equation (183) we derive  $-2 = p_0 - \frac{3}{2} p_1$  and get  $p_1 = 0$ . We could easily check that for these values of the  $p_i$  in each equation all terms have the same order with respect to  $\tilde{t}$ .

Now, we insert (186) into the equations. For some terms we have to take into account lower order terms, since the highest order terms vanish. We find

$$g_0 \tilde{R} \sim q_1 \left(1 - \frac{1}{2}q_1 + \frac{1}{2}q_0\right) t^{-2} \tilde{w}^{-2} \quad (187)$$

and

$$\begin{aligned} g_0 \tilde{\mathcal{L}} \sim & t^{-2} \left[ \frac{1}{2} (\nu^2 \bar{v}^2 \tilde{w}^{2q_3} + 1) \left( q_2 - \frac{1}{2}q_0 + \frac{3}{2}q_1 \right) \bar{g}_0^{-1} \bar{u}_o^2 \tilde{w}^{-q_0+2q_2-2} \right. \\ & + \frac{1}{2} \mu \bar{u}_o t_0 \tilde{w}^{q_2} - \frac{9}{4} \mu^2 \bar{u}_o^2 t_0^2 \tilde{w}^{2q_2} \\ & \left. + \frac{1}{2} p_3^2 \bar{v}^2 \tilde{w}^{2q_3-2} + \frac{1}{2} m^2 \bar{g}_0 \bar{v}^2 t_0^2 \tilde{w}^{q_0+2q_3} \right] \end{aligned} \quad (188)$$

Equation (183) gives

$$\begin{aligned} & t^{-2} \left[ \mu \left( q_2 + \frac{3}{2}q_1 - \frac{1}{2}q_0 \right) \bar{u}_o t_0 \tilde{w}^{q_2-1} - \frac{9}{2} \mu^2 \bar{u}_o^2 t_0^2 \tilde{w}^{2q_2} \right. \\ & \left. + \frac{1}{2} p_3^2 \bar{v}^2 \tilde{w}^{2q_3-2} \right] \sim t^{-2} \kappa c^2 M_0 \bar{g}_0 \bar{g}_1^{-\frac{3}{2}} t_0^2 \tilde{w}^{q_0-\frac{3}{2}q_1} \end{aligned} \quad (189)$$

Multiplying (178) and (179) by  $g_0$  we get

$$\begin{aligned} & t^{-3} \left( 6 + \frac{9}{2} \mu^2 \bar{g}_0 t_0^2 \tilde{w}^{q_0} \right) \bar{u}_o t_0 \tilde{w}^{q_2} \\ & + t^{-2} \nu^2 \tilde{w}^{q_0} \frac{d}{dt} \left[ \left( q_2 + \frac{3}{2}q_1 - \frac{1}{2}q_0 \right) \bar{u}_o \bar{v}^2 t_0 \tilde{w}^{q_2+2q_3-q_0} \right] \\ \sim & t^{-3} \left[ \left( 6 + \frac{9}{2} \mu^2 \bar{g}_0 t_0^2 \tilde{w}^{q_0} \right) \bar{u}_o t_0 \tilde{w}^{q_2} \right. \\ & + \nu^2 \left( q_2 + \frac{3}{2}q_1 - \frac{1}{2}q_0 \right) \left( q_2 + 2q_3 - q_0 \right) \bar{u}_o \bar{v}^2 t_0 \tilde{w}^{q_2+2q_3-1} \left. \right] \\ \sim & t^{-3} \frac{3}{2} \mu q_1 \bar{g}_0 t_0^2 \tilde{w}^{q_0-1} \end{aligned} \quad (190)$$

and

$$\begin{aligned} & t^{-2} \bar{v} \tilde{w}^{q_3} \left[ q_3 \tilde{w}^{-1} + m^2 \bar{g}_0 t_0^2 \tilde{w}^{q_0} \right. \\ & \left. + \nu^2 \left( q_2 + \frac{3}{2}q_1 - \frac{1}{2}q_0 \right)^2 \bar{g}_0^{-1} \bar{u}_o^2 \tilde{w}^{2q_2-q_0-2} \right] \sim 0 \end{aligned} \quad (191)$$

Now, we search values for  $q_0, \dots, q_3$  so that the relations (184) and (189) - (191) can be solved up to terms of lower order, i.e. the greatest exponents in powers of  $\tilde{w}$  should occur at least two times while all other are smaller in each of the relations. A solution can be found only, if we set  $\bar{v} = 0$  so that (191) is satisfied automatically and  $q_3$  is irrelevant. The other exponents are  $q_0 = -1$ ,  $q_1 = \frac{4}{3}$  and  $q_2 = -2$ . Then, the values of  $\bar{g}_0$ ,  $\bar{g}_1$  and  $\bar{u}_o$  are obtained from a comparison of coefficients for the terms of highest order with respect to  $\tilde{w}$ .

$$\frac{1}{2} \mu \bar{u}_o t_0 = -\frac{1}{3}, \quad \frac{1}{2} \mu \bar{u}_o t_0 = \kappa c^2 M_0 \bar{g}_0 \bar{g}_1^{-\frac{3}{2}} t_0^2, \quad 6 \bar{u}_o t_0 = 2 \mu \bar{g}_0 t_0^2 \quad (192)$$

Finally, we get

$$\begin{aligned} g_0(t) &\sim -\frac{2}{\mu^2} \frac{1}{t^2} \left[ \ln \left( \frac{t}{t_0} \right) \right]^{-1}, & g_1(t) &\sim \left( \frac{6}{\mu^2} \kappa c^2 M_0 \right)^{\frac{2}{3}} \left[ \ln \left( \frac{t}{t_0} \right) \right]^{\frac{4}{3}}, \\ u_o(t) &\sim -\frac{2}{3\mu} \frac{1}{t} \left[ \ln \left( \frac{t}{t_0} \right) \right]^{-2}, & v &\sim 0 \end{aligned} \quad (193)$$

We may now discuss the meaning of this result for the expansion of the universe and the observable redshift in comparison with the Robertson-Walker metric using the representation of the line element

$$(ds)^2 = g_{\alpha\beta} dx^\alpha dx^\beta = \chi_1^2 (dx)^2 - \chi_0^2 (dt)^2 \quad (194)$$

In our case  $\chi_0^2(t) = -g_0(t)$  and  $\chi_1^2(t) = g_1(t)$ , while for the Robertson-Walker metric  $\chi_0^{(RW)} = 1$ .

If we consider two points  $P_1$  and  $P_2$  at a fixed time  $t$ , we have  $dt = 0$  and hence  $ds = \chi_1 dx$ . Integrating from  $P_1$  to  $P_2$  gives for the distance

$$\Delta s = \chi_1 \Delta x, \quad \frac{d}{dt} \Delta s = \dot{\chi}_1 \Delta x \quad \text{and hence} \quad \frac{\frac{d}{dt} \Delta s}{\Delta s} = \frac{\dot{\chi}_1}{\chi_1} =: H(t) \quad (195)$$

$H$  describes how fast the distance of two points increases in relation to its value. From (193) we get

$$H(t) \sim \frac{2}{3} \frac{1}{t} \left[ \ln \left( \frac{t}{t_0} \right) \right]^{-1} \quad (196)$$

What we actually observe is the redshift of light, from which we conclude to the velocity of the emitter. Since light moves with velocity  $c$ , it satisfies  $ds = 0$  and (194) gives

$$dx = \frac{\chi_0}{\chi_1} dt \quad (197)$$

Considering light sent from a point  $P_0$  with wavelength  $\lambda_0$  at time  $t_0 + \Delta t_0$  to  $P_1$  with wavelength  $\lambda_1$  at time  $t_1 + \Delta t_1$  we can assume that the distance between  $P_0$  and  $P_1$  does not change during short times  $\Delta t_0$  resp.  $\Delta t_1$ . Integration of (197) for  $\Delta t_i = 0$  and  $\Delta t_i = \lambda_i$  ( $i = 0, 1$ ) gives

$$\int_{P_0}^{P_1} dx = \int_{t_0}^{t_1} \frac{\chi_0(t)}{\chi_1(t)} dt = \int_{t_0+\lambda_0}^{t_1+\lambda_1} \frac{\chi_0(t)}{\chi_1(t)} dt \quad (198)$$

Subtracting the integral from  $t_0 + \lambda_0$  to  $t_1$  we get

$$\lambda_0 \frac{\chi_0(t_0)}{\chi_1(t_0)} \approx \int_{t_0}^{t_0+\lambda_0} \frac{\chi_0(t)}{\chi_1(t)} dt = \int_{t_1}^{t_1+\lambda_1} \frac{\chi_0(t)}{\chi_1(t)} dt \approx \lambda_1 \frac{\chi_0(t_1)}{\chi_1(t_1)} \quad (199)$$

and finally

$$\frac{\lambda_1}{\lambda_0} \approx \frac{\chi_0(t_0) \chi_1(t_1)}{\chi_1(t_0) \chi_0(t_1)} \quad (200)$$

i.e. the redshift of observed light is determined by the function

$$L(t) = \frac{\chi_1(t)}{\chi_0(t)} \sim \left( \frac{3}{2^{\frac{1}{2}}} \mu \kappa c^2 M_0 \right)^{\frac{1}{3}} t \left[ \ln \left( \frac{t}{t_0} \right) \right]^{\frac{7}{6}} \quad (201)$$

Interpreting this observation in the sense of the Robertson-Walker metric,  $L$  would correspond to  $\chi_1^{(RW)}$ . Since the second derivative of  $L$  is positive, this interpretation suggests an accelerated expansion of the universe. A time contraction in the past could also explain the observations of objects in the early universe, which seem to have developed much faster than expected.

Reminding the identification of  $t$  with  $x^0$  to get a 'real' time one has to replace  $t$  by  $\frac{1}{c}t$ .

## 7 Discussion

In the case of a central mass the investigation of the scalar field in a first approximation showed a behavior of the generated gravity for great distances from the center, which very good coincides with observations in galaxies, while a second approximation gave a qualitatively stronger gravity for larger distances, which could possibly describe galaxy clusters without the need of dark matter.

In cosmology the long term development showed a logarithmic expansion of the universe, but a time contraction in the past, which results in redshifts of observations, that can be interpreted as an accelerated expansion. Thus, no dark energy would be needed.

For the verification of the theory in galaxies and galaxy clusters and the determination of the coupling constants  $\mu$ ,  $\nu$  the system (117) - (122) with at least four unknown functions or more generally (47) - (49) has to be solved numerically and compared to observed motions. The obtained coupling constants should be sufficiently small such that in the solar system the deviations from general relativity are negligible in the sense of (51).

Since matter is coupled only to the tensor field the theory should not be falsified by observations of gravity waves [9].

Stability was examined in the flat case for the system of vector and scalar fields, which can be generalized to a small warping of spacetime. A further investigation of spherically symmetric perturbations in the sense of [19], [18] was not possible, since the simplifications derived there could not be achieved.

At least, in the two considered cases we got solutions for the highest order terms of a power series Ansatz with logarithmic terms. Therefore, we expect no exponential growth for any of the participated fields.

The results about solutions in the central symmetric case and in cosmology are only valid for sufficiently large distances from the center or a long term, respectively. So it would be important to find a description of the behavior of gravity near great masses and in the beginning of the universe. Furthermore, rotating objects should be investigated.

## 8 Acknowledgment

I thank Ronald Dittrich for inspiration and helpful discussions.

## References

- [1] J. D. Bekenstein. Relativistic gravitation theory for the modified Newtonian dynamics paradigm. *Phys. Rev. D* **70**, 083509, 2004.
- [2] J. D. Bekenstein. Relativistic gravitation theory for the MOND paradigm. arXiv: astro-ph/0403694, 2005.
- [3] C. Brans and R. H. Dicke. Machs principle and a relativistic theory of gravitation. *Phys. Rev.*, **124**:925935, 1961.
- [4] A. Guarnizo, L. Castaneda and J. M. Tejeiro. Boundary Term in Metric f(R) Gravity: Field Equations in the Metric Formalism. arXiv: 1002.0617, 2010.
- [5] T. Fließbach. *Allgemeine Relativitätstheorie*. Springer Spektrum Akademischer Verlag, 2012.
- [6] L. Heisenberg. A systematic approach to generalisations of General Relativity and their cosmological implications. arXiv: 1807.01725, 2018.
- [7] L. Hörmander. *The Analysis of Linear Partial Differential Operators II. Differential Operators with Constant Coefficients*. Springer-Verlag Berlin Heidelberg New York Tokyo, 1983.
- [8] S. Hossenfelder. A Covariant Version of Verlinde’s Emergent Gravity. arXiv: 1703.01415, 2017.
- [9] S. Boran, S. Desai, E. O. Kahya, and R. P. Woodard. GW170817 falsifies Dark Matter Emulators. arXiv: 1710.06168, 2018.

- [10] P. Kroupa. The dark matter crisis: falsification of the current standard model of cosmology. arXiv: 1204.2546, 2012.
- [11] S. S. McGaugh and F. Lelli. The Radial Acceleration Relation in Rotationally Supported Galaxies. arXiv: 1609.05917, 2016.
- [12] M. Milgrom. A modification of the Newtonian dynamics - Implications for galaxies. *Astrophysical Journal, Vol. 270*, 1983.
- [13] J. W. Moffat. Acceleration in Modified Gravity (MOG) and the Mass-Discrepancy Baryonic Relation. arXiv: 1610.06909, 2016.
- [14] J. W. Moffat and V. T. Toth. Fundamental parameter-free solutions in Modified Gravity. arXiv: 0712.1796, 2009.
- [15] J. W. Moffat and V. T. Toth. Modified Gravity: Cosmology without dark matter or Einstein's cosmological constant. arXiv: 0710.0364, 2012.
- [16] J.A. Isenberg, J.M. Nester. The effect of gravitational interaction on classical fields: A Hamilton-Dirac analysis. *Annals of Physics, Volume 107, Pages 56-81*, 1977.
- [17] F. Lelli, S. S. McGaugh, J. M. Schombert, M. S. Pawlowski. One Law To Rule Them All: The Radial Acceleration Relation of Galaxies. arXiv: 1610.08981, 2017.
- [18] M. D. Seifert. Stability of spherically symmetric solutions in modified theories of gravity. arXiv: gr-qc/0703060, 2007.
- [19] M. D. Seifert and R. M. Wald. A general variational principle for spherically symmetric perturbations in diffeomorphism covariant theories. arXiv: gr-qc/0612121, 2007.
- [20] T. Clifton, P. G. Ferreira, A. Padilla, C. Skordis. Modified Gravity and Cosmology. arXiv: 1106.2476, 2012.
- [21] R. M. Wald. *General Relativity*. University of Chicago Press, 1984.
- [22] C. Skordis, T. Zlosnik. A new relativistic theory for Modified Newtonian Dynamics. arXiv: 2007.00082, 2021.