

# TESTING THE D.M. by QUANTUM GRAVITATION THEORY BY THE MASS ASSOCIATED TO THE TURNAROUND RADIUS

**Author: Manuel Abarca Hernandez**

## ABSTRACT

The DM by Quantum Gravitation theory (DMbQG hereafter), is a novel theory developed in the two main papers [1] and [2] Abarca, M. (2024). In the present work, in the framework of DMbQG theory it will be derived the formula for the mass associated to the turnaround radius depending on the turnaround radius and it will be shown that the new expression has a deeper physical meaning regarding the same formula in the current  $\Lambda$ CDM paradigm.

The DMbQG theory states a direct relation between the virial radius and the turnaround radius as well as the virial mass and the mass associated to the turnaround radius.

The most important contribution of this work is placed in the chapter 2 where it has been demonstrated that the formula for the mass associated to the turnaround radius, using the DMbQG theory, is the same that the one in the current  $\Lambda$ CDM paradigm. However the new formula has a deeper physical meaning because its dimensionless factor is simpler regarding the one in the current  $\Lambda$ CDM paradigm. Also it is remarkable the fact that the both dimensionless factors depend on exclusively from  $\Omega_m$  or  $\Omega_{DE}$  and the both ones differ only a 5%.

The fact that in the framework of DMbQG theory has been found a better formula for the mass associated to the turnaround radius is a strong support for the DMbQG theory whose hypothesis about the origin of DM is exclusively the gravitational field.

The chapters 3, 4 and 5 are devoted to test the DMbQG theory in the clusters: Virgo, Coma and the Local Group of galaxies using recent data published about these clusters. The tests are mainly focused on the turnaround radius and its mass associated and all of them have been a complete success by the DMbQG theory.

## 1. INTRODUCTION

The Lynden-Bell (1981) formula  $M_0 = R_0^3 \cdot H^2 \frac{\pi^2}{8 \cdot G}$  gives the mass associated to the turnaround radius ( $R_0$ ) in the case of a spherical over density with the cosmological parameter  $\Lambda = 0$ .

Currently, in the framework of  $\Lambda$ CDM the previous formula becomes

$$M_0 = R_0^3 \cdot H^2 \frac{\pi^2}{8 \cdot G} / f^2(\Omega_m) \quad (1.1)$$

Where  $f(\Omega_m) = (1 - \Omega_m)^{-1} - (\Omega_m/2)(1 - \Omega_m)^{-3/2} \cdot \text{arc cosh}[\frac{2}{\Omega_m} - 1]$

The reader can consult the previous expressions in the introduction of paper [3] Karachentsev.

Assuming  $\Omega_m = 0.3$  then  $f^2(0.3) = 0.6541465$  and

$$M_0 = \frac{R_0^3 \cdot H^2}{G} \cdot 1.88597 \quad (1.2)$$

and assuming  $H = 70 \text{ km/s/Mpc}$  the dimensionless expression becomes

$$M_0 / M_{\odot} = 2.14738 \cdot 10^{12} \cdot \left(\frac{R_0}{\text{Mpc}}\right)^3 \quad (1.3)$$

where  $M_{\odot}=1.99 \cdot 10^{30}$  kg and  $\text{Mpc}=3.0857 \cdot 10^{22}$  m

At the present work it will be obtained the same formula in the framework of DMbQG so it is needed to consult the papers [1] and [2] Abarca, M. in order to understand several formulas cited in this work.

Now it will be referred the main formulas used in this work that has been demonstrated in [1] and [2] papers.

### Formulas derived from the DMbQG theory used in this work

#### 1° Zero gravity radius

It is the formula (13.8), paper [1],

$$R_{ZG} = \left[ \frac{a^2}{H^2 \cdot \Omega_{DE}} \right]^{2/5} \quad (1.4)$$

The parameter  $a^2$  is specific for each galaxy or galaxy cluster in the framework of DMbQG theory.

In the sphere with this radius, the total mass (baryonic plus DM) is counterbalanced with the DE. In the paper [4] Chernin, A. it is defined the gravitating mass as the addition of total mass plus a negative mass associated to DE so into the sphere with radius  $R_{ZG}$  the gravitating mass is zero.

#### 2° Parameter $a^2$

In the chapter 3, paper [2], it is demonstrated the formula:

$$a^2 = (G \cdot M_{200})^{5/6} \cdot (10 \cdot H)^{1/3} \quad (1.5)$$

This formula is derived from the called Direct mass applied into the sphere with radius  $R_{200}$  and mass  $M_{200}$

#### 3° Ratio zero gravity radius versus virial radius

In the chapter 4, paper [2] it was got the ratio

$$\frac{R_{ZG}}{R_{VIR}} = U^2 \text{ being } U = \left[ \frac{100}{\Omega_{DE}} \right]^{1/5} \text{ and assuming } \Omega_{DE} = 0.7 \text{ then } U \approx 2.7 \text{ and}$$

$$R_{ZG} / R_{VIR} \approx 7.277 \quad (1.6)$$

#### 4° Gravitating mass

In the chapter 4, paper [2], using the gravitating mass concept, defined by [4] Chernin, A and merging with the Direct mass formula, it is got the formula for the gravitating mass:

$$M_G(< R) = M_G(< f \cdot R_{ZG}) = \left[ \sqrt{f} - f^3 \right] \cdot U \cdot M_{200} \quad (1.7)$$

Where  $f = \text{Radius} / R_{ZG}$

For example: As at the zero velocity radius  $f = 0.602$  then

$$M_G (<R_{ZV}) \approx 1.5 \cdot M_{200} \quad (1.8)$$

5° Ratio zero velocity radius versus zero gravity radius

In the chapter 5, paper [2], was demonstrated the *zero velocity radius theorem*, that states;

$$f = R_{ZV} / R_{ZG} = 0.602 \quad (1.9)$$

6° Ratio turnaround radius versus virial radius

Merging the ratio (1.9) with the ratio (1.6) it is got

$$R_{ZV} / R_{200} = 0.602 \cdot 7.277 = 4.38 \quad (1.10)$$

In the following epigraph these previous formulas will be cited frequently. The reader may study more in deep such formulas in the papers [1] and [2], where it is developed fully the DMbQG theory.

## 2. FORMULA FOR THE GRAVITATING MASS ASSOCIATED TO THE TURNAROUND RADIUS IN THE DMbQG THEORY

In the paper [2] Abarca, M.(2024), see chapter 2, was shown that for galaxy clusters is a good approximation to consider  $M_{VIR} \approx M_{200}$  where  $M_{200} = \frac{100H^2R_{200}^3}{G}$  (2.1)

i.e. the sphere with radius  $R_{200}$  has an average density equal to 200 times the Universal density  $\rho_c = \frac{3H^2}{8\pi G}$ .

In the paper [2] Abarca, M.(2024) was demonstrated the formula

$$a^2 = (G \cdot M_{200})^{5/6} \cdot (10 \cdot H)^{1/3} \quad \text{See chapter 3.} \quad (2.2)$$

In the paper [1] Abarca, M.(2024) was demonstrated the expression for the zero gravity radius.

$$R_{ZG} = \left[ \frac{a^2}{H^2 \cdot \Omega_{DE}} \right]^{2/5} \quad \text{See chapter 13.} \quad (2.3)$$

At this radius the total mass is counterbalanced by the DE, so the called gravitating mass is zero. Do not mislead this radius with the zero velocity radius or turnaround radius.

In the paper [2] Abarca, M.(2024) was demonstrated the *Zero velocity radius theorem* (see chapter 5) that assuming  $\Omega_{DE} = 0.7$  and  $H = 70$  km/s/Mpc stated that  $R_{ZV} / R_{ZG} \approx 0.602016$ .

So calling  $f = R_{ZV} / R_{ZG} \approx 0.602$ . Then  $R_{ZV} = f \cdot R_{ZG}$  (2.4)

Merging the expression (2.3) into (2.4) it is got  $R_{ZV} = f \cdot \frac{a^{4/5}}{H^{4/5} \Omega_{DE}^{2/5}}$

Clearing up  $a^{4/5}$  from this expression and doing its cubic power it is obtained:

$$a^{12/5} = R_{ZV}^3 \cdot \frac{H^{12/5} \Omega_{DE}^{6/5}}{f^3} \quad (2.5)$$

Clearing up from (2.2) the virial mass it is got:

$$M_{200} = \frac{a^{12/5}}{G \cdot 10^{2/5} \cdot H^{2/5}} \quad (2.6)$$

Merging (2.5) into (2.6) it is got

$$M_{200} = R_{ZV}^3 \cdot \frac{H^2 \Omega_{DE}^{6/5}}{G \cdot 10^{2/5} f^3} \quad (2.7)$$

The expression (1.7) gives the gravitating mass associated to the turnaround radius i.e.

$M_G(< R_{ZV}) = M_G(< f \cdot R_{ZG}) = [\sqrt{f} - f^3] \cdot U \cdot M_{200}$  where  $U = \left[\frac{100}{\Omega_{DE}}\right]^{1/5}$  and  $f = R_{ZV} / R_{ZG} = 0.602$  by the zero velocity radius theorem, (see in the epigraph 5.4, paper [2], how the ratio  $f=0.602$  was got assuming  $\Omega_{DE} = 0.7$ ).

So merging (2.7) in the later expression it is obtained:

$$M_G(< R_{ZV}) = M_0 = \frac{H^2 \cdot R_{ZV}^3}{G} \cdot \frac{10^{2/5} \cdot \Omega_{DE}^{6/5} [\sqrt{f} - f^3]}{10^{2/5} \cdot \Omega_{DE}^{1/5} f^3} = \frac{H^2 \cdot R_{ZV}^3}{G} \cdot \frac{\Omega_{DE} [\sqrt{f} - f^3]}{f^3} \quad (2.8)$$

Assuming  $\Omega_{DE} = 0.7$ ,  $f = 0.602$  and writing  $R_{ZV}$  as  $R_0$  it is got the expression:

$$M_0 = \frac{H^2 \cdot R_0^3}{G} \cdot 1.789 \quad (2.9)$$

This formula is a new crucial test for the DMbQG theory because its dimensionless factors are extremely simples: the universal  $\Omega_{DE}$ , the  $\sqrt{f}$  that comes from the direct mass and the  $f^3$  that shows how the DE increases with the cubic power into the space. However the same formula in the framework of  $\Lambda$ CDM, see the expression (1.1), contains a quite complex function: Namely  $f(\Omega_m) = (1 - \Omega_m)^{-1} - (\Omega_m/2)(1 - \Omega_m)^{-3/2} \cdot \text{arc cosh}\left[\frac{2}{\Omega_m} - 1\right]$

It is remarkable the fact that the both dimensionless factors:

$\frac{\pi^2}{8 \cdot f^2(\Omega_m)} = 1.886$  (current  $\Lambda$ CDM paradigm) and  $\frac{\Omega_{DE} [\sqrt{f} - f^3]}{f^3} = 1.789$  (DMbQG theory) depend on exclusively from  $\Omega_m$  or  $\Omega_{DE}$  and the both ones differ only a 5%.

The formula (2.9) may be written as a dimensionless expression:

$$M_0 / M_\odot = 2.037 \cdot 10^{12} \cdot \left(\frac{R_0}{Mpc}\right)^3 \quad (2.10)$$

This expression is the same than (1.3) with the numerical factor only a 5% lower which is virtually negligible.

To demonstrate the formula (2.9) has been used all the most important formulas previously obtained in the DMbQG theory. Namely these formulas are:

1° parameter  $a^2$  depending on  $M_{200}$ . See formula (1.5)

2° Zero gravity radius depending on  $a^2$ . See formula (1.4)

3° Zero velocity theorem. See formula (1.9)

4° Gravitating mass associated to the turnaround radius. See formulas (1.7) and (1.8).

Into the DMbQG theory each galaxy or cluster has a specific parameter  $a^2$  and consequently most of the formulas contain such parameter, as it is shown in the process of the demonstration to obtain the formula (2.8) although in this formula the parameter  $a^2$  has disappeared. This formula is a step forward regarding the same formula in the current  $\Lambda$ CDM framework because the former contains simple dimensionless factors whereas the later contains quite complex dimensionless ones, and additionally the both ones differ only a 5%.

This theoretical finding is a strong support for the DMbQG theory whose hypothesis about the origin of DM is exclusively the gravitational field.

### 3. TESTING THE TURNAROUND FORMULAS IN THE VIRGO CLUSTER

The data have been taken from [5] Kashibadze, O. et al. 2020. Namely the data of table 1 has been taken from the abstract and the introduction epigraphs.

Table 1 Virgo data. Source. [5] Kashibadze et al.		
Virial mass	Turnaround radius or $R_{ZV}$	Turnaround mass or $M_0$
$(6.3 \pm 0.9) \cdot 10^{14} M_{\odot}$	$7.8 \pm 0.3$ Mpc	$(7.4 \pm 0.9) \cdot 10^{14} M_{\odot}$

Assuming that  $M_{VIR} = M_{200}$  is a good approximation, as  $M_{200} = \frac{100H^2R_{200}^3}{G}$  then knowing  $M_{200}$  it is right to get  $R_{200} = 1.77$  Mpc so by the expression (1.10) it is right to get

$$R_{ZV} = 7.75 \text{ Mpc} \quad (3.1)$$

That virtually matches with the data published by Kashibadze.

On the other side, by the expression (1.8) the mass associated to the turnaround radius is

$$M_0 = 1.5(6.3 \pm 0.9) \cdot 10^{14} M_{\odot} = (9.45 \pm 1.35) \cdot 10^{14} M_{\odot} \quad (3.2)$$

This result matches in the data range given by Kashibadze, although this result does not match so well than the turnaround radius.

A second way to calculate the turnaround mass in the framework of DMbQG theory is using the formula (2.10).

$$M_0 / M_{\odot} = 2.037 \cdot 10^{12} \cdot \left(\frac{R_0}{\text{Mpc}}\right)^3 \quad (2.10)$$

Considering  $R_{ZV} = 7.75$  Mpc calculated in the framework of DMbQG theory and using (2.10) it is got  $M_0 = 9.48 \cdot 10^{14} M_{\odot}$  (3.3)

The result (3.3) virtually matches with (3.2) because the formulas (1.8) and (2.10) are mathematically equivalents in the framework of DMbQG theory.

## 4. TESTING THE TURNAROUND FORMULAS IN THE COMA CLUSTER

### 4.1 THE SOURCE OF THE COMA CLUSTER DATA

The source of Coma data is [6] Benisty 2026. In this epigraph are summarized some important data such as virial radius and mass, the turnaround radius and its mass associated. In case the reader wants to look for these data in the original paper, in the following epigraphs each data is properly referred.

In his paper, the authors consider  $h = 0.73$  so the formulas (4.3) and (4.4) stated a lower bound turnaround radius equal to 6.7 Mpc and an upper bound one equal to 11.8 Mpc.

In the Discussion epigraph, page 26, the authors give an estimate for the virial radius equal to 2.67 Mpc

In the epigraph 4.3 Virial theorem, the authors give for the virial mass the range  $(1.45 \text{ to } 1.72) \cdot 10^{15} M_{\odot}$ . Also they refer that this value is consistent with other studies of  $M_{200}$  equal to  $1.3 \cdot 10^{15} M_{\odot}$  obtained by weak lensing method. This statement means that the authors consider the virial mass closely related to  $M_{200}$  as is usual in clusters.

In the page 20 the equation (4.6) gives the range  $(0.7 \text{ to } 3.9) \cdot 10^{15} M_{\odot}$  for the mass associated to the turnaround radius. In the table 2 are summarized the previous Coma data.

Table 2 Coma data source: [6] Benisty 2026			
Virial mass	Virial Radius	Upper bound for the Turnaround radius	Turnaround mass or $M_0$
1.45 to 1.72 $M_{\odot}$	2.67 Mpc	11.8 Mpc	$(0.7 \text{ to } 3.9) \cdot 10^{15} M_{\odot}$

The average virial mass is  $1.59 \cdot 10^{15} M_{\odot}$  and by the expression (2.1) it is right to get  $R_{200} = 2.41$  Mpc that differs a 10 % regarding the virial radius given by Benisty. As the DMbQG theory is developed using  $M_{200}$  and  $R_{200}$ , in this test will be used these ones.

### 4.2 TURNAROUND RADIUS AND MASS

By the expression (1.10) it is right to get  $R_{ZV} = 4.38 \cdot 2.41 = 10.55$  Mpc

$$R_{ZV} = 10.55 \text{ Mpc} \quad (4.1)$$

that it is 1.3 Mpc lower than the turnaround upper bound radius.

By the expression (1.8)  $M_0 = 1.5 \cdot 1.59 \cdot 10^{15} M_{\odot} = 2.38 \cdot 10^{15} M_{\odot}$

$$M_0 = 2.38 \cdot 10^{15} M_{\odot} \quad (4.2)$$

that is just the average value of the mass range  $(0.7 \text{ to } 3.9) \cdot 10^{15} M_{\odot}$  given by Benisty.

A second way to calculate the turnaround mass in the framework of DMbQG theory is using the formula (2.10).

$$M_0 / M_{\odot} = 2.037 \cdot 10^{12} \cdot \left(\frac{R_0}{\text{Mpc}}\right)^3 \quad (2.10)$$

Considering  $R_{ZV} = 10.55$  Mpc calculated in the framework of DMbQG theory and using (2.10) it is got

$$M_0 = 2.39 \cdot 10^{15} M_{\odot} \quad (4.3)$$

This result virtually matches with (4.2) because the formulas (1.8) and (2.10) are mathematically equivalents in the framework of DMbQG theory.

## 5. TESTING THE DMbQG THEORY IN THE LOCAL GROUP OF GALAXIES

In this epigraph it will be used two different data sources for the Local Group. Namely the papers are [7] Shaya, E et al. (2017) and [8] Santos-Santos et al. (2023).

### 5.1 TESTING THE TOTAL MASS OF THE LOCAL GROUP UP TO 770 KPC

In the abstract of [7] Shaya, E et al. paper, the authors affirm that the mass for the MW plus M31 is equal to  $M_{LG} = (5.15 \pm 0.35) \cdot 10^{12} M_{\odot}$

In the paper [1] Abarca, the DMbQG theory is introduced and developed fully. The DM theory is initially developed into the halo region of the galaxies and the paper ended showing a way to extend the theory to the clusters. According to this theory each cluster of galaxies has a specific parameter  $a^2$  which may be obtained by the addition of the specific parameter associated to each galaxy. For example in the table 19, the LG parameter  $a^2$  is  $4.28 \cdot 10^{21} \text{ m}^{2.5}/\text{s}^2$ , obtained by the addition of the parameters ones associated to MW, M31, M33 and LMC.

The DMbQG theory stated that the total mass is unbounded and the total mass at a specific radius is calculated by the Direct mass:  $M_{TOTAL}(r) = \frac{a^2 \cdot \sqrt{r}}{G} \quad (5.1)$

In the last column of the table 20 of paper [1] Abarca, it is calculated the total mass at different distances for the L.G. using the value for the parameter  $a^2 = 4.28 \cdot 10^{21} \text{ m}^{2.5}/\text{s}^2$ . Namely for a radius equal to 770 kpc (distance between MW and M31) the total mass is equal to  $4.97 \cdot 10^{12} M_{\odot}$  that match fully with the data provided by [7] Shaya, E.

Even at this scale the DE is not negligible regarding the DM and may be quantified around a 6% so it is not significant.

In DMbQG theory each galaxy has a specific parameter  $a^2$  and may be added in order to study the DM into the external region of the cluster, assuming spherical symmetry. It is clear that the spherical symmetry for the LG is an approximation too rough. However it will be shown in the following epigraph that the turnaround radius calculated by the theory match very well with measures published.

### 5.2 TESTING THE DMbQG THEORY BY THE L.G. TURNAROUND RADIUS

In the abstract of paper [8] Santos-Santos et al. the authors stated that the LG turnaround radius is 1.25 Mpc. In this epigraph it will be compared that measure with its theoretical value in the framework of DMbQG theory.

In the chapter 13 of paper [1] Abarca, it is obtained the formula (13.8) for the zero gravity radius  $R_{ZG}$ , a concept introduced by [4] Chernin, A., but calculated specifically in the framework of the DMbQG theory. By the formula (1.4) the  $R_{ZG}$  for the LG is 2.19 Mpc. Do not mislead zero gravity radius with turnaround radius, called zero velocity radius as well.

By the zero velocity radius theorem, formula (1.9),  $R_{ZV} = 0.6 \cdot R_{ZG}$ , so the  $R_{TA} = 0.6 \cdot 2.19 = 1.3$  Mpc that match almost mathematically with the result published by [8] Santos-Santos  $R_{TA} = 1.25$  Mpc.

Additionally, also in the abstract of the paper [8] Santos-Santos et al. the authors affirm that the pure Hubble flow (i.e. where the radial velocity becomes comparable to  $H \cdot d$ ) is not reached out to at least 3 Mpc.

This result may be explained by the concept of zero gravitating radius (2.2 Mpc for the LG), because at such distance the total mass (DM plus baryonic matter) is counterbalanced by the DE obtaining a gravitating mass equal to zero and consequently at 2.2 Mpc the galaxies cannot have an adequate Hubble flow so it is needed an extra distance to reach the Hubble flow regime.

### 5.3 TURNAROUND RADIUS AND ITS MASS VERSUS $R_{200}$ AND $M_{200}$

From the expression (2.2) it is possible to clear up  $M_{200}$

$$M_{200} = \frac{a^{12/5}}{G \cdot (10 \cdot H)^{2/5}} \quad (5.2)$$

Merging the expression (2.1) and (5.2) it is possible to clear up  $R_{200}$

$$R_{200} = \left[ \frac{a^2}{100 \cdot H^2} \right]^{2/5} \quad (5.3)$$

Assuming that for the L.G.  $a^2 = 4.28 \cdot 10^{21} \text{ m}^{2.5}/\text{s}^2$  then  $M_{200} = 3.1 \cdot 10^{12} M_{\odot}$  and  $R_{200} = 301 \text{ kpc}$ . By the ratio  $R_{ZV}/R_{200} = 4.38$ , formula (1.10), it is possible to test newly the turnaround radius,  $R_{ZV} = 301 \text{ kpc} \cdot 4.38 = 1.32 \text{ Mpc}$  and by the formula (1.8) the turnaround mass:

$$M_0 = 1.5 \cdot M_{200} = 4.65 \cdot 10^{12} M_{\odot} \quad (5.4)$$

Notice how for a radius equal to 1.32 Mpc the DE is not negligible and the gravitating mass associated to  $R_{ZV}$  is lower than the total mass estimated for the LG when it is studied the gravitational interaction MW+LMC between M31+M33 where the total mass is equal to  $4.97 \cdot 10^{12} M_{\odot}$ .

The second method to calculate the turnaround mass is the formula (2.10) that for  $R = 1.32 \text{ Mpc}$  is  $M_0 = 4.68 \cdot 10^{12} M_{\odot}$  (5.5)

The results (5.4) and (5.5) are virtually equals because the formulas (1.8) and (2.10) are mathematically equivalents in the framework of DMbQG theory.

## 6. CONCLUDING REMARKS

This work, specially the chapter 2 is a new and crucial test for the novel DMbQG theory. The hypothesis of this theory is that the DM is generated by the own gravitational field and the total mass (baryonic plus DM), under spherical symmetry, grows up with the square root of the radius, its formula is called Direct mass. The theory is fully developed in the papers [1] and [2].

In the chapter 2 is placed the most important finding of this work where it has been demonstrated that the formula for the mass associated to the turnaround radius, using the DMbQG theory, is the same that the one in the current  $\Lambda$ CDM paradigm. Its dimensionless factors differ only a 5%. However the new formula has a deeper physical meaning because its

dimensionless factor is simpler regarding the one in the other formula. Namely the factors are:  $\frac{\Omega_{DE}[\sqrt{f}-f^3]}{f^3} = 1.789$ , where  $f = R_{ZV}/R_{ZG} \approx 0.602$  in the DMbQG theory and  $\frac{\pi^2}{8 \cdot f^2(\Omega_m)} = 1.886$  in the current  $\Lambda$ CDM paradigm, where the expression  $f^2(\Omega_m)$  is a quite complex function that assuming  $\Omega_m = 0.3$  then  $f^2(0.3) = 0.6541465$ .

It is remarkable the fact that the both dimensionless factors depend on exclusively from  $\Omega_m$  or  $\Omega_{DE}$  and the both ones differ only a 5%.

The fact that in the framework of DMbQG theory has been found a better formula for the mass associated to the turnaround radius is a strong support for the DMbQG theory whose hypothesis about the origin of DM is exclusively the gravitational field.

The chapters 3,4 and 5 are devoted to test the DMbQG theory in the clusters, Virgo, Coma and the Local Group of galaxies using recent data published about these clusters. The tests are mainly focused on the turnaround radius and its mass associated and all of them have been a complete success by the DMbQG theory.

## 7. BIBLIOGRAPHY

- [1] Abarca, M.(2024). *A dark matter theory by quantum gravitation for galaxies and clusters*. Journal of High Energy Physics, Gravitation and Cosmology. Vol.10 No.4  
[doi.org/10.4236/jhepgc.2024.104100](https://doi.org/10.4236/jhepgc.2024.104100)
- [2] Abarca, M.(2024). *Solving the Conundrum of Dark Matter and Dark Energy in Galaxy Clusters*. Journal of High Energy Physics, Gravitation and Cosmology. Vol.10  
[doi.org/10.4236/jhepgc.2024.104101](https://doi.org/10.4236/jhepgc.2024.104101)
- [3] Karachentsev,I.BrentTully,R. et al. 2014. *Infall of nearby galaxies into the Virgo cluster as traced with HST*. arXiv:1312.6769v2
- [4] Chernin, A.D., et al . (2013) *Dark Energy and the Structure of the Coma Cluster of Galaxies*. Astronomy & Astrophysics , 553, A101.  
[doi.org/10.1051/0004-6361/201220781](https://doi.org/10.1051/0004-6361/201220781)
- [5] Olga Kashibadze, Igor Karachentsev et al.2020. *On structure and kinematics of the Virgo cluster of galaxies*. arXiv:2002.12820v1
- [6] D. Benisty, J.Wagner, S. Haridasu, P. Salucci. 2025. *Unveiling the Coma Cluster Structure: From the Core to the Hubble Flow* arXiv: 2504.04135v3
- [7] Shaya,E. Tully,R. Hoffman,Y.Pomarede,D. 2017. *Action dynamics of the local supercluster*. arXiv:1710.08935v2
- [8] Santos-Santos,I., Navarro J., McConnachie, A.2023. *Anisotropies in the spatial distribution and kinematics of dwarf galaxies in the Local Group and beyond* arXiv:2310.02464v1