Electrodynamics in a Uniformly Accelerating Frame in Schwarzschild Spacetime

Sangwha Yi
Daejeon University, Daejeon, Republic of Korea
leeshwa@edu.dju.ac.kr

November 23, 2026

Abstract

We investigate the classical electrodynamics of a static observer—which acts as a uniformly accelerating frame—in Schwarzschild spacetime. By constructing an orthonormal tetrad basis adapted to the observer's 4-velocity, we rigorously derive the 3+1 decomposed Maxwell's equations. We demonstrate that the proper acceleration of the observer and the spatial curvature of the black hole introduce effective charge and current density terms into the field equations, fundamentally distinguishing this scenario from the flat-spacetime Rindler effect. Furthermore, we derive the covariant electromagnetic wave equation and the effective potential governing the propagation of light in this accelerating frame. The asymptotic behavior of the waves near the event horizon and at spatial infinity is analyzed. This framework provides a mathematical foundation for understanding phenomena such as the Unruh effect, Hawking radiation, and gravitational lensing in the vicinity of a black hole.

1 Introduction

The unification of general relativity and quantum mechanics remains one of the most profound challenges in modern theoretical physics. A key conceptual tool in bridging these frameworks is the study of quantum fields in curved spacetime. A fundamental insight from general relativity is the Equivalence Principle, which states that a uniformly accelerating frame in flat spacetime is locally indistinguishable from a frame at rest in a uniform gravitational field.

In flat Minkowski spacetime, a uniformly accelerating observer is described by Rindler coordinates. The quantization of fields in this frame leads to the celebrated Unruh effect, where the observer detects a thermal bath of particles with temperature $T_U = \hbar a/(2\pi c k_B)$. However, in curved spacetime, particularly in the Schwarzschild geometry describing a non-rotating black hole, a globally uniform accelerating frame does not exist due to the presence of tidal forces (spacetime curvature).

Instead, we define a "uniformly accelerating frame" as the local rest frame of a *static observer*. This observer hovers at a constant radial coordinate r and must continuously accelerate outward with a proper acceleration a to resist the black hole's gravity. In this paper, we formulate the classical electrodynamics as measured by such an observer.

The structure of the paper is as follows: In Section 2, we review the geometric setup of Schwarzschild spacetime. In Section 3, we construct the tetrad formalism for the static observer. In Section 4, we derive the modified Maxwell's equations. In Section 5, we derive the electromagnetic wave equation and analyze the effective potential. Finally, we discuss the physical implications and conclude.

2 Geometric Setup of Schwarzschild Spacetime

The Schwarzschild metric in spherical coordinates (t, r, θ, ϕ) is given by:

$$ds^2 = -f(r)c^2dt^2 + f(r)^{-1}dr^2 + r^2d\theta^2 + r^2\sin^2\theta d\phi^2, \tag{1}$$

where $f(r) = 1 - \frac{r_s}{r}$ and $r_s = \frac{2GM}{c^2}$ is the Schwarzschild radius. We adopt the signature $(-,+,+,+)$ and set $c = 1$ for the intermediate calculations, restoring it when necessary.

The non-zero Christoffel symbols $\Gamma^\mu_{\nu\lambda}$ for this metric are:

$$\Gamma^t_{tr} = \Gamma^t_{rt} = \frac{f'}{2f}, \tag{2}$$

$$\Gamma^r_{tt} = \frac{ff'}{2}, \quad \Gamma^r_{rr} = -\frac{f'}{2f}, \tag{3}$$

$$\Gamma^r_{\theta\theta} = -rf, \quad \Gamma^r_{\phi\phi} = -rf\sin^2\theta, \tag{4}$$

$$\Gamma^\theta_{r\theta} = \Gamma^\theta_{\theta r} = \frac{1}{r}, \quad \Gamma^\theta_{\phi\phi} = -\sin\theta\cos\theta, \tag{5}$$

$$\Gamma^\phi_{r\phi} = \Gamma^\phi_{\phi r} = \frac{1}{r}, \quad \Gamma^\phi_{\theta\phi} = \Gamma^\phi_{\phi\theta} = \cot\theta. \tag{6}$$

where $f' = \partial_r f = r_s/r^2$.

Because the Schwarzschild metric is a vacuum solution to the Einstein field equations, the Ricci tensor and the Ricci scalar vanish identically:

$$R_{\mu\nu} = 0, \quad R = 0. \tag{7}$$

However, the Riemann curvature tensor $R^\mu{}_{\nu\rho\sigma}$ is non-zero, indicating the presence of tidal forces.

3 The Static Observer and Tetrad Formalism

A static observer maintains a constant radial coordinate r, as well as constant θ and ϕ. The 4-velocity $u^\mu = \frac{dx^\mu}{d\tau}$ (where τ is the proper time) must satisfy $u^\mu u_\mu = -1$. For a static observer:

$$u^\mu = \left(\frac{1}{\sqrt{f}}, 0, 0, 0\right), \quad u_\mu = \left(-\sqrt{f}, 0, 0, 0\right). \tag{8}$$

The 4-acceleration is defined as $a^\mu = u^\nu\nabla_\nu u^\mu$. Since the observer is static, the only non-zero component is:

$$a^r = u^t\nabla_t u^r = u^t(\partial_t u^r + \Gamma^r_{tt}u^t) = u^t\Gamma^r_{tt}u^t = \frac{ff'}{2}. \tag{9}$$

The magnitude of the proper acceleration (the acceleration felt by the observer) is:

$$a = \sqrt{a^\mu a_\mu} = \sqrt{g_{rr}(a^r)^2} = \sqrt{f^{-1}\left(\frac{ff'}{2}\right)^2} = \frac{f'}{2\sqrt{f}} = \frac{GM}{r^2\sqrt{1-\frac{r_s}{r}}}. \tag{10}$$

Notice that as $r \to r_s$, $a \to \infty$. At spatial infinity ($r \to \infty$), $a \to 0$.

To perform local physical measurements, the observer carries an orthonormal tetrad (a local basis of four mutually orthogonal unit vectors) $e_{\hat{\mu}}$, satisfying $e_{\hat{\mu}} \cdot e_{\hat{\nu}} = \eta_{\hat{\mu}\hat{\nu}}$, where $\eta_{\hat{\mu}\hat{\nu}}$ is the Minkowski metric. A natural choice adapted to the static observer is:

$$e_{\hat{0}} = \frac{1}{\sqrt{f}}\partial_t, \tag{11}$$

$$e_{\hat{1}} = \sqrt{f}\partial_r, \tag{12}$$

$$e_{\hat{2}} = \frac{1}{r}\partial_\theta, \tag{13}$$

$$e_{\hat{3}} = \frac{1}{r\sin\theta}\partial_\phi. \tag{14}$$

The physical electric field $\mathbf{E}$ and magnetic field $\mathbf{B}$ measured by the observer are defined via the electromagnetic field tensor $F_{\mu\nu}$:

$$E_{\hat{i}} = F_{\hat{i}\hat{0}}, \quad B_{\hat{i}} = \frac{1}{2}\epsilon_{\hat{i}\hat{j}\hat{k}}F_{\hat{j}\hat{k}}, \tag{15}$$

where $\epsilon_{\hat{i}\hat{j}\hat{k}}$ is the 3D Levi-Civita symbol.

4 Maxwell's Equations in the Accelerating Frame

The general covariant Maxwell equations are:

$$\nabla_\mu F^{\mu\nu} = \mu_0 J^\nu, \tag{16}$$

$$\nabla_{[\lambda}F_{\mu\nu]} = 0, \tag{17}$$

where $J^\nu = (\rho c, \mathbf{J})$ is the 4-current density. To project these equations onto the observer's local frame, we use the 3+1 decomposition (ADM formalism).

By substituting the tetrad components and the Christoffel symbols into the covariant derivative, we obtain the following 3D vector equations as measured by the static observer.

4.1 Gauss's Law

The time component of the source equation yields Gauss's law:

$$\nabla \cdot \mathbf{E} + \frac{2}{r}E_{\hat{r}} + \frac{a}{c^2}E_{\hat{r}} = \frac{\rho}{\varepsilon_0}. \tag{18}$$

Here, $\nabla \cdot \mathbf{E} = \frac{1}{r^2}\partial_r(r^2 E_{\hat{r}}) + \dots$ is the standard flat-space divergence. The term $2/r$ arises from the spatial curvature (the spherical geometry), while the term a/c^2 represents an *effective charge density* induced by the observer's proper acceleration.

4.2 Ampere-Maxwell Law

The spatial components of the source equation yield the Ampere-Maxwell law:

$$\nabla \times \mathbf{B} - \frac{1}{c^2}\frac{\partial \mathbf{E}}{\partial \tau} + \frac{a}{c^2}(\hat{r} \times \mathbf{B}) = \mu_0 \mathbf{J}. \tag{19}$$

The term $\frac{a}{c^2}(\hat{r} \times \mathbf{B})$ acts as an *effective current density* generated by the acceleration.

4.3 Gauss's Law for Magnetism

The time component of the homogeneous equation gives:

$$\nabla \cdot \mathbf{B} + \frac{2}{r}B_{\hat{r}} + \frac{a}{c^2}B_{\hat{r}} = 0. \tag{20}$$

4.4 Faraday's Law

The spatial components of the homogeneous equation yield:

$$\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial \tau} = 0. \tag{21}$$

Notice that Faraday's law remains completely unchanged by acceleration or spatial curvature, reflecting its topological nature.

5 The Electromagnetic Wave Equation

In vacuum ($J^\mu = 0$), we introduce the electromagnetic 4-potential $A^\mu = (\Phi/c, \mathbf{A})$ such that $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. We impose the Lorenz gauge condition in curved spacetime:

$$\nabla_\mu A^\mu = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} A^\mu) = 0. \tag{22}$$

With this gauge choice, Maxwell's equations reduce to the covariant wave equation:

$$\Box A^\mu - R^\mu{}_\nu A^\nu = \mu_0 J^\mu, \tag{23}$$

where $\Box = \nabla_\mu \nabla^\mu$ is the d'Alembertian operator. Since the Schwarzschild spacetime is Ricci-flat ($R^\mu{}_\nu = 0$), the curvature coupling term vanishes, yielding:

$$\Box A^\mu = 0. \tag{24}$$

Expanding the d'Alembertian in the Schwarzschild metric yields:

$$-\frac{1}{fc^2}\frac{\partial^2 A^\mu}{\partial t^2} + \frac{1}{r^2}\frac{\partial}{\partial r}\left(r^2 f \frac{\partial A^\mu}{\partial r}\right) + \frac{1}{r^2}\nabla_\Omega^2 A^\mu = 0, \tag{25}$$

where ∇_Ω^2 is the angular momentum operator:

$$\nabla_\Omega^2 = \frac{1}{\sin\theta}\partial_\theta(\sin\theta\,\partial_\theta) + \frac{1}{\sin^2\theta}\partial_\phi^2. \tag{26}$$

To solve this equation, we employ the *tortoise coordinate* r^*, defined by:

$$\frac{dr^*}{dr} = \frac{1}{f(r)} \implies r^* = r + r_s \ln\left|\frac{r}{r_s} - 1\right|. \tag{27}$$

The tortoise coordinate maps the event horizon $r \to r_s$ to $r^* \to -\infty$, and spatial infinity $r \to \infty$ to $r^* \to \infty$.

We introduce the separation of variables ansatz:

$$A^\mu(t, r, \theta, \phi) = \sum_{l,m} \frac{u_{lm}(r^*)}{r} Y_{lm}(\theta, \phi) e^{-i\omega t}, \tag{28}$$

where Y_{lm} are the spherical harmonics. Substituting this into the wave equation yields a 1-dimensional Schrödinger-like equation:

$$\frac{d^2 u_{lm}}{dr^{*2}} + \left(\omega^2 - V_{\text{eff}}(r)\right) u_{lm} = 0, \tag{29}$$

where the effective potential $V_{\text{eff}}(r)$ is given by:

$$V_{\text{eff}}(r) = f(r)\left[\frac{l(l+1)}{r^2} + \frac{r_s}{r^3}\right]. \tag{30}$$

5.1 Analysis of the Effective Potential

The effective potential consists of two terms:

1. The **centrifugal barrier** term $\frac{l(l+1)}{r^2} f(r)$, which dominates at large r.
2. The **curvature term** $\frac{r_s}{r^3} f(r)$, which arises due to the spatial curvature of the black hole and dominates near the horizon.

At the event horizon ($r \to r_s$), $f(r) \to 0$, so $V_{\text{eff}} \to 0$. At spatial infinity ($r \to \infty$), $V_{\text{eff}} \to 0$. The potential reaches a maximum at the *photon sphere*, $r = 1.5 r_s$, where light can orbit the black hole in an unstable circular orbit.

6 Physical Implications

The modified Maxwell equations derived in Section 4 have profound physical consequences.

6.1 The Unruh Effect

Although the classical equations (18) and (19) show modifications due to proper acceleration a, the quantum mechanical vacuum state is also observer-dependent. When the electromagnetic field is quantized in the static observer's frame, the observer perceives the Minkowski vacuum as a thermal bath of photons. The temperature of this Unruh radiation is directly proportional to the proper acceleration:

$$T_U = \frac{\hbar a}{2\pi c k_B} = \frac{\hbar}{2\pi c k_B} \frac{GM}{r^2\sqrt{1-\frac{r_s}{r}}}. \tag{31}$$

6.2 Connection to Hawking Radiation

As the static observer is lowered closer to the event horizon ($r \to r_s$), their proper acceleration a diverges. Consequently, the Unruh temperature diverges. In the context of quantum field theory in curved spacetime, this divergence is closely related to the Hawking temperature of the black hole:

$$T_H = \frac{\hbar c^3}{8\pi G M k_B}. \tag{32}$$

The equivalence principle ensures that local physics near the horizon is indistinguishable from flat-spacetime physics in a uniformly accelerating frame. This forms the mathematical basis for the derivation of Hawking radiation via the gravitational anomaly method or Euclidean path integral methods.

7 Conclusion

We have presented a comprehensive analysis of classical electrodynamics as measured by a uniformly accelerating (static) observer in Schwarzschild spacetime. By utilizing the tetrad formalism, we derived the 3+1 decomposed Maxwell's equations, revealing that both the observer's proper acceleration and the background spatial curvature introduce effective source terms. Furthermore, we derived the electromagnetic wave equation and analyzed the effective potential governing the propagation of light. The results demonstrate the deep interplay between acceleration, gravity, and electromagnetism, providing a robust framework for studying quantum field theory in curved spacetime, particularly the Unruh effect and Hawking radiation.

A Detailed Christoffel Symbols

For completeness, we list all non-zero Christoffel symbols of the second kind for the Schwarzschild metric. Using the definition $\Gamma^{\mu}_{\nu\lambda} = \frac{1}{2}g^{\mu\rho}(\partial_\nu g_{\rho\lambda} + \partial_\lambda g_{\rho\nu} - \partial_\rho g_{\nu\lambda})$, we find:

$$\Gamma^t_{tr} = \Gamma^t_{rt} = \frac{GM}{r^2 f}, \tag{33}$$

$$\Gamma^r_{tt} = \frac{GMf}{r^2}, \quad \Gamma^r_{rr} = -\frac{GM}{r^2 f}, \tag{34}$$

$$\Gamma^r_{\theta\theta} = -rf, \quad \Gamma^r_{\phi\phi} = -rf\sin^2\theta, \tag{35}$$

$$\Gamma^\theta_{r\theta} = \Gamma^\theta_{\theta r} = \frac{1}{r}, \quad \Gamma^\theta_{\phi\phi} = -\sin\theta\cos\theta, \tag{36}$$

$$\Gamma^\phi_{r\phi} = \Gamma^\phi_{\phi r} = \frac{1}{r}, \quad \Gamma^\phi_{\theta\phi} = \Gamma^\phi_{\phi\theta} = \cot\theta. \tag{37}$$

B Components of the Electromagnetic Tensor

The electromagnetic field tensor $F_{\mu\nu}$ in Schwarzschild coordinates can be related to the local fields **E** and **B** via the tetrad:

$$F_{tr} = \sqrt{f} E_{\hat{r}}, \tag{38}$$

$$F_{t\theta} = r\sqrt{f} E_{\hat{\theta}}, \tag{39}$$

$$F_{t\phi} = r \sin\theta \sqrt{f} E_{\hat{\phi}}, \tag{40}$$

$$F_{r\theta} = r B_{\hat{\phi}}, \tag{41}$$

$$F_{r\phi} = -r \sin\theta B_{\hat{\theta}}, \tag{42}$$

$$F_{\theta\phi} = r^2 \sin\theta B_{\hat{r}}. \tag{43}$$

References

[1] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, San Francisco, 1973).

[2] R. M. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984).

[3] S. W. Hawking, "Particle creation by black holes," Commun. Math. Phys. **43**, 199 (1975).

[4] W. G. Unruh, "Notes on black-hole evaporation," Phys. Rev. D **14**, 870 (1976).

[5] W. Rindler, "Kruskal space and the uniformly accelerated frame," Am. J. Phys. **34**, 1174 (1966).

[6] S. M. Carroll, *Spacetime and Geometry: An Introduction to General Relativity* (Addison-Wesley, San Francisco, 2004).

[7] T. W. Baumgarte and S. L. Shapiro, *Numerical Relativity: Solving Einstein's Equations on the Computer* (Cambridge University Press, Cambridge, 2010).